PROTECTION OF IMIDE GROUP OF URACIL MOIETY BY MEANS OF 2,2,2-TRICHLORO-TERT-BUTYLOXYCARBONYL CHLORIDE: A SELECTIVE SYNTHESIS OF 2'-O-METHYLURIDINE

Takashi KAMIMURA, Tsukio MASEGI, and Tsujiaki HATA\* Department of Life Chemistry, Tokyo Institute of Technology, Nagatsuta, Midoriku, Yokohama 227

Summary: 2,2,2-Trichloro-tert-butyloxycarbonyl group (TCBOC) was useful protecting group for the protection of imide function of uracil moiety. Starting from N<sup>3</sup>-TCBOC uridine derivative, 2'-0-methyluridine was selectively synthesized without formation of  $N^3$ -methyluridine derivatives.

There have been several protecting groups for amino and hydroxyl functions in nucleoside chemistry. However, little attention<sup>1)</sup> has been paid to imide group of nucleoside bases. Recently, Reese<sup>2)</sup> and our laboratory<sup>3)</sup> have demonstrated some of side reactions which may proceed during the coupling reactions for the synthesis of oligonucleotides. Synthesis of 2'-O-methyluridine exists in the same situation. When the methylation reaction of appropriately protected uridine was performed, methyl groups were introduced not only at the 2'-Ohydroxyl group but also the  $N^3$ -imide function.<sup>4)</sup> In order to avoid such side reactions, we have to tackle to find out an appropriate protecting group for these groups of nucleoside bases.

After several screenings 2,2,2-trichloro-t-butyloxycarbonyl group (TCBOC) was found to be the most suitable protecting group for the imide function of uracil residue. This group can be introduced in high yield on  $N^3$ -position of uridine by means of 2,2,2-trichloro-t-butyloxycarbonyl chloride (TCBOC-Cl) which was used as a reagent for protection of amino group of amino acids in peptide synthesis<sup>5)</sup> and is removed by treatment with zinc powder under very mild conditions.

The outline for the synthesis of 2'-O-methyluridine is shown in Scheme 1.

Procedure of each step is described as follows: When uridine (0.63 g, 2.57 mmol) was treated with 1,3-dichloro-1,1,3,3-tetraisopropyldisiloxane (TIPDSiCl<sub>2</sub>) (0.85 g, 2.70 mmole) in dry pyridine (20 ml) at room temperature overnight, the corresponding 3',5'-cyclic silylether derivative of uridine (1)<sup>6)</sup> was obtained in 95% yield (1.19 g).



Scheme I

Compound 1 (0.88 g, 1.81 mmole) was further treated with phenoxyacetic anhydride (1.55 g, 5.42 mmole)<sup>7)</sup> in dry pyridine (10 ml) at 0°C. 2'-O-Phenoxyacetyl-3',5'cyclic silylether derivative (2) was obtained in 87% yield (0.98 g) as a white powder.

<sup>1</sup>H-NMR(CDCl<sub>3</sub>)  $\delta$ =1.04, 1.08, 1.10(28, si-CH(CH<sub>3</sub>)<sub>2</sub>), 3.86-4.42(4, H-3',4',5'), 4.72  $(s, 2, C(0)-CH_2-OPh), 5.48(d, 1, J_{2'-3'}=4Hz, H-2'), 5.66(d, 1, J_{5-6}=8Hz, H-5),$ 5.76(s, 1, H-1'), 6.82-7.30(m, 5, O-Ph), 7.68(d, 1, H-6), 10.10(br.s, 1, H-3). Found: C, 53.09; H, 7.22; N, 4.74%. Calcd for C<sub>29</sub>H<sub>44</sub>O<sub>9</sub>N<sub>2</sub>Si<sub>2</sub>·2H<sub>2</sub>O: C, 53.03; H, 7.37; N, 4.26%.

Compound 2 (0.98 g, 1.58 mmole) was allowed to react with TCBOC-Cl (0.95 g, 3.94 mmole) in dry pyridine at room temperature overnight to afford the corresponding  $N^3$ -TCBOC uridine derivative (3) in 83% yield (1.07 g) as a white powder. <sup>1</sup>H-NMR(CDCl<sub>3</sub>)  $\delta$ =1.06, 1.10, 1.12(28, Si-CH(CH<sub>3</sub>)<sub>2</sub>), 2.10(s, 6, C(CH<sub>3</sub>)<sub>2</sub>CCl<sub>3</sub>), 3.88-4.44(4, H-3',4',5'), 4.70(s, 2, C(0)-CH<sub>2</sub>-OPh),  $5.50(d, 1, J_{2'-3'}=4Hz, H-2')$ , 5.68 (d, 2, J<sub>5-6</sub>=8Hz, H-5), 5.72(s, 1, H-1'), 6.82-7.30(m, 5, O-Ph), 7.66(d, 1, H-6). Found: C, 48.55; H, 6.13; N, 3.33; Cl, 12.48%. Calcd for C<sub>34</sub>H<sub>49</sub>O<sub>11</sub>N<sub>2</sub>Si<sub>2</sub>Cl<sub>3</sub>·H<sub>2</sub>O: C, 48.48; H, 6.10; N, 3.33; Cl, 12.63%.

The phenoxyacetyl group was selectively removed from the fully protected uridine derivative (3) (0.96 g, 1.16 mmole) by treatment with t-butylamine (0.255 g, or 0.366 ml, 3.49 mmole) in methanol (25 ml) at room temperature for 20 min. The

corresponding 2'-hydroxyl derivative (4) was obtained in 77% yield (0.614 g) as a white powder. <sup>1</sup>H-NMR(CDCl<sub>3</sub>)  $\delta$ =1.04, 1.08, 1.10(28, Si-CH(CH<sub>3</sub>)<sub>2</sub>), 2.08(s, 6, C(CH<sub>3</sub>)<sub>2</sub>CCl<sub>3</sub>), 3.12 (s, 1, OH-2'), 4.04-4.40(5, H-2',3',4',5'), 5.70(s, 1, H-1'), 5.72(d, 1, J<sub>5-6</sub>=8Hz, H-5), 7.68(d, 1, H-6). Found: C, 45.08; H, 6.36; N, 4.07; Cl, 16.06. Calcd for C<sub>26</sub>H<sub>43</sub>O<sub>9</sub>N<sub>2</sub>Si<sub>2</sub>Cl<sub>3</sub>: C, 45.25; H, 6.28; N, 4.06; Cl, 15.41%.

Compound 4 (0.5 g, 0.724 mmol) was methylated by use of excess methyl iodide (15.06 g, 0.106 mole) in the presence of silver oxide<sup>4a)</sup> (0.912 g, 3.93 mmole) under reflux for 7 h. 2'-O-Methylated product (5) was obtained in 93% yield (0.472 g) as a white powder.

<sup>1</sup>H-NMR(CDCl<sub>3</sub>)  $\delta$ =1.08, 1.12(28, Si-CH(CH<sub>3</sub>)<sub>2</sub>), 2.10(s, 6, C(CH<sub>3</sub>)<sub>2</sub>CCl<sub>3</sub>), 3.64(s, 3, 2'-OCH<sub>3</sub>), 3.74(d, 1, J<sub>2'-3</sub>=4Hz, H-2'), 4.04-4.32(4, H-3',4'5'), 5.68(d, 1, J<sub>5-6</sub>=8Hz, H-5), 5.72(s, 1, H-1'), 7.90(d, 1, H-6). Found: C, 46.21; H, 6.42; N, 3.94; Cl, 14.51%. Calcd for C<sub>27</sub>H<sub>45</sub>O<sub>9</sub>N<sub>2</sub>Si<sub>2</sub>Cl<sub>3</sub>: C, 46.05; H, 6.44; N, 3.98; Cl, 15.10%.

The TCBOC group was removed by treatment of 5 (0.408 g, 0.579 mmole) with zinc powder (0.568 g, 8.69 mmole) in the presence of acetylacetone<sup>8)</sup> (0.87 g, 8.69 mmole) in dry pyridine (15 ml) at room temperature for 15 min. The deblocked material (6) was obtained in 86% yield (0.25 g) as a white powder. <sup>1</sup>H-NMR(CDCl<sub>3</sub>)  $\delta$ =1.08, 1.12(28, Si-CH(CH<sub>3</sub>)<sub>2</sub>, 3.66(s, 3, 2'-OCH<sub>3</sub>), 3.72(d, 1, J<sub>2'-3</sub>' = 4Hz, H-2'), 4.02-4.30(4, H-3',4',5'), 5.64(d, 1, J<sub>5-6</sub>=8Hz, H-5), 5.72(s, 1, H-1'), 7.88(d, 1, H-6), 10.10(br.s, 1, H-3). Found: C, 52.85; H, 8.34; N, 5.44%. Calcd for C<sub>22</sub>H<sub>40</sub>O<sub>7</sub>N<sub>2</sub>Si<sub>2</sub>: C, 52.77; H, 8.05; N, 5.59%.

Desilylation from 6 (93 mg, 0.186 mmole) was performed by use of potassium fluoride (62.5 mg, 1.12 mmole) and tetraethylammonium chloride (234 mg, 1.12 mmole) in aqueous acetonitrile(67:1 v/v, 4.06 ml) at 53°C for 0.5 h<sup>9)</sup>. 2'-O-Methyluridine (7) was obtained in 69% yield which was estimated by UV absorption.  $(\xi_{263}=10,100^{4b})$  in 95% EtOH).

<sup>1</sup>H-NMR( $D_2O/t$ -BuOH internal)  $\delta$  =3.60(s, 3, 2'-OCH<sub>3</sub>), 3.72-4.44(5, H-2',3',4',5'), 5.94(d, 1,  $J_{5-6}$ =8Hz, H-5), 6.00(d, 1,  $J_{1'-2'}$ =4Hz, H-1'), 7.96(d, 1, H-6).

Quite recently, Reese<sup>10)</sup> has reported the protection of imide and amide group with aryl group. The application of TCBOC-protected uridine unit to the oligoribonucleotide synthesis is now in progress.

```
968
```

References

- J. F. W. McOmie, Ed. "Protective Groups in Organic Chemistry", Plenum Press, London, (1973), p. 403.
  - M. Miyaki, A. Saito, and B. Shimizu, Chem. Pharm. Bull., 18, 2459 (1970).

T. Azuma and K. Isono, Chem. Pharm. Bull., 25, 3347 (1977).

- J. A. Montgomery and H. J. Thomas, J. Org. Chem., <u>46</u>, 594 (1981).
- 2) P. K. Bridson, W. T. Markiewicz, and C. B. Reese, J. Chem. Soc. Chem. Commun., <u>1977</u>, 791.
  C. B. Reese and A. Ubasawa, Nucleic Acids Res. Symposium Series No. 7, 5 (1980).
  - C. B. Reese and A. Ubasawa, Tetrahedron Lett., 21, 2265 (1980).
- B. Rayner, C. B. Reese, and A. Ubasawa, J. Chem. Soc. Chem. Commun., 1980, 972.
- 3) H. P. Daskalov, M. Sekine, and T. Hata, Tetrahedron Lett., <u>21</u>, 3899 (1980). Idem, Bull. Chem. Soc. Jpn., <u>54</u>, 3076 (1981).
- 4) a) Y. Furukawa, K. Kobayashi, Y. Kanai, and M. Honjo, Chem. Pharm. Bull., <u>13</u>, 1273 (1965).
  - b) D. M. G. Martin, C. B. Reese, and G. F. Stephenson, Biochemistry, <u>7</u>, 1406 (1968).
  - c) J. T. Kusmierek, J. Giziewicz, and D. Shugar, Biochemistry, 12, 194 (1973).
  - d) A. H. Hains, Tetrahedron, 29, 2807 (1973).
  - e) M. J. Robins, S. R. Naik, and A. S. K. Lee, J. Org. Chem., 39, 1891 (1974).
- 5) H. Eckert, M. Listl, and I. Ugi, Angew. Chem. Int. Ed. Engl., <u>17</u>, 361 (1978).
- W. T. Markiewicz and M. Wiewiórowski, Nucleic Acids Res., Spec. publ. No. 4, s 185 (1978).
  - W. T. Markiewicz, J. Chem. Res. (s), 24 (1979).
- 7) C. B. Reese and J. C. M. Stewart, Tetrahedron Lett., 1968, 4273.
- 8) R. W. Adamiak, E.B. K. Grześkowiak, R. Kierzek, A. Kraszewski, W. T. Markiewicz, J. Stawiński, and M. Wiewiórowski, Nucleic Acids Res., <u>4</u>, 2321 (1977).
- 9) S. Honda, K. Terada, Y. Sato, M. Sekine, and T. Hata, Chem. Lett., 1982, 15.
- 10) S. S. Jones, C. B. Reese, S. Sibanda, and A. Ubasawa, Tetrahedron Lett., <u>22</u>, 4755 (1981).

(Received April 10, 1982)