

A CONVENIENT SYNTHESIS OF PHENYL SELENOCYANATE



Shuji TOMODA,* Yoshito TAKEUCHI, and Yujiro NOMURA

Department of Chemistry, College of General Education
The University of Tokyo, Komaba, Meguro-Ku, Tokyo 153

Treatment of benzeneselenenyl chloride with trimethylsilyl cyanide in various organic solvents provided phenyl selenocyanate in virtually quantitative yield.


The reaction presents a very simple preparation of the useful selenenylation agent.

Organic synthesis using organoselenium compounds is of current interest because of the facilities of transformations of these compounds to effect selective functional group interconversion.¹⁾ Among them, phenyl selenocyanate (1) occupies a unique position; it is used successfully for selective conversion of primary²⁾ and secondary^{2),3)} alcohols into corresponding selenides via S_N2 mechanism³⁾ and aldehydes⁴⁾ and carboxylic acids⁵⁾ are converted into cyanoselenides and selenocarboxylate in high yields, respectively (Eq. 1-3). More recently, a high-yield copper-catalyzed oxyselenenylation of olefins using 1 has been reported.⁶⁾



Current method for the preparation of 1 employs diazo reaction.⁷⁾ This requires two consecutive steps and tedious purification procedure (ethereal extraction and fractional distillation) which presents a nuisance not only because of unpleasant odor of 1⁸⁾ but also because of the fatal low yield of the reaction. Herein we report a very simple synthesis of 1, which provides 1 in virtually quantitative yield.

As shown in eq. 4, treatment of benzeneselenenyl chloride⁹⁾ with 20% excess trimethylsilyl cyanide¹⁰⁾ in dry tetrahydrofuran or in dry dichloromethane at room temperature for 5 to 10 min afforded a 96% yield of 1 after evaporation of solvent and the trimethylsilyl chloride produced.¹¹⁾ The



phenyl selenocyanate thus obtained as a light yellow oil¹²⁾ is sufficiently pure for most practical purposes as shown by thin layer chromatographic analysis.¹³⁾ This new method is particularly attrac-

tive for small-scale laboratory operations. Experimental details are as following:

Synthesis of Phenyl Selenocyanate: A solution of benzeneselenenyl chloride(212 mg, 1.11 mmol)⁹⁾ in dry THF(distilled from sodium diphenylketyl under argon, 2 ml) was added dropwise over 3 min to a solution of trimethylsilyl cyanide(131 mg, 1.33 mmol)¹⁰⁾ in dry THF(2 ml) with stirring at room temperature under argon. After addition, the mixture was stirred for 10 min and the system was evacuated to remove all volatile components, leaving 194 mg of practically pure phenyl selenocyanate (1) as a light yellow oil(96% yield); bp 104 °C/ 4 torr(lit⁷⁾) bp 118 °C/11 torr); IR(liquid film) 3312(w), 2120(m, CN), 1564 (m), 1470(s), 1431(s) cm⁻¹; ¹H-NMR(CDCl₃) δ=7.39(multiplet, 5H); ¹³C-NMR(CDCl₃) δ=132.7(d), 130.4(d), 129.7(d), 121.8(s), 101.4(s). If analytically pure 1 is desired, silica-gel column chromatography eluting with a hexane-dichloromethane(2:1) mixture or simple distillation under reduced pressure using Kugel-rohr is recommended.

References

- 1) For recent reviews on organoselenium chemistry, see; a) D.L.J. Clive, *Tetrahedron*, **34**, 1049(1978); b) H.J. Reich, *Acc. Chem. Res.*, **12**, 22(1979); c) H.J. Reich, "Oxidation in Organic Chemistry, Pt. C", W. Trahanovsky, Ed., Academic Press, New York, 1978, p.1.
- 2) P.A. Grieco, S. Gilman, and M. Nishizawa, *J. Org. Chem.*, **41**, 1485(1976).
- 3) M. Sevrin and A. Krief, *J. Chem. Soc. Chem. Commun.*, **1980**, 656.
- 4) P.A. Grieco and Y. Yokoyama, *J. Am. Chem. Soc.*, **99**, 5210(1977).
- 5) P.A. Grieco, Y. Yokoyama, and E. Williams, *J. Org. Chem.*, **43**, 1283(1978).
- 6) A. Toshimitsu, T. Aoai, S. Uemura, and M. Okano, *J. Org. Chem.*, **45**, 1953(1980).
- 7) O. Behaghel and H. Seibert, *Ber.*, **65**, 812(1932).
- 8) P.A. Grieco, J.Y. Jaw, D.A. Claremon, and K.C. Nicolaou, *J. Org. Chem.*, **46**, 1215(1981).
- 9) Purchased from Aldrich Chem. Co., New Jersey, U.S.A.
- 10) Prepared from trimethylsilyl chloride and silver cyanide; D.A. Evans, G.L. Carrol, and L.K. Truesdal, *J. Org. Chem.*, **39**, 914(1974). This reagent is commercially available from Aldrich Chem. Co., New Jersey, U.S.A. More economical syntheses of this reagent appeared recently in; a) J.K. Rasmussen and S.M. Heilman, *Synthesis*, **1979**, 523; b) S. Hünig and G. Wehner, *ibid*, **1979**, 522.
- 11) The reaction in acetonitrile at room temperature was extremely fast and afforded slightly less clean product mixture. In this case the temperature had to be lowered to -20 °C in order to minimize side reactions.
- 12) Compound 1 was identified by comparison of spectral data(IR and NMR) and physical constants(bp and TLC) with those of an authentic sample prepared according to reference 7.
- 13) The only minor impurity(less than 4%) was identified as diphenyl diselenide by TLC and IR spectral comparison with an authentic sample.

(Received June 9, 1981)