Article

# Methylenespiro[2.3]hexanes via Nickel-Catalyzed Cyclopropanations with [1.1.1]Propellane

Songjie Yu, Adam Noble, Robin B. Bedford, and Varinder K. Aggarwal J. Am. Chem. Soc., Just Accepted Manuscript • Publication Date (Web): 30 Nov 2019 Downloaded from pubs.acs.org on November 30, 2019

## **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

8 9

10 11

12 13

14 15 16

17 18

19

20

21

22

23

24

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

# Methylenespiro[2.3]hexanes via Nickel-Catalyzed Cyclopropanations with [1.1.1]Propellane

Songjie Yu, Adam Noble, Robin B. Bedford,\* and Varinder K. Aggarwal\*

School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, U.K.

Supporting Information Placeholder

**ABSTRACT:** [1.1.1]Propellane is a highly strained tricyclic hydrocarbon whose reactivity is dominated by addition reactions across the central inverted bond to provide bicyclo[1.1.1]pentane derivatives. These reactions proceed under both radical and twoelectron pathways, hence providing access to a diverse array of products. Conversely, transition metal-catalyzed reactions of [1.1.1]propellane are underdeveloped and lack synthetic utility, with reported examples generally yielding mixtures of ring-opened structural isomers, dimers, and trimers, often with poor selectivity. Herein, we report that nickel(0) catalysis enables the use of [1.1.1]propellane as a carbene precursor in cyclopropanations of a range of functionalized alkenes to give methylenespiro[2.3]hexane products. Computational studies provide support for initial formation of a Ni(0)-[1.1.1]propellane complex followed by concerted double C–C bond activation to give the key 3-methylenecyclobutylidene-nickel intermediate.

#### 1. Introduction

Catalytic cyclopropanation is a powerful method for generating three-membered carbocycles from alkenes and appropriate carbene precursors.<sup>1</sup> Key to the catalytic process is the formation of a transition metal carbene intermediate (Scheme 1), which then reacts with an alkene in either a concerted manner (pathway A),<sup>2,3</sup> wherein the alkene attacks the carbene moiety directly to form two C–C bonds simultaneously, or in a stepwise fashion via the formation of a metallacyclobutane intermediate (pathway B).<sup>4–6</sup> Irrespective of the pathway adopted, the formation of a carbene complex is regarded as pivotal to the transformation. Typically, this is accomplished by taking advantage of highly reactive compounds, such as diazoalkanes,<sup>7</sup> containing weak C–X bonds that are easily cleaved by the metal catalyst.<sup>8</sup>





A less common approach to metal carbenes is via C–C bond activation,<sup>9</sup> which is challenging due to the inertness of C–C

bonds and their lack of interaction with transition metal catalysts. Typically, catalytic C-C bond activation protocols require the use directing groups<sup>10</sup> or activated strained ring systems, including cyclopropanes<sup>11,12</sup> or cyclobutanones,<sup>13</sup> and proceed via cleavage of a single C-C bond to generate a metallacycle intermediate. Conversely, access to metal carbene complexes requires a double C-C bond activation of cyclopropanes, via a formal retro-cyclopropanation. While the use of simple cyclopropanes as carbene precursors is rare,<sup>14</sup> systems that impart additional ring-strain as a crucial driving force have been exploited to promote the challenging double C-C bond activation.<sup>15,16</sup> For example, bicyclo[1.1.0]butanes have been employed as metal carbene precursors under rhodium17 and nickel catalysis.<sup>18</sup> In contrast, [1.1.1]propellane,<sup>19</sup> which is considerably more strained than bicyclo[1.1.1]butane, has found little application as a metal carbene precursor.

[1.1.1]Propellane (1) is a highly strained tricyclic hydrocarbon, with a calculated strain energy of between 98 and 113 kcal/mol.<sup>20</sup> This is attributed to the unusual 'inverted' central  $\sigma$ -bond between its bridgehead carbons. As a result, the reactivity of 1 has been dominated by strain-release, wherein additions across the bridgehead bond vield bicyclo[1.1.1]pentane derivatives either via single-electron<sup>21</sup> or two-electron pathways (Scheme 2a).<sup>22,23</sup> Despite the growing number of synthetic methods for accessing functionalized bicyclo[1.1.1]pentanes from 1, reports of transition metal-catalyzed reactions of 1 are rare. However, its high ring-strain should provide the required driving force to facilitate metal-catalyzed retro-cyclopropanation, in which double C–C bond activation furnishes a metal carbene species.

## Scheme 2. Transformations of [1.1.1]propellane

(a) Bicyclo[1.1.1]pentane formation for [1.1.1]propellane (well-established)



(b) Transition metal-catalyzed reactions of [1.1.1]propellane (rare)



(c) Nickel-catalyzed cyclopropanations with [1.1.1]propellane (this work)



To date, the limited reports of carbene formation from 1 have resulted in mixtures of isomerization and oligomerization products, thus lacking synthetic utility. Thermal ring-opening of 1 under flow conditions at 430 °C yields 1,2dimethylenecyclopropane,<sup>24</sup> a reaction subsequently shown to carbene occur via a free intermediate (methylenecyclobutylidene) with an activation energy of 39.6 kcal/mol.<sup>25</sup> Wiberg explored ring-opening reactions of **1** in the presence of various transition metal catalysts (Scheme 2b).<sup>26</sup> With Rh(I) or Pd(II) an almost instantaneous reaction occurred to give the ring-opened dimer 2, whereas Ag(I), Ir(I), Pt(II), and Pt(0) led to slower reactions to produce varying amounts of dimer 2, methylene cyclobutene (3), and various trimeric products. The authors proposed that these were formed via metal carbene intermediates 4. This was supported by a cyclopropanation reaction with methyl acrylate catalyzed by  $[Rh(CO)_2Cl]_2$ , which gave methylenespiro[2.3]hexane 5 as a 1:1.5 mixture with 2. However, formation of 5 required the use of methyl acrylate as the solvent, presumably to make cyclopropanation competitive with the rapid dimerization of the carbene intermediate.

We envisaged that the carbenes generated by metalmediated ring-opening of **1** could be exploited more generally in cyclopropanation reactions with alkenes. Key to achieving this would be to suppress undesired isomerization and oligomerization reactions of the metal carbene intermediate. Herein, we report that nickel catalysis efficiently promotes cyclopropanation reactions of alkenes using **1** as a carbene precursor (Scheme 2c). The reaction proceeds via double C–C bond activation of **1** to generate nickel carbene **6** and provides access to methylenespiro[2.3]hexanes **7** through cyclopropanation of a broad array of alkenes. Density functional theory (DFT) calculations provide insight into the key interaction between **1** and the Ni(0) catalyst and support the formation of the proposed 3-methylenecyclobutylidene nickel complex **6**.

#### 2. Results and Discussion

Initial investigations focused on cyclopropanation of 4-vinylbiphenyl (8a) with 1 in the presence of various transition metal catalysts (Table 1). In line with the report by Wiberg,<sup>26</sup> a Rh(I) catalyst resulted in rapid formation of the dimerization product bis(3-methylenecyclobutylidene) (2) with no observable cyclopropanation (entry 1). Similarly, Ir(I), Pd(0), and Cu(I) catalysts failed to promote cyclopropanation (entries 2-4). Pleasingly, switching to Ni(cod)<sub>2</sub> as the catalyst with the N-heterocyclic carbene (NHC) ligand SIMes (generated in situ from 1,3-bis(2,4,6trimethylphenyl)imidazolinium chloride [SIMes·HCl] and NaO'Bu) suppressed the dimerization pathway and resulted in formation of methylenespiro[2.3]hexane derivative 7a in 64% yield after 20 h at 50 °C (entry 5).<sup>27</sup> Further evaluation of the

#### Table 1. Optimization studies<sup>a</sup>

Ν		catalyst (5–10 mol SIMes•HCI ( <i>x</i> mol <sup>6</sup>	%)	
<b>1</b> (2.0 equiv)	+ Ph 8a (1.0 equiv)	base (20 mol%) toluene, 50 °C, 20 h		Ph 7a
entry	catalyst (mol%)	x	base	$3 (\%)^b$
1	Rh(cod) <sub>2</sub> Cl (5)	-	-	0
2	$Ir(cod)_2Cl(5)$	-	-	0
3	Pd(dba)3 (5)	-	-	0
4	Cu(MeCN)4PF6 (10)	_ c	-	0
5	Ni(cod) <sub>2</sub> (10)	12	NaO <sup>t</sup> Bu	64
6	Ni(cod) <sub>2</sub> (10)	12	LiOMe	81
7	Ni(cod) <sub>2</sub> (10)	12	LiO <sup>t</sup> Bu	67
8	Ni(cod) <sub>2</sub> (10)	12	KO'Bu	69
9	Ni(cod) <sub>2</sub> (10)	20	LiOMe	87 (76)
10	Ni(cod) <sub>2</sub> (10)	12	LiOMe	(89) <sup>d</sup>
11	Ni(cod) <sub>2</sub> (10)	12	LiOMe	$(65)^{d,e}$
12	Ni(cod) <sub>2</sub> (10)	-	-	$(75)^{d}$
13	-	12	LiOMe	0

<sup>*a*</sup> Reaction conditions: **8a** (0.20 mmol), **1** (0.7–0.9 M in Et<sub>2</sub>O, 0.40 mmol), catalyst (5–10 mol%), SIMes·HCl (0–20 mol%), base (20 mol%), and solvent (3 mL) at 50 °C for 20 h. <sup>*b*</sup> Yields were determined by <sup>1</sup>H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. Numbers in parenthesis are isolated yields of the purified product. <sup>*c*</sup> 2,2-Bis((4*S*)-4-isopropyloxazoline)propane (12 mol%) was used as the ligand. <sup>*d*</sup> With 4.0 equiv **1**. <sup>*e*</sup> Using IMes·HCl (12 mol%) in place of SIMes·HCl.

58 59

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32 33 34

35

36

42

53

54 55

56

57 58 59

60

base used to generate the NHC ligand revealed LiOMe to be optimal, affording 7a in 81% yield (entries 6-8). Using a higher ligand loading of 20 mol% resulted in marginal increase in yield (entry 9). However, a significant improvement was observed upon increasing the amount of 1 from 2 equiv to 4 equiv, which allowed isolation of 7a in 89% (entry 10). Changing the NHC ligand to 1,3-bis(2,4,6trimethylphenyl)imidazol-2-ylidene (IMes) resulted in a slightly decreased yield (entry 11). Interestingly, Ni(cod)<sub>2</sub> in the absence of NHC ligand was still an effective catalyst for cyclopropanation, providing 7a in a slightly reduced yield of 75% (entry 12). Finally, a control experiment demonstrated the importance of the Ni(0) catalyst as no product was observed in its absence (entry 13), which is in line with the high energy barrier to free carbene formation via thermal rearrangement of 1.25

With optimal conditions for the nickel-catalyzed cyclopropanation established, the scope of the reaction was assessed (Scheme 3). A wide range of styrene derivatives reacted efficiently to provide cyclopropanes 7a-7k in 47-89% vield, including those bearing electron-donating (7c) and electron-withdrawing substituents (7d-7g). It is notable that several reactive functional groups, including a pinacol boronic ester (7h), chloride (7i and 7j), and fluoride (7k), are tolerated under the reaction conditions, despite them being known to nickel-catalyzed transformations.<sup>28</sup> undergo Alkenes substituted with electron-rich aromatic groups, such as naphthyl and ferrocenyl gave the spirocyclic products 71 and 7m in 84% and 44% yield, respectively. Heteroaromatic moieties are ubiquitous in bioactive molecules but their functionalization can be challenging using transition metal catalysis due to the potentially strong ligation to metal centers. In this catalytic system, a broad range of heteroaryl substituted

## Scheme 3. Alkene scope<sup>a</sup>

alkenes were tolerated, such as pyridine (7n), quinoline (7o), indole (7p), furan (7q), and benzothiophene (7r), undergoing cyclopropanation in good to excellent yields. In addition to alkenes with aromatic substituents, conjugated 1,3-diene and 1,3-enyne substrates were also tolerated. These substrates displayed higher reactivity and reactions proceeded efficiently at room temperature, delivering alkenyl or alkynyl-substituted spirocyclic products 7t–7x in good yield and with complete regioselectivity for reaction at the terminal alkene. We also investigated the reaction of terminal alkynes with 1 (Scheme 4). Interestingly, 4-ethynylbiphenyl (9) reacted via a carbene/alkyne metathesis rather than the expected cyclopropanation,<sup>29</sup> giving bis(3-methylenecyclobutylidene) product 10 in 20% yield.

## Scheme 4. Reaction of terminal alkynes



The presence of a conjugated  $\pi$ -system on the alkene substrate was found to be crucial for productive cyclopropanation, as alkyl-substituted alkenes (e.g., allylbenzene) were unreactive under our optimized conditions. Unfortunately, the cyclopropanation reaction also failed for  $\alpha$ and  $\beta$ -substituted styrenes. Therefore, we sought alternative alkene substrates that would undergo cyclopropanation while providing a synthetic handle that would allow subsequent substitution reactions. Boronic esters are highly versatile



<sup>*a*</sup> Reaction conditions: alkene (0.20 mmol), propellane (0.7–0.9 M in Et<sub>2</sub>O, 4.0 equiv), Ni(cod)<sub>2</sub> (10 mol%), SIMes·HCl (12 mol%), LiOMe (20 mol%), and toluene (3 mL) at 50 °C for 20 h. Yield are of isolated products. <sup>*b*</sup> Reaction performed at room temperature.

moieties that can readily be transformed into a diverse array of functional groups.<sup>30</sup> Thus, we examined the nickel-catalyzed cyclopropanation reaction with alkenylboronic esters 11 (Scheme 5).<sup>31</sup> We were pleased to find that vinylboronic acid pinacol ester underwent productive reaction, providing borylated methylenespiro[2.3]hexane **12a** in good yield.<sup>32</sup> Furthermore, in contrast to the styrene derivatives, alkenylboronic esters with  $\beta$ -substitution were also suitable substrates. Various alkyl groups at the  $\beta$ -position were tolerated, providing products 12b-12f in moderate to good yields and with excellent diastereoselectivity when E-alkene substrates were used. The stereospecificity of the reaction was examined by using (Z)-1-propenylboronic acid pinacol ester, which selectively provided the syn diastereomer of 12b but only with a modest 32:68 trans/cis ratio (see below for further discussion). In addition to  $\beta$ -substitution,  $\alpha$ -substituted alkenylboronic esters were efficiently cyclopropanated, providing tertiary benzylic (12g) and tertiary non-benzylic (12h and 12i) cyclopropylboronic esters in good yields. Unfortunately, trisubstituted alkenes proved unreactive, with 1-cyclohexenylboronic ester pinacol ester failing to yield tricyclic product 12j. Finally, the use of benzyl acrylate provided cyclopropyl carboxylate ester 13 in 40% yield.

## Scheme 5. Alkenylboronic ester scope<sup>a</sup>



<sup>*a*</sup> Yields are of isolated products. The *trans:cis* ratio was determined by GC/MS. <sup>*b*</sup> Using (*E*)-1-propenylboronic acid pinacol ester. <sup>*c*</sup> Using (*Z*)-1-propenylboronic acid pinacol ester

We next sought to highlight the synthetic utility of the methylenespiro[2.3]hexane skeleton (Scheme 6). Methylenecyclobutanes have previously found application in natural product total synthesis due to the versatility of the exocyclic alkene.33 Thus, we used this moiety as a handle for diversification of cyclopropanation product 7a. Transformation into synthetically useful organoboronic ester products was achieved via cross metathesis<sup>34</sup> and

hydroboration,<sup>35</sup> providing alkenylboronic ester **14** and alkylboronic ester **15**, respectively. Spiro[2.3]hexan-5-one derivative **16** could be prepared in excellent yield via ozonolysis. Finally, spirocyclic propagations were performed to access different sized heterocycles, including a carboalumination/thioetherification<sup>36</sup> to form tetrahydrothiophene **17**, and a [2+2] cycloaddition with chlorosulfonyl isocyanate to give  $\beta$ -lactam **18**.<sup>33c</sup>

### Scheme 6. Products derivatizations



<sup>*a*</sup> Prop-1-en-1-ylboronic acid pinacol ester, Grubbs 2<sup>nd</sup> generation catalyst (5 mol%), CH<sub>2</sub>Cl<sub>2</sub>, 50 °C. <sup>*b*</sup> Products formed in 52:48–56:44 d.r. <sup>*c*</sup> CuCl (5 mol%), DTBM-segphos (6 mol%), KO'Bu (20 mol%), HBpin (1.2 equiv), toluene, r.t. <sup>*d*</sup> O<sub>3</sub>, MeOH/CH<sub>2</sub>Cl<sub>2</sub>, then PPh<sub>3</sub> (1.5 equiv). <sup>*e*</sup> Cp<sub>2</sub>ZrCl<sub>2</sub> (5 mol%), Et<sub>3</sub>Al (1.2 equiv), toluene, then sulfur (2.4 equiv). <sup>*f*</sup> Chlorosulfonyl isocyanate (1.0 equiv), Et<sub>2</sub>O, r.t.

## 3. Mechanistic Studies

Our investigations into the mechanism of this nickelcatalyzed cyclopropanation initially focused on elucidating if the reaction proceeded via a single-electron pathway. This was considered due to the propensity of the bridgehead bond of 1 to react with free radicals<sup>37</sup> and because nickel-catalyzed reactions commonly involve single-electron transfer processes.<sup>28</sup> Furthermore, Wiberg has previously shown that 1 undergoes cyclopropanation reactions with highly activated electron-deficient  $\pi$ -systems and proposed that they proceed via a radical mechanism.<sup>26,38</sup> In our nickel-catalyzed process, if a stepwise radical mechanism was operative, cyclopropanation of 1,2-disubstituted alkenes would be expected to proceed with poor stereocontrol, whereas two-electron pathways of a metal carbene intermediate (e.g., pathways A and B in Scheme 1) should be stereospecific. Therefore, we investigated the stereospecificity of the cyclopropanation reaction using both (E)- and (Z)- $\beta$ -deutero-4-vinylbiphenyl (19) (Schemes 7a and 7b). Reaction of (E)-19 (>99:1 E/Z) with 1 under the standard conditions afforded *trans-20* as the major product with a 95:5 ratio of isotopomers, while the reaction of (Z)-19 (>99:1 Z/E) gave cis-20 as the major product in a 97:3 ratio. The high stereoselectivity of these reactions support a stereoretentive cycloaddition of a nickel carbene rather than a radical pathway. We postulated that the small drop in selectivity in the reactions of (E)-19 and (Z)-19, and the poor diastereoselectivity observed with (Z)-11b (see product syn-12b, Scheme 5), could be caused by isomerization of the

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22 23

24

25

26

27

28

29

30 31 32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24 25

26

27

28 29

30

31

32

33

34

35

36

37

38 39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

starting alkene prior to cyclopropanation. Indeed, submitting (*E*)-**19** and (*Z*)-**11b** to the standard reactions conditions but in the absence of **1** resulted in substantial isomerization, giving a 50:50 ratio of isomers of **19** and a 91:9 *E/Z* ratio for **11b** (Schemes 7c and 7d). In addition, when **8a** was treated with the nickel catalyst in the presence of methanol- $d_4$ , approximately 5% deuterium incorporation was observed in the recovered alkene (Scheme 7e), which reveals the alkene isomerization stems from a nickel-hydride species generated from oxidative addition of Ni(0) to methanol.<sup>39</sup> Based on these experiments, we concluded that the cyclopropanation proceeds via a two-electron pathway involving reaction of the alkene substrate with a nickel carbene intermediate, as shown in Scheme 1.

#### Scheme 7. Mechanistic experiments



While metal carbenes have been proposed as intermediates in transition metal-catalyzed reactions of [1.1.1]propellane,<sup>26</sup> their mechanism of formation and subsequent reactivity has not been studied. Therefore, we used DFT (B3LYP-D3/6-311++G\*\*//B3LYP-D3/6-31+G\*; SMD, solvent = toluene, see Supporting Information for full details) to model the nickel catalytic cycle,<sup>40</sup> where the smaller NHC ligand L was used as a representative catalyst (Figure 1).<sup>41,42</sup> Loss of a styrene from the 16-electron Ni(0) species **RS**, generating the 14-electron intermediate **I**<sup>A</sup>, is an endergonic process, consistent with **RS** acting as an off-cycle resting state. With intermediate **I**<sup>A</sup> + **1** + styrene defining the free-energy zero point of the cycle, coordination of propellane to **I**<sup>A</sup> to give the Ni(0) intermediate **I**<sup>B</sup> is thermodynamically favorable, with a  $\Delta G$  of -8.3 kcal/mol. An alternative diradical ground state for the adduct  $\mathbf{I}^{\mathbf{B}}$  ( $\mathbf{I}^{\mathbf{B}^*}$ ) was explored but was found to be too high in energy to be relevant (see Supporting Information for full details). The propellane adduct I<sup>B</sup> then undergoes a concerted ring-opening where two C-C bonds are cleaved to generate the carbene intermediate  $I^{C}$  via transition state  $TS^{BC}$ . The subsequent cyclopropanation was found to proceed through a stepwise pathway (see pathway B, Scheme 1), with C-C bond TS<sup>CD</sup>. formation, via transition state giving metallacyclobutane intermediate I<sup>D</sup>. The regioselectivity of the formation of metallacyclobutane  $I^D$  favored C-Ni bondformation at the benzylic carbon, as the alternative transition state TS<sup>CD'</sup>, in which C-C bond-formation occurs at the benzylic carbon, is 3.5 kcal/mol higher in free energy than transition state TS<sup>CD</sup>. Reductive elimination of the cyclopropane moiety, via  $TS^{DE}$ , then yields  $I^{E}$ , the Ni(0) adduct of the product, 7b, which is in turn liberated by reaction with styrene, regenerating  $I^{A,43}$  It appears that  $\pi$ coordination of the arene during the reductive elimination facilitates the process, a finding in accord with the experimental observation that replacing the arene with a more strongly coordinating alkene function allows the reaction temperature to be lowered (Scheme 3), whereas no reaction occurs when the coordinating group is removed (e.g., alkylsubstituted alkenes).



Figure 1. DFT-calculated catalytic cycle and associated free energy level diagram. \*The individual ligand substitution steps ( $I^E$  to  $I^A$ ) were not modelled.

Public

Intermediate  $I^B$  can also undergo smooth, reversible oxidative addition to give an off-cycle metalla-[2.1.1]propellane intermediate  $I^F$ , a process calculated to be thermodynamically and kinetically competitive with the concerted ring-opening to form  $I^C$  (Figure 1). While two alternative pathways for cyclopropanation that proceed via  $I^F$ may be envisaged, these were found to be energetically less likely, at least in our model system (see Supporting information for full details).

The calculated intermediate  $I^{B}$  represents a rare example of a coordinated [1.1.1]propellane,<sup>44</sup> while intermediate  $I^{F}$  can formally be considered as a metalla-[2.1.1]propellane. To the best of our knowledge, while rare examples of transition metal-based metalla-[1.1.1]propellanes are known,<sup>45</sup> [2.1.1]analogues have not been reported. The nature of the 'inverted' C–C bonds of small ring propellanes has been the subject of much debate in the literature,<sup>20a,46</sup> with recent *ab initio* valence bond calculations by Wu and co-workers revealing that the inverted bond has predominant 'charge-shift' bonding character with little or no diradical contribution.<sup>47,48</sup> Accordingly, a closer examination of the electronic structures of intermediates  $I^{B}$  and  $I^{F}$  (geometry optimization at the B3LYP/6-311++G\*\* level of theory) was of interest.

The calculated C-C bond length of the inverted bond of the propellane ligand in I<sup>B</sup> of 1.600 Å is marginally longer than either that calculated for free propellane at the same level of theory (1.582 Å) or measured for a closely related derivative (1.579 Å),<sup>49</sup> but is significantly shorter than the bridgehead carbon distance in bicyclo[1.1.1]pentane (1.845 Å),<sup>50</sup> which is indicative of a C-C bonding interaction. The coordination of the propellane in  $I^B$  is offset from linear, with a calculated angle between the inverted C-C bond axis and the nickel center of 151°. Meanwhile, the calculated Ni-C bond length to the propellane ligand (2.042 Å) is longer than that of the Ni-C bond for the carbene ligand (1.935 Å). Figure 2 shows both the bonding molecular orbitals (MO) between the propellane ligand and a d-orbital on nickel - an orbital that also shows significant involvement of the NHC donor - and a summary of key topological data, calculated using the Atoms in Molecules (AIM) method,<sup>51</sup> along with Wiberg Bond Indices (WBI) for key bonds in I<sup>B</sup> and propellane. Comparing the data, it is clear that the inverted C-C bond of propellane remains largely intact on coordination. The positive value for the Laplacian,  $\nabla^2 \rho$ , is atypical of covalent C–C bonds, which tend to return negative values, but is in line with the charge-shift description of propellane inverted bonds.49

The intermediate  $\mathbf{I}^{\mathrm{F}}$  could be viewed as containing either a metalla-[2.1.1]propellane (form  $\mathbf{A}$  in Figure 3) or, at the other extreme, a nickel carbene complex with an intramolecularly coordinated alkene moiety (form  $\mathbf{B}$ ). Indeed, as described in the introduction, metal carbene intermediates are ubiquitous in catalytic cyclopropanations. However, the calculated Ni–C<sub>a</sub> bond length (1.968 Å) is too long for any significant carbenic character, indeed it is longer than the Ni–C(NHC) bond (1.926 Å), which is generally accepted to be a single bond with very little retro-donation. By comparison, structurally characterized Ni(II) carbene complexes without heteroatom functionality on the carbene carbon display Ni=C bond lengths in the range 1.844 – 1.906 Å,<sup>52</sup> while the Ni–C bond lengths of the very limited number of nickelacyclobutanes reported are 1.927 – 1.956 Å.<sup>53</sup> Meanwhile, despite being marginally shorter than a

typical C-C single bond, Cb-Cc is significantly longer than a double bond, even one coordinated to Ni, as highlighted by the calculated C-C bond length of 1.361 Å for the coordinated styrene ligand in I<sup>F</sup>. Taken together, these bond metrics favor the metalla-[2.1.1]propellane form A, which would be expected to have an inverted Ca-Cb bond. Indeed, the separation of these carbon atoms is very similar to that calculated for [2.1.1]propellane at the same level of theory (Figure 3). An NBO analysis shows a  $\sigma$ -symmetry bonding orbital located between Ca and Cb while the equivalent MO has further density associated with the metal-C<sub>c</sub> fragment. The WBI of the C<sub>a</sub>–C<sub>b</sub> bond clearly indicate a bonding interaction, however, an AIM analysis does not identify a bond critical point (bcp) between these two carbon atoms. This is perhaps not surprising in view of the extended nature of the MO, and the observation that organometallic ligands with such conjugation to metal centers do not always reveal bcp's.<sup>54</sup> In contrast, the delocalization index ( $\delta(AB)$ ) for this interaction, also obtained from the AIM calculation, is clearly indicative of significant bonding.



**Figure 2.** MO of the Ni-propellane  $\sigma$ -bond of intermediate  $I^B$  (isovalue = 0.05 (electron/bohr<sup>3</sup>)<sup>1/2</sup>); AIM-calculated bond critical points (bcp, small green spheres), associated electron density ( $\rho$ ) and Laplacian ( $\nabla^2 \rho$ ) values at the bcp's and Wiberg Bond Indices (WBI) of the bonds.



**Figure 3.** NBO and MO associated with the inverted C-C bond of intermediate  $I^F$  (isovalue = 0.05 (electron/bohr<sup>3</sup>)<sup>1/2</sup>); WBI and AIM-calculated metrics (as defined in Figure 2) as well as AIM-derived delocalisation index ( $\delta$ (AB)) values.

## 4. Conclusions

We have developed a new approach to methylenespiro[2.3]hexane derivatives via a nickel-catalyzed cyclopropanation of alkenes with [1.1.1]propellane. The

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

reaction involves a rare example of double C-C bond provide activation of [1.1.1]propellane to а methylenecyclobutylidene-metal intermediate. Nickel catalysis proved essential in preventing deleterious side reactions of the metal-carbene and enabled the successful application of a wide range of alkene substrates, including styrenes, 1,3-dienes, 1,3eynes, and alkenyl boronic esters. The formation of a nickel carbene was supported both experimentally, by the high stereospecificity of the cyclopropanation of 1,2-disubstituted alkenes; and computationally, with DFT calculations showing the carbene is formed via a concerted double C-C bond cleavage of a nickel-[1.1.1]propellane complex. Given that transformations of propellane are dominated by additions across the bridgehead  $\sigma$ -bond to form bicyclo[1.1.1]pentanes, this nickel-catalyzed process unveils a new reaction mode that will enhance the application of [1.1.1]propellane in the synthesis of complex molecules.

## ASSOCIATED CONTENT

#### Supporting Information.

Experimental procedures, full characterization data for new compounds, and details of computational studies are supplied in the Supporting Information. This material is available free of charge via the Internet at http://pubs.acs.org.

## AUTHOR INFORMATION

#### Corresponding Author

\* r.bedford@bristol.ac.uk

\* v.aggarwal@bristol.ac.uk

#### Notes

The authors declare no competing financial interests.

#### ACKNOWLEDGMENT

We thank the EPSRC (EP/S017801/1) for financial support. S.Y. thanks the EU for an H2020 Marie Skłodowska-Curie Fellowship (grant no. 792439).

## REFERENCES

(1) For selected reviews, see: (a) Wu, W.; Lin, Z.; Jiang, H. Recent Advances in the Synthesis of Cyclopropanes. Org. Biomol. Chem. 2018, 16, 7315. (b) Thankachan, A. P.; Sindhu, K. S.; Krishnan, K. K.; Anilkumar, G. Recent Advances in the Syntheses, Transformations and Applications of 1,1-Dihalocyclopropanes. Org. Biomol. Chem. 2015, 13, 8780. (c) Bos, M.; Poisson, T.; Pannecoucke, X.; Charette A. B.; Jubault, P. Recent Progress Toward the Synthesis of Trifluoromethyl- and Difluoromethyl-Substituted Cyclopropanes. Chem. Eur. J. 2017, 23, 4950. (d) Ebner, C.; Carreira, E. M. Cyclopropanation Strategies in Recent Total Syntheses. Chem. Rev. 2017, 117, 11651. (e) Tabolin A. A.; Ioffe, S. L. Small Ring Compounds and N-Oxides: Cycloadditions and Related Processes. Isr. J. Chem. 2016, 56, 385. (f) Trushkov, I. V. Indole-derived Donoracceptor Cyclopropanes. Isr. J. Chem. 2016, 56, 369. (g) Ganesh, V.; Sridhar, P. R.; Chandrasekaran, S. Synthetic Applications of Carbohydrate-derived Donor-Acceptor Cyclopropanes. Isr. J. Chem. 2016, 56, 417. (h) Reiser, O. Catalytic Conversion of Furans and Pyrroles to Natural Products and Analogues Utilizing Donor-Acceptor Substituted Cyclopropanes as Key Intermediates. Isr. J. Chem. 2016, 56, 531. (i) Candish, L.; Gillard, R. M.; Fernando, J. E. M.; Levens, A.; Lupton, D. W. All-Carbon N-Heterocyclic Carbene-Catalyzed (3+2) Annulation using Donor-Acceptor Cyclopropanes. Isr. J. Chem. 2016, 56, 522. (j) Qian D.; Zhang, J. Gold-Catalyzed Cyclopropanation Reactions Using a Carbenoid Precursor Toolbox. Chem. Soc. Rev. 2015, 44, 677.

(2) (a) Bonge, H. T.; Hansen, T. Computational Study of Cyclopropanation Reactions with Halodiazoacetates. *J. Org. Chem.* **2010**, *75*, 2309. (b) Hansen, J.; Autschbach, J.; Davies, H. M. L. Computational Study on the Selectivity of Donor/Acceptor-Substituted Rhodium Carbenoids. *J. Org. Chem.* **2009**, *74*, 6555. (c) Nowlan, D. T., III; Gregg, T. M.; Davies, H. M. L.; Singleton, D. A. Isotope Effects and the Nature of Selectivity in Rhodium-Catalyzed Cyclopropanations. *J. Am. Chem. Soc.* **2003**, *125*, 15902.

(3) (a) Suenobu, K.; Itagaki, M.; Nakamura, E. Reaction Pathway and Stereoselectivity of Asymmetric Synthesis of Chrysanthemate with the Aratani *C*<sub>1</sub>-Symmetric Salicylaldimine–Copper Catalyst. *J. Am. Chem. Soc.* **2004**, *126*, 7271. (b) Fraile, J. M.; García, J. I.; Gil, M. J.; Martínez-Merino, V.; Mayoral, J. A.; Salvatella, L. Theoretical Insights into the Role of a Counterion in Copper-Catalyzed Enantioselective Cyclopropanation Reactions. *Chem. Eur. J.* **2004**, *10*, 758. (b) Fraile, J. M.; García, J. I.; Martínez-Merino, V.; Mayoral, J. A.; Salvatella, L. Theoretical (DFT) Insights into the Mechanism of Copper-Catalyzed Cyclopropanation Reactions. Implications for Enantioselective Catalysis. *J. Am. Chem. Soc.* **2001**, *123*, 7616.

(4) (a) Berthon-Gelloz, G.; Marchant, M.; Straub, B. F.; Marko, I. E. Palladium-Catalyzed Cyclopropanation of Alkenyl Silanes by Diazoalkanes: Evidence for a Pd<sup>0</sup> Mechanism. *Chem. Eur. J.* **2009**, *15*, 2923. (b) Bernardi, F.; Bottoni, A.; Miscione, G. P. DFT Study of the Palladium-Catalyzed Cyclopropanation Reaction. *Organometallics* **2001**, *20*, 2751.

(5) Rosenberg, M. L.; Krapp, A.; Tilset, M. On the Mechanism of Cyclopropanation Reactions Catalyzed by a Rhodium(I) Catalyst Bearing a Chelating Imine-Functionalized NHC Ligand: A Computational Study. *Organometallics* **2011**, *30*, 6562.

(6) Wang, F.; Meng, Q.; Li, M. Density Functional Computations of the Cyclopropanation of Ethene Catalyzed by Iron (II) Carbene Complexes Cp(CO)(L)Fe=CHR, L = CO, PMe<sub>3</sub>, R = Me, OMe, ph, CO<sub>2</sub>Me. *Int. J. Quantum Chem.* **2008**, *108*, 945.

(7) (a) Doyle, M. P.; McKervey M. A.; Tao, Y. In Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides. Wiley-VCH: Weinheim, 1998, pp. 1–65. (b) Ford, A.; Miel, H.; Ring, A.; Slattery, C. N.; Maguire A. R.; McKervey, M. A. Modern Organic Synthesis with α-Diazocarbonyl Compounds. Chem. Rev. 2015, 115, 9981. (c) Mix, K. A.; Aronoff M. R.; Raines, R. T. Diazo Compounds: Versatile Tools for Chemical Biology. ACS Chem. Biol. 2016, 11, 3233.

(8) Jia, M.; Ma, S. New Approaches to the Synthesis of Metal Carbenes. *Angew. Chem. Int. Ed.* **2016**, *55*, 9134.

(9) (a) Jones, W. D. The Fall of the C–C Bond. *Nature* **1993**, *364*, 676. (b) Rybtchinski, B.; Milstein, D. Metal Insertion into C–C Bonds in Solution. *Angew. Chem., Int. Ed.* **1999**, *38*, 870. (c) Souillart, L.; Cramer, N. Catalytic C–C Bond Activations via Oxidative Addition to Transition Metals. *Chem. Rev.* **2015**, *115*, 9410. (d) Chen, P.-H.; Billett, B.; Tsukamoto, T.; Dong, G. "Cut and Sew" Transformations via Transition-Metal-Catalyzed Carbon–Carbon Bond Activation. *ACS Catal.* **2017**, *7*, 1340. (e) Fumagalli, G.; Stanton, S.; Bower, J. F. Recent Methodologies That Exploit C–C Single-Bond Cleavage of Strained Ring Systems by Transition Metal Complexes. *Chem. Rev.* **2017**, *117*, 9404. (f) Song, F.; Gou, T.; Wang, B.-Q.; Shi, Z.-J. Catalytic Activations of Unstrained C–C Bond involving Organometallic Intermediates. *Chem. Soc. Rev.* **2018**, *47*, 7078.

(10) (a) Rathbun, C. M.; Johnson, J. B. Rhodium-Catalyzed Acylation with Quinolinyl Ketones: Carbon–Carbon Single Bond Activation as the Turnover-Limiting Step of Catalysis. J. Am. Chem. Soc. 2011, 133, 2031. (b) Wentzel, M. T.; Reddy, V. J.; Hyster, T. K.; Douglas, C. J. Chemoselectivity in Catalytic C–C and C–H Bond Activation: Controlling Intermolecular Carboacylation and Hydroarylation of Alkenes. Angew. Chem., Int. Ed. 2009, 48, 6121. (c) Zeng, R.; Dong, G. Rh-Catalyzed Decarbonylative Coupling with Alkynes via C–C Activation of Isatins. J. Am. Chem. Soc. 2015, 137, 1408. (d) Shaw, M. H.; Melikhova, E. Y.; Kloer, D. P.; Whittingham, W. G.; Bower, J. F. Directing Group Enhanced Carbonylative Ring Expansions of Amino-Substituted Cyclopropanes: Rhodium-Catalyzed Multicomponent Synthesis of N-Heterobicyclic Enones. J. Am. Chem. Soc. 2013, 135, 4992. (e) Wang, G.-W.; McCreanor, N. G.; Shaw, M. H.; Whittingham, W. G.; Bower, J. F. New Initiation Modes for Directed Carbonylative C–C Bond Activation: Rhodium-Catalyzed (3+1+2) Cycloadditions of Aminomethylcyclopropanes. J. Am. Chem. Soc. 2016, 138, 13501.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

(11) (a) Liu, L.; Montgomery, J. Dimerization of Cyclopropyl Ketones and Crossed Reactions of Cyclopropyl Ketones with Enones as an Entry to Five-Membered Rings. J. Am. Chem. Soc. 2006, 128, 5348. (b) Tamaki, T.; Ohashi, M.; Ogoshi, S. [3+2] Cycloaddition Reaction of Cyclopropyl Ketones with Alkynes Catalyzed by Nickel/Dimethylaluminum Chloride. Angew. Chem., Int. Ed. 2011, 50, 12067. (c) Ogoshi, S.; Nagata, M.; Kurosawa, H. Formation of Nickeladihydropyran by Oxidative Addition of Cyclopropyl Ketone. Key Intermediate in Nickel-Catalyzed Cycloaddition. J. Am. Chem. Soc. 2006, 128, 5350. (d) Jiao, L.; Lin, M.; Yu, Z.-X. Density Functional Theory Study of the Mechanisms and Stereochemistry of the Rh(I)-Catalyzed Intramolecular [3+2] Cycloadditions of 1-Ene-and 1-Yne-Vinylcyclopropanes. J. Am. Chem. Soc. 2011, 133, 447.

(12) (a) Lautens, M.; Ren, Y. Transition Metal Catalyzed Stereospecific Intramolecular [3 + 2] Cycloadditions of Methylenecyclopropanes with Alkynes. J. Am. Chem. Soc. 1996, 118, 9597. (b) Saito, S.; Masuda, M.; Komagawa, S. Nickel-Catalyzed Intermolecular [3 + 2 + 2] Cocyclization of Ethyl Cyclopropylideneacetate and Alkynes. J. Am. Chem. Soc. 2004, 126, 10540. (c) Saya, L.; Bhargava, G.; Navarro, M. A.; Gulías, M.; López, F.; Fernández, I.; Castedo, L.; Mascareñas, J. L. Nickel-Catalyzed [3+2+2]Cycloadditions between Alkynylidenecyclopropanes and Activated Alkenes. Angew. Chem., Int. Ed. 2010, 49, 9886. (d) Evans, P. A.; Inglesby, P. A. Diastereoselective Rhodium-Catalyzed Ene-Cycloisomerization Reactions of Alkenylidenecyclopropanes: Total Synthesis of (-)-a-Kainic Acid. J. Am. Chem. Soc. 2012, 134, 3635. (e) Terao, J.; Tomita, M.; Singh, S. P.; Kambe, N. Nickel-Catalyzed Regioselective Carbomagnesation of Methylenecyclopropanes through a Site-Selective Carbon-Carbon Bond Cleavage. Angew. Chem., Int. Ed. 2010, 49, 144. (f) Simaan, S.; Marek, I. Hydroformylation Reaction of Alkylidenecyclopropane Derivatives: A New Pathway for the Formation of Acyclic Aldehydes Containing Quaternary Stereogenic Carbons. J. Am. Chem. Soc. 2010, 132, 4066.

(13) (a) Juliá-Hernández, F.; Ziadi, A.; Nishimura, A.; Martin, R. Nickel-Catalyzed Chemo-, Regio- and Diastereoselective Bond Formation through Proximal C–C Cleavage of Benzocyclobutenones. *Angew. Chem., Int. Ed.* 2015, 54, 9537. (b) Thakur, A.; Facer, M. E.; Louie, J. Nickel-Catalyzed Cycloaddition of 1,3-Dienes with 3-Azetidinones and 3-Oxetanones. *Angew. Chem., Int. Ed.* 2013, 52, 12161. (c) Masuda, Y.; Hasegawa, M.; Yamashita, M.; Nozaki, K.; Ishida, N.; Murakami, M. Oxidative Addition of a Strained C–C Bond onto Electron-Rich Rhodium(I) at Room Temperature. *J. Am. Chem. Soc.* 2013, *135*, 7142. (d) Souillart, L.; Parker, E.; Cramer, N. Highly Enantioselective Rhodium(I)-Catalyzed Activation of Enantiotopic Cyclobutanone C–C Bonds. *Angew. Chem., Int. Ed.* 2014, *53*, 3001. (e) Ko, H. M.; Dong, G. Cooperative Activation of Cyclobutanones and Olefins Leads to Bridged Ring Systems by a Catalytic [4 + 2] Coupling. *Nat. Chem.* 2014, *6*, 739.

45 (14) (a) Asako, S.; Kobashi, T.; Takai, K. Use of Cyclopropane as 46 C1 Synthetic Unit by Directed Retro-Cyclopropanation with Ethylene Release J. Am. Chem. Soc. 2018, 140, 15425. (b) Gassman, P. G.; 47 Johnson, T. H. Retrocarbene Additions. Dissection of Alkyl-48 Substituted Cyclopropanes Under Metathesis Conditions. J. Am. 49 Chem. Soc. 1976, 98, 6057. (c) Gassman, P. G.; Johnson, T. H. 50 Cyclopropane-Olefin Cross Metathesis. J. Am. Chem. Soc. 1976, 98, 6058. (d) Kawamura, T.; Kawaguchi, Y.; Sugikubo, K.; Inagaki, F.; 51 Mukai, C. Rhodium(I)-Catalyzed Cycloisomerization of Allene-52 Allenylcyclopropanes. Eur. J. Org. Chem. 2015, 2015, 719.

Anenyrcyclopropanes. Eur. J. Org. Chem. 2015, 2017, 719.
(15) Yoshizaki, S.; Shibata, Y.; Tanaka, K. Fulvene Synthesis by Rhodium(I)-Catalyzed [2+2+1] Cycloaddition: Synthesis and Catalytic Activity of Tunable Cyclopentadienyl Rhodium(III)
Complexes with Pendant Amides. Angew. Chem., Int. Ed. 2017, 56, 3590. (16) For retro-cyclopropanations driven by aromatization, see: (a) Solorio-Alvarado, C. R.; Echavarren, A. M. Gold-Catalyzed Annulation/Fragmentation: Formation of Free Gold Carbenes by Retro-Cyclopropanation. J. Am. Chem. Soc. **2010**, *132*, 11881. (b) Wang, Y.; McGonigal, P. R.; Herlé, B.; Besora, M.; Echavarren, A. M. Gold(I) Carbenes by Retro-Buchner Reaction: Generation and Fate. J. Am. Chem. Soc. **2014**, *136*, 801.

(17) (a) Walczak, M.; Krainz, T.; Wipf, P. Ring-Strain-Enabled Reaction Discovery: New Heterocycles from Bicyclo[1.1.0]butanes. *Acc. Chem. Res.* **2015**, *48*, 1149. (b) Walczak, M.; Wipf, P. Rhodium(I)-Catalyzed Cycloisomerizations of Bicyclobutanes. *J. Am. Chem. Soc.* **2008**, *130*, 6924.

(18) (a) Noyori, R.; Suzuki, T.; Kumagai, Y.; Takaya, H. Nickel(0)-Catalyzed Reaction of Bicyclo[1.1.0]butanes with Olefins. *J. Am. Chem. Soc.* **1971**, *93*, 5894. (b) Noyori, R. On the Nature of Carbenoids Generated from Bicyclo[1.1.0]butanes and Transition Metal Complexes. *Tetrahedron Lett.* **1973**, *14*, 1691. (c) Takaya, H.; Suzuki, T.; Kumagai, Y.; Hosoya, M.; Kawauchi, H.; Noyori, R. Nickel(0)-Catalyzed Reactions of Bicyclo[1.1.0]butanes. Geminal Two-Bond Cleavage Reaction and the Stereospecific Olefin Trapping of the Carbenoid Intermediate. *J. Org. Chem.* **1981**, *46*, 2854.

(19) (a) Wiberg, K. B.; Walker, F. H. [1.1.1]Propellane. J. Am. Chem. Soc. **1982**, 104, 5239. (b) Semmler, K.; Szeimies, G.; Belzner, J. Tetracyclo[ $5.1.0.0^{1.6}.0^{2.7}$ ]octane, a [1.1.1]Propellane Derivative, and a New Route to the Parent Hydrocarbon. J. Am. Chem. Soc. **1985**, 107, 6410.

(20) The calculated strain energy is dependent on the level of theory used. See: (a) Wiberg, K. B. Strain Energies of Small Ring Propellanes. J. Am. Chem. Soc. **1983**, 105, 1227. (b) Khoury, P. R.; Goddard, J. D.; Tam, W.; Ring Strain Energies: Substituted Rings, Norbornanes, Norbornenes and Norbornadienes. Tetrahedron **2004**, 60, 8112. (c) Polo, V.; Andres, J.; Silvi, B. New Insights on the Bridge Carbon–Carbon Bond in Propellanes: A Theoretical Study Based on the Analysis of the Electron Localization Function. J. Comput. Chem. **2007**, 28, 857. (d) Ioffe, A. I.; Svyatkin, V. A.; Nefedov, O. M. Molecular-Mechanical Analysis of the Structure of Strained Organic Molecules. 6. [m.n.k]-Propellanes. Bull. Acad. Sci. USSR Div. Chem. Sci. **1988**, 37, 1827.

(21) (a) Kaszynski, P.; Friedli, A. C.; Michl, J. Toward a Molecular-Size Tinkertoy Construction Set. Preparation of Terminally Functionalized [n]Staffanes from [1.1.1]Propellane. J. Am. Chem. Soc. 1992, 114, 601. (b) Kaszynski, P.; McMurdie, N. D.; Michl, J. Synthesis of Doubly Bridgehead Substituted Bicyclo[1.1.1]pentanes. Radical Transformations of Bridgehead Halides and Carboxylic Acids. J. Org. Chem. 1991, 56, 307. (c) Messner, M.; Kozhushkov, S. I.; de Meijere, A. Nickel- and Palladium-Catalyzed Cross-Coupling Reactions at the Bridgehead of Bicyclo[1.1.1]pentane Derivatives - A Convenient Access to Liquid Crystalline Compounds Containing Bicyclo[1.1.1]pentane Moieties. Eur. J. Org. Chem. 2000, 1137. (d) Kanazawa, J.; Maeda, K.; Uchiyama, M. Radical Multicomponent Carboamination of [1.1.1]Propellane. J. Am. Chem. Soc. 2017, 139, 17791. (e) Shelp, R. A.; Walsh, P. J. Synthesis of BCP Benzylamines From 2-Azaallyl Anions and [1.1.1]Propellane. Angew. Chem., Int. Ed. 2018, 57, 15857. (f) Caputo, D. F. G.; Arroniz, C.; Dürr, A. B.; Mousseau, J. J.; Stepan, A. F.; Mansfield, S. J.; Anderson, E. A. Synthesis and Applications of Highly Functionalized 1-Halo-3substituted Bicyclo[1.1.1]pentanes. Chem. Sci. 2018, 9, 5295. (g) Nugent, J.; Arroniz, C.; Shire, B. R.; Sterling, A. J.; Pickford, H. D.; Wong, M. L. J.; Mansfield, S. J.; Caputo, D. F. J.; Owen, B.; Mousseau, J. J.; Duarte, F.; Anderson, E. A. A General Route to Bicyclo[1.1.1]pentanes through Photoredox Catalysis. ACS Catal. 2019, 9, 9568. (h) Bär, R. M.; Kirschner, Nieger, S. M.; Bräse, S. Alkyl and Aryl Thiol Addition to [1.1.1]Propellane: Scope and Limitations of a Fast Conjugation Reaction. Chem. Eur. J. 2018, 24, 1373.

(22) (a) Gianatassio, R.; Lopchuk, J. M.; Wang, J.; Pan, C.-M.; Malins, L. R.; Prieto, L.; Brandt, T. A.; Collins, M. R.; Gallego, G. M.; Sach, N. W.; Spangler, J. E.; Zhu, H.; Zhu, J.; Baran, P. S. Strain-Release Amination. *Science* **2016**, *351*, 241. (b) Makarov, I. S.;

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

1 Brocklehurst, C. E.; Karaghiosoff, K.; Koch, G.; Knochel, P. 2 Synthesis of Bicyclo[1.1.1]pentane Bioisosteres of Internal Alkynes and para-Disubstituted Benzenes from [1.1.1]Propellane. Angew. 3 Chem., Int. Ed. 2017, 56, 12774. (c) Trongsiriwat, N.; Pu, Y.; Nieves-4 Quinones, Y.; Shelp, R. A.; Kozlowski, M. C.; Walsh, P. J. Reactions 5 of 2-Aryl-1,3-Dithianes and [1.1.1]Propellane. Angew. Chem. Int. Ed. 6 2019, 58, 13416. (d) Hughes, J. M. E.; Scarlata, D. A.; Chen, A. C.-7 Y.; Burch, J. D.; Gleason, J. L. Aminoalkylation of [1.1.1]Propellane Enables Direct Access to High Value 3-Alkylbicyclo[1.1.1]pentan-1-8 amines. Org. Lett. 2019, 21, 6800. 9 (23) For review articles, see: (a) Wiberg, K. Small Ring 10

Propellanes. Chem. Rev. 1989, 89, 975. (b) Levin, M. D.; Kaszynski,
P.; Michl, J. Bicyclo[1.1.1]pentanes, [n]Staffanes, [1.1.1]Propellanes, and Tricyclo[2.1.0.0<sup>2.5</sup>]pentanes. Chem. Rev. 2000, 100, 169. (c)
Dilmaç, A. M.; Spuling, E.; de Meijere, A.; Bräse, S. Propellanes—
From a Chemical Curiosity to "Explosive" Materials and Natural
Products. Angew. Chem., Int. Ed. 2017, 56, 5684. (d) Kanazawa, J.;
Uchiyama, M. Recent Advances in the Synthetic Chemistry of Bicyclo[1.1.1]pentane. Synlett 2019, 30, 1.

(24) Belzner, J.; Szeimies, G. Thermal Rearrangement of Some [1.1.1] Propellanes. *Tetrahedron Lett.* **1986**, *27*, 5839.

(25) (a) Pierini, A. B.; Reale, H. F.; Medrano, J. A. A Novel Interpretation of Bonding in [1.1.1] Propellane and Theoretical Calculation of its Unimolecular Rearrangement. *J. Mol. Struct.* (*THEOCHEM*) **1986**, *148*, 109. (b) Jarosch, O.; Walsh, R.; Szeimies, G. Kinetics and Mechanism of the Thermal Rearrangement of [1.1.1]Propellane. *J. Am. Chem. Soc.* **2000**, *122*, 8490.

(26) Wiberg, K.; Waddell, S. T. Reactions of [1.1.1]Propellane. J. Am. Chem. Soc. **1990**, *112*, 2194.

(27) In the absence of styrene, [1.1.1]propellane dimerized instantaneously in the presence of  $[Rh(CO)_2Cl]_2$  but <5% dimerization was observed after 20 h at 50 °C in the presence of Ni(cod)<sub>2</sub> (see Supporting Information for details).

(28) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. Recent Advances in Homogeneous Nickel Catalysis. *Nature* **2014**, *509*, 299.

(29) (a) Le Paih, J.; Vovard-Le Bray, C.; Dérien, S.; Dixneuf, P. H. Ruthenium-Catalyzed Synthesis of Functional Conjugated Dienes via Addition of Two Carbene Units to Alkynes. *J. Am. Chem. Soc.* **2010**, *132*, 7391. For a review, see: (b) Pei, C.; Zhang, C.; Qian, Y.; Xu, X. Catalytic Carbene/Alkyne Metathesis (CAM): A Versatile Strategy for Alkyne Bifunctionalization. *Org. Biomol. Chem.* **2018**, *16*, 8677.

(30) Sandford, C.; Aggarwal, V. K. Stereospecific Functionalizations and Transformations of Secondary and Tertiary Boronic Esters. *Chem. Commun.* **2017**, *53*, 5481.

(31) For selected examples of transition metal-catalyzed cyclopropanations of alkenylboronic esters, see: (a) Fontani, P.; Carboni, B.; Vaultier, M.; Carrié, R. A Convenient Highly Stereoselective Synthesis of Cycloproylboronates. *Tetrahedron Lett.* 1989, 30, 4815. (b) Carreras, J.; Caballero, A.; Pérez, P. J. Enantio-and Diastereoselective Cyclopropanation of 1-Alkenylboronates: Synthesis of 1-Boryl-2,3-Disubstituted Cyclopropanes. *Angew. Chem., Int. Ed.* 2018, *57*, 2334.

(32) For reactions with alkenylboronic esters, higher yields were obtained when NaO'Bu was used in place of LiOMe.

(33) (a) Tanino, K.; Takahashi, M.; Tomata, Y.; Tokura, H.; Uehara, T.; Narabu, T.; Miyashita, M. Total Synthesis of Solanoeclepin A. *Nat. Chem.* **2011**, *3*, 484. (b) Sun, M.; Li, W.-D. Z.; Qiu, F. G. Asymmetric Synthesis of the DEFG Rings of Solanoeclepin A. *Org. Lett.* **2019**, *21*, 644. (c) Palomo, C.; Oiarbide, M.; Bindi, S. A Concise  $\beta$ -Lactam Route to Short Peptide Segments Containing  $\beta$ , $\beta$ -Disubstituted  $\beta$ -Amino Acids. *J. Org. Chem.* **1998**, *63*, 2469.

(34) Morrill, C.; Grubbs, R. H. Synthesis of Functionalized Vinyl Boronates via Ruthenium-Catalyzed Olefin Cross-Metathesis and Subsequent Conversion to Vinyl Halides. *J. Org. Chem.* **2003**, *68*, 6031.

(35) Jang, W. J.; Song, S. M.; Moon, J. H.; Lee, J. Y.; Yun, J. Copper-Catalyzed Enantioselective Hydroboration of Unactivated 1,1-Disubstituted Alkenes. *J. Am. Chem. Soc.* **201**7, *139*, 13660.

(36) D'yakonov, V. A.; Trapeznikova, O. A.; Ibragimov, A. G.; Dzhemilev, U. M. Synthesis and Transformations of Metallacycles: New Method for the Synthesis of Spiro[3.3]heptanes and Spiro[3.4]octanes by Cycloalumination of Methylidenecyclobutanes with  $Et_3Al$  in the Presence of a Cp<sub>2</sub>ZrCl<sub>2</sub> Catalyst. *Russ. Chem. Bull. Int. Ed.* **2009**, *58*, 948.

(37) Dockery, K. P.; Bentrude, W. G. Free Radical Chain Reactions of [1.1.1]Propellane with Three-Coordinate Phosphorus Molecules. Evidence for the High Reactivity of the Bicyclo[1.1.1]pent-1-yl Radical. *J. Am. Chem. Soc.* **1997**, *119*, 1388.

(38) Wiberg, K. B.; Waddell, S. T. [1.1.1]Propellane: Reaction with Electron Deficient Alkenes and Alkynes. *Tetrahedron Lett.* **1987**, 28, 151.

(39) Xiao, L.-J.; Cheng, L.; Feng, W.-M.; Li, M.-L.; Xie, J.-H.; Zhou, Q.-L. Nickel(0)-Catalyzed Hydroarylation of Styrenes and 1,3-Dienes with Organoboron Compounds. *Angew. Chem., Int. Ed.* **2018**, 57, 461.

(40) See Supporting Information for full details.

(41) This ligand has been successfully employed as a computational surrogate in related computational investigations, see: (a) Menezes da Silva, V. H.; Braga, A. A. C.; Cundari, T. R. N-Heterocyclic Carbene Based Nickel and Palladium Complexes: A DFT Comparison of the Mizoroki-Heck Catalytic Cycles. Organometallics 2016, 35, 3170. (b) Guest, D.; Menezes da Silva, V. H.; de Lima Batista, A. P.; Roe, S. M.; Braga, A. A. C.; Navarro, O. (N-Heterocyclic Carbene)-Palladate Complexes in Anionic Mizoroki-Coupling Cycles: A Combined Experimental and Heck Computational Study. Organometallics 2015, 34, 2463. (c) Shen, A.; Hu, Y.-C.; Liu, T.-T.; Ni, C.; Luo, Y.; Cao, Y.-C. Supporting Ligand-N-Heterocyclic Carbene Palladium Complexes: Assisted Characterization, Computation, and Catalytic Activity in Suzuki-Miyaura Cross Coupling between Aryl and Heteroaromatic Chlorides and Various Boronic Acids. Tetrahedron Lett. 2016, 57, 2055.

(42) Using 1,3-dimethylimidazolium iodide in place of SIMes·HCl in the reaction of 4-vinylbiphenyl (8a) (see Table 1, entry 10) provided 7a in 71% isolated yield, along with 26% recovered 4-vinylbiphenyl.

 $\left(43\right)$  The individual steps of the alkene substitution were not modelled.

(44) For computed examples of bridging propellane ligands, see: Zhang, C.; Ning, L.; Li, J. Evidence for Charge-Shift Bonding in [1.1.1]Propellanes  $C_5H_6$  and  $E_5Me_6$  (E = Si, Ge, Sn): A Theoretical Investigation. *Organometallics* **2013**, *32*, 7083.

(45) (a) Schwager, H.; Krüger, C.; Neidlein, R.; Wilke, G. Nickel Complexes with a Propellane Structure. *Angew. Chem., Int. Ed.* **1987**, 26, 65. (b) Schwager, H.; Benn, R.; Wilke, G. Reactions of  $(\eta^3$ -Allyl) $(\eta^5$ -cyclopentadienyl)palladium with Cyclopropabenzenes. *Angew. Chem., Int. Ed.* **1987**, 26, 67.

(46) (a) Stohrer, W. D.; Hoffmann, R. Electronic Structure and Reactivity of Strained Tricyclic Hydrocarbons. J. Am. Chem. Soc. 1972, 94, 779. (b) Newton, M. D.; Schulman, J. M. Theoretical Studies of Tricyclo[1.1.1.0<sup>1,3</sup>]pentane and Bicyclo[1.1.1]pentane. J. Am. Chem. Soc. 1972, 94, 773. (c) Chakrabarti, P.; Seiler, P.; Dunitz, J. D.; Schlüter, A. D.; Szeimies, G. Experimental Evidence for the Absence of Bonding Electron Density between Inverted Carbon Atoms. J. Am. Chem. Soc. 1981, 103, 7378. (d) Jackson, J. E.; Allen, L. C. The Carbon 1-Carbon 3 Bond in [1.1.1]Propellane. J. Am. Chem. Soc. 1984, 106, 591. (e) Politzer, P.; Jayasuriya, K. J. Mol. Struct. (THEOCHEM) 1986, 135, 245. (f) Jarret, R. M.; Cusumano, L. <sup>13</sup>C-<sup>13</sup>C Coupling in [1.1.1]Propellane. Tetrahedron Lett. 1990, 31, 171. (g) Galasso, V. Theoretical Study of the  ${}^{1}J({}^{13}C{}^{13}C)$  Indirect Nuclear Spin-Spin Coupling Constant of the Bridgehead Bond in [1.1.1]Propellane and Related Systems. Chem. Phys. Lett. 1994, 230, 387. (h) Werner, M.; Stephenson, D. S.; Szeimies, G. Synthesis of [1.1.1]Propellanes by Bridging of Bicyclo[1.1.0]butanes. Liebigs Ann. Chem. 1996, 1705.

(47) Wu, W.; Gu, J.; Song, J.; Shaik, S.; Hiberty, P. C. The Inverted Bond in [1.1.1]Propellane is a Charge-Shift Bond. *Angew. Chem.*, *Int. Ed.* **2009**, *48*, 1407.

Public

(48) Shaik, S.; Danovich, D.; Wu, W.; Hiberty, P. C. Charge-Shift Bonding and its Manifestations in Chemistry. *Nat. Chem.* **2009**, *1*, 443.

(49) Messerschmidt, M.; Scheins, S.; Grubert, L.; Pätzel, M.; Szeimies, G.; Paulmann, C.; Luger, P. Electron Density and Bonding at Inverted Carbon Atoms: An Experimental Study of a [1.1.1]Propellane Derivative. *Angew. Chem., Int. Ed.* **2005**, *44*, 3925.

(50) Chiang, J. F.; Bauer, S. H. Molecular Structure of Bicyclo[1.1.1]pentane. J. Am. Chem. Soc. **1970**, *92*, 1614.

(51) (a) Bader, R. F. W. *Atoms in Molecules: A Quantum Theory*. Oxford University Press: Oxford, 1990. (b) Popelier, P. *Atoms in Molecules: An Introduction*. Prentice Hall: Harlow, 2000.

(52) (a) LaPierre, E. A.; Piers, W. E.; Gendy, C. Redox-State Dependent Activation of Silanes and Ammonia with Reverse Polarity (PC<sub>carbene</sub>P)Ni Complexes: Electrophilic vs. Nucleophilic Carbenes. *Dalton Trans.* 2018, 47, 16789. (b) LaPierre, E. A.; Piers, W. E.; Gendy, C. Divergent Reactivity of CO<sub>2</sub>, CO, and Related Substrates at the Nickel Carbon Double Bond of (PC<sub>carbene</sub>P)Ni(II) Pincer Complexes. *Organometallics* 2018, 37, 3394. (c) Lash, T. D.; Ferrence, G. M. Metalation and Selective Oxidation of Diphenyl-23-oxa-, -thia-, and -selena-21-carbaporphyrins. *Inorg. Chem.* 2017, 56, 11426. (d) Stander-Grobler, E.; Strasser, C. E.; Schuster, O.; Cronje, S.; Raubenheimer, H. G. Amine-Substituted α-N(Standard)- and δ-N(Remote)-Pyridylidene Complexes: Synthesis and Bonding. *Inorg.*

*Chim. Acta* **2011**, *376*, 87. (e) Stander-Grobler, E.; Schuster, O.; Heydenrych, G.; Cronje, S.; Tosh, E.; Albrecht, M.; Frenking, G.; Raubenheimer, H. G. Pyridine-Derived N-Heterocyclic Carbenes: An Experimental and Theoretical Evaluation of the Bonding in and Reactivity of Selected Normal and Abnormal Complexes of Nickel(II) and Palladium(II). *Organometallics* **2010**, *29*, 5821. (f) Schneider, S. K.; Julius, G. R.; Loschen, C.; Raubenheimer, H. G. Frenking, G.; Herrmann, W. A. A First Structural and Theoretical Comparison of Pyridinylidene-Type rNHC (Remote N-Heterocyclic Carbene) and NHC Complexes of Ni(II) Obtained by Oxidative Substitution. *Dalton Trans.* **2006**, 1226.

(53) (a) Harrison, D. J.; Daniels, A. L.; Korobkov, I.; Baker, R. T. d<sup>10</sup> Nickel Difluorocarbenes and Their Cycloaddition Reactions with Tetrafluoroethylene. *Organometallics* **2015**, *34*, 5683. (b) Hughes, R. P.; Carl, R. T.; Samkoff, D. E.; Davis R. E.; Holland, K. D. Transition-Metal Chemistry of Octafluorocyclooctatetraene. Synthesis and X-ray Structure of a Novel Five-Coordinate *cis*-Dialkylnickel Complex. *Organometallics* **1986**, *5*, 1053.

(54) Farrugia, L. J.; Evans, C.; Tegel, M. Chemical Bonds without "Chemical Bonding"? A Combined Experimental and Theoretical Charge Density Study on an Iron Trimethylenemethane Complex. *J. Phys. Chem. A* **2006**, *110*, 7952.

Public

