

* Present address: Institute of Scientific and Industrial Research, Osaka Univ., Sakai, Osaka.

TABLE I. REACTION OF SYNTHETIC GAS WITH BENZALDEHYDEANILS

Anil	Solvent	Reaction temp., °C	Reaction time, min.	Product	Yield, %
$C_6H_5N=CHC_6H_5$	B	135	110	$C_6H_5NHCH_2C_6H_5$	80
$C_6H_5N=CHC_6H_5$	A B	120	60	$C_6H_5NHCH_2C_6H_5$	79
$p-ClC_6H_4N=CHC_6H_5$	B	150	70	$p-ClC_6H_4NHCH_2C_6H_5$	79
$p-ClC_6H_4N=CHC_6H_5$	A B	130	130	$p-ClC_6H_4NHCH_2C_6H_5$	81
$p-MeC_6H_4N=CHC_6H_5$	B	140	90	$p-MeC_6H_4NHCH_2C_6H_5$	82
$p-MeC_6H_4N=CHC_6H_5$	A B	120	60	$p-MeC_6H_4NHCH_2C_6H_5$	80
$p-MeOC_6H_4N=CHC_6H_5$	B	130	60	$p-MeOC_6H_4NHCH_2C_6H_5$	83
$p-MeOC_6H_4N=CHC_6H_5$	A B	120	50	$p-MeOC_6H_4NHCH_2C_6H_5$	82
$p-NO_2C_6H_4N=CHC_6H_5$	B	130	80	$p-NO_2C_6H_4NHCH_2C_6H_5$	80
$p-NO_2C_6H_4N=CHC_6H_5$	A B	120	70	$p-NO_2C_6H_4NHCH_2C_6H_5$	78

Catalyst, $[Co(CO)_4]_2$, 0.03 millimol./ml.; Solvent, B=benzene, AB=alcohol-benzene (1:1); Gas, CO(100 atm.) + H_2 (100 atm.).

TABLE II. FORMATION OF DIPHENYLUREAS BY THE REACTION OF AZOBENZENES AND NITROBENZENE WITH SYNTHETIC GAS

	$C_6H_5NHCONHC_6H_5$ %	$R-C_6H_4NHCONHC_6H_5$ %	$R-C_6H_4NHCONHC_6H_4-R$ %
$C_6H_5NO_2$	5~6		
$C_6H_5N=NC_6H_5$	15~20		
$C_6H_5NH-NHC_6H_5$	25~30		
$p-ClC_6H_4N=NC_6H_5$	trace	5	3
$p-MeC_6H_4N=NC_6H_5$	trace	12	5

Reaction conditions are the same in Table I.



noteworthy, because it demonstrates, that some new kind of CO-reaction has occurred, and especially the fact that a formation of symmetrically substituted urea derivatives were observed from asymmetrically substituted azo-compounds was of special note.

As the reaction appeared to be a new one, the authors planned further experiments so as to obtain more detailed information about the high pressure reaction of carbon monoxide.

Hydrazobenzene gave 50% yield of azobenzene under the similar condition, at the reaction temperature which was lower than that required for the reduction (80~90°C). From this experiment it could be postulated that urea could have formed from hydrazobenzene via azobenzene. However, the yields of diphenylurea were in all cases higher when hydrazobenzene was used instead of azobenzene. In the case of nitrobenzene, the formation of urea derivative was also observed and its formation could be interpreted by the intermediate formation of either hydrazobenzene or azobenzene.

In order to discuss the possible mechanism for the formation of urea derivative,

one has to consider either 1) fission of hydrazo linkage followed by addition of CO, or 2) that the azo group remains without cleavage until CO is added across the double bond and then the -N-N- bond is subjected to fission by hydrogenolysis as will be shown below (Eq. 2). Eq. 1 was suggested earlier by W. Reppe¹⁾ as a possible mechanism of the oxo-reaction and the carboxylation of olefins. Accordingly one might propose Eq. 2 for the urea formation.



Since azobenzene is reduced for the most part to aniline, it would be an acceptable argument that the formation of urea could have resulted from the reaction of

1) W. Reppe, *Ann.*, 582, 40 (1953).

aniline and carbon monoxide. Pino²⁾ discussed the possibility of formation of urea derivatives from two moles of aniline and one mole of carbon monoxide under a dehydrating condition, and Buckley³⁾ reported the formation of formylaniline by the reaction of aniline and carbon monoxide.

However, all the authors' tests with aniline and with formylaniline failed to show that the urea-formation is the result of the reaction of aniline and carbon monoxide.

An alternate mechanism would presumably be a fission of the azo-compound by means of carbon monoxide and hydrogen, leading to the primary formation of phenylisocyanate and aniline. However, the authors failed to isolate such an intermediate to prove this mechanism.

As another possibility, hydrazobenzene might well be accounted for the reaction. In this regard the authors will discuss it in later papers.

Experimental

1) Reduction of Benzaldehydeanil with Synthetic Gas.—Benzaldehydeanil (5 g.) was dissolved in 50 ml. of benzene or benzene-alcohol mixture (1:1) and charged in an autoclave of 100 ml. content made of 18-8 stainless steel together with cobalt carbonyl (1.0 g.). After replacement of air, the synthetic gas was compressed up to 200 atm. (CO, 100 atm. and H₂, 100 atm.). The autoclave was heated at 120~130°C, when the pressure began to descend, and after completion of gas absorption it was allowed to cool. The content of the autoclave was heated in a flask on a water bath and after decomposition of the cobalt catalyst and filtration the filtrate was distilled in vacuo, affording 4 g. of benzyaniline, m. p. 38°C, which was identified by the mixed melting point test with an authentic sample.

The essential feature of the experimental procedures are all the same as described above, and thus *p*-chloro, *p*-methyl-, *p*-methoxy- and *p*-nitro-benzaldehydeanil were subjected to exami-

nation. The results are given in Table I. The melting points and the boiling points all agree with the values described in the literatures. *p*-Nitrocompound was reduced under the present condition selectively at the anil double bond and the nitro group remained unchanged.

2) Reaction of 3-Phenyl-2-thioxopropionic Acid and Carbon Monoxide.—3-Phenyl-2-thioxopropionic acid (3 g.) in benzene (50 ml.) was allowed to react with carbon monoxide and hydrogen of 200 atm. in the presence of cobalt catalyst (1 g.) at 120~130°C for 2 hr. No gas absorption was observed, but after cooling and degassing, it was found that hydrogen sulfide gas had formed. A definite product could not be isolated except 2-phenylpropionic acid amide melting at 102°C.

3) Formation of Diphenylurea as a By-product in the Reaction of Azobenzene and Synthetic Gas.—Azobenzene (5 g.) yielded 1.1 g. of diphenylurea (20% yield) in a form of a difficultly soluble product, which was recrystallized from alcohol.

4) Urea Derivatives from *p*-Chloroazobenzene.—*p*-Chloroazobenzene (5 g.) was treated as in exp. 1 and from the reaction product 0.6 g. (12%) of a mixture of urea compounds was obtained, and was separated into two parts: *p*,*p'*-dichlorodiphenylurea as the difficultly soluble part in alcohol in colorless prisms, m. p. 286°C (with decomposition), and as the soluble part *p*-chlorodiphenylurea in scaly crystals, m. p. 244°C.

Anal. Found: C, 56.66; H, 3.70; Cl, 25.20; N, 10.36. Calcd. for C₁₃H₁₀Cl₂N₂O: C, 56.93; H, 3.64; Cl, 25.26; N, 10.21%.

Anal. Found: C, 64.06; H, 4.64; Cl, 14.32; N, 11.55. Calcd. for C₁₃H₁₁ClN₂O: C, 64.19; H, 4.52; Cl, 14.40; N, 11.52%.

5) Urea Derivatives from *p*-Methylazobenzene.—*p*-Methylazobenzene (5 g.) was treated as in exp. 4 and from the filtrate of the reaction mixture a mixture (1.2 g., 22%) of urea was separated. From a difficultly soluble portion *p*,*p'*-dimethylphenylurea (0.3 g.) was obtained in colorless needles, m. p. 262°C and from the soluble portion *p*-dimethyldiphenylurea (0.7 g., 12%) was obtained, and further a small quantity of unsubstituted diphenylurea was obtained from the last portion of the soluble residue.

Anal. Found: C, 74.81; H, 6.70; N, 11.78. Calcd. for C₁₅H₁₆N₂O: C, 75.00; H, 6.66; N, 11.66%.

Anal. Found: C, 74.15; H, 6.20; N, 12.51. Calcd. for C₁₄H₁₄N₂O: C, 74.33; H, 6.19; N, 12.38%.

Department of Chemistry
Faculty of Science
Osaka University
Nakanoshima, Osaka

2) P. Pino and C. Paleari, *Gazz. Chim. Ital.*, **81**, 646 (1951); *Chem. Abstr.*, **46**, 7063 (1952).

3) G. D. Buckley and N. H. Ray., *J. Chem. Soc.*, **1949**, 1151.