mixture was kept overnight at 70 °C and worked up as described for 14 above. The orange oil obtained (0.45 g) was chromatographed on dry silica column (Woelm TSC) by using toluene as the eluant. Crystallization from EtOH gave 0.35 g (65%) of colorless crystals of isopropyl dimesitylacetate: mp 90 °C; UV (hexane) λ_{max} 207 nm (log ϵ = 4.56), 265 (2.89); IR (Nujol) ν_{max} 1725 (C=O), 1610 (C=C) cm⁻¹; ¹H NMR $(CD_2Cl_2, room temperature) \delta 1.16 (6 H, d, J = 6.3 Hz, 2Me), 1.95 (12)$ H, s, 4Me), 2.14 (6 H, s, 2Me), 4.98 (1 H, m, J = 6.3 Hz, CH), 5.18 (1 H, s, CH), 6.70 (4 H, s, Mes-H); mass spectrum, m/z 338 (M, 13%), 251 (Mes₂CH⁺, 100%), 236 (Mes₂CH⁺ - Me, 4%), 221 (Mes₂CH⁺ 2Me, 10%), 206 (Mes₂CH⁺ - 3Me, 6%), 43 (C₃H₇, 8%). Anal. Calcd for C₂₃H₃₀O₂: C, 81.61; H, 8.93. Found: C, 81.97; H, 9.05.

Equilibration Studies. Approximately 20 mg of the ketone or the enol was dissolved in 25 mL of spectroscopic hexane, and the solution was divided into several pressure ampules which were wrapped in aluminum foil and were kept in an oil bath in the dark at 80.3 ± 0.1 °C. If the ampules of the enols were kept at daylight, decomposition took place as reflected by the appearance of new (non-ketonic) signals in the NMR. For example, the reaction of 5 gave signals (in CDCl₃) at 0.99 (s), 2.19 (s), 4.7 (AB q), 6.6 (s), and 9.54 (s) ppm.

Samples were withdrawn after a few hours, the solvent was evaporated, and the ¹H NMR spectra in CDCl₃ were recorded. The relative ratios of the two species were determined by integration of the methyls of the alkyl groups for each one of the keto/enol pairs. In several cases, a corroboration for this ratio was obtained from the integration of the ketonic CH and enolic OH protons, but this was impossible when traces of CF₃COOH still remained, since exchange with the OH resulted in a higher integration. Care was taken to ensure complete relaxation of the various hydrogens in order to obtain reliable integration. When two or three samples taken at different reaction times gave a similar enol/ketone composition, this was regarded as the equilibrium value. In all cases, the equilibrium compositions obtained by starting from the enol or from the ketone were identical within the accuracy of the integration.

Acknowledgment. We are indebted to Silvio Biali for discussions and for some experiments with 13 and 14, to Prof. H. Schwarz for the mass spectra of 14, to Dr. S. Cohen for the X-ray crystallography of compound 14, and to Dr. L. Radom for discussions. This work was supported by a grant from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel, to whom we are grateful.

Registry No. 1, 54288-04-9; 2, 89959-15-9; 3, 96040-90-3; 4, 96040-91-4; 5, 89959-16-0; 6, 94203-58-4; 7, 96040-92-5; 8, 96040-93-6; 9, 96040-94-7; 10, 96040-95-8; 11, 87871-33-8; 12, 96040-98-1; 13, 96040-97-0; 14, 96040-96-9; dimesitylacetic acid, 5740-42-1; isopropyl bromide, 75-26-3; tert-butyl bromide, 507-19-7; ethyl bromide, 74-96-4.

Supplementary Material Available: Tables S1-S4 giving the crystallographic data for compound 14 (6 pages). Ordering information is given on any current masthead page.

Stereo- and Regioselective Palladium-Catalyzed, 1,4-Acetoxychlorination of 1,3-Dienes. 1-Acetoxy-4-chloro-2-alkenes as Versatile Synthons in Organic Transformations

Jan-E. Bäckvall,* Jan-E. Nyström, and Ruth E. Nordberg

Contribution from the Department of Organic Chemistry, Royal Institute of Technology, S-100 44 Stockholm, Sweden. Received November 8, 1984

Abstract: Palladium-catalyzed oxidation of 1,3-dienes in the presence of LiCl and LiOAc produces 1-acetoxy-4-chloro-2-alkenes with high selectivity. The reaction is stereospecific and cyclic dienes give an overall cis 1,4-addition (>97-98% cis). The stereospecificity of the reaction also holds in acyclic systems as shown by the oxidation of (E,E)- and (E,Z)-2,4-hexadiene to (R^*, R^*) - and (R^*, S^*) -10, respectively. The mechanism of the acetoxychlorination is now well understood. It proceeds via a trans acetoxypalladation of the diene to produce an intermediate (4-acetoxy-1,2,3-ŋ³-allyl)palladium species, followed by an oxidation-induced nucleophilic attack at C-1 with inversion. Kinetic studies indicate that p-benzoquinone, which is a unique oxidant for the reaction, not only serves as an oxidant but also acts as a ligand to palladium. The chloroacetate products are useful synthons in organic transformations. Sequential substitutions of the chloro and acetoxy groups allow a regiochemical choice, and the fact that the allylic chloro group can be substituted with either clean retention or inversion allows a stereochemical choice. These principles are demonstrated in a number of cases. It is shown that the acetoxychlorination approach allows a complete control of the 1,4-relative stereochemistry with a unique choice of functionality at the asymmetric center.

Regio- and stereocontrolled 1.4-addition to conjugated dienes is of synthetic interest. Methods for achieving such additions include cycloaddition of singlet oxygen^{1,2} and nitroso compounds^{3,4}

followed by reduction.⁵ We recently reported a few stereo- and regioselective 1,4-additions to conjugated dienes.^{6,7} These oxi-

0002-7863/85/1507-3676\$01.50/0 @ 1985 American Chemical Society

^{(1) (}a) Balci, M. Chem. Rev. 1981, 81, 91. (b) Schenck, G. O.; Dunlop, D. E. Angew. Chem. 1956, 68, 248.

^{(2) (}a) Kaneko, C.; Sugimoto, A.; Tanaka, S. Synthesis 1974, 876. (b) Gollnick, K.; Griesbeck, A. Tetrahedron Lett. 1983, 24, 3303

^{(3) (}a) Belleau, B.; Au-Young, Y. K. J. Am. Chem. Soc. 1963, 85, 64. (b)
Krezse, G.; Dittel, W.; Melzer, H. Liebiqs Ann. Chem. 1981, 224 and references cited. (c) Keck, G. E.; Fleming, S. A. Tetrahedron Lett. 1978, 4763.
(4) (a)Streith, J.; Augelmann, G.; Fritz, H.; Strub, H. Tetrahedron Lett.
1982, 23, 1909. (b) Becker, Y.; Eisenstadt, A.; Shvo, Y. Tetrahedron 1978, 24 and 1978.

^{34, 801.}

^{(5) 1,4-}Additions to conjugated dienes have been obtained in a limited number of electrophilic and radical additions. Most of these reactions, however, are not general, and furthermore, the reactions are either not stereoselective or the stereochemistry of the additions is unknown: (a) Neale, R. S. Synthesis 1971, 1. (b) Hassner, A.; Keogh, J. Tetrahedron Lett. 1975, 1575. (c) Truce, W. E.; Goralski, C. T.; Christensen, L. W.; Bavra, R. M. J. Org.
 Chem. 1970, 35, 4217. (d) Adams, D. R.; Bhatuagar, S. P.; Cookson, R. C.
 Tetrahedron Lett. 1974, 3197. (e) Matsumoto, H.; Nateaur, T.; Nikaido, T.;
 Nagai, Y. Chem. Lett. 1978, 115.
 (d) Rescher Lett. 1978, 125. (6) Bäckvall, J. E. Acc. Chem. Res. 1983, 16, 335.

 Table I.
 Palladium(II)-Catalyzed 1,4-Acetoxychlorination of Conjugated Dienes

Entry	Diene	Method ^a ,reac (addition ti	tion time me) h	Product(s)		Yield %
1a b		A, 26 (0) B, 20 (16)		CI OAC	(E/Z = 9/1)	81 ^b 78 ^b
2a b		A, 22 (0) B, 26 (16)		CIOAc	(E/Z=36/1)	76 [°] 74 [°]
3a	<u></u>	A, 20 (0)		CI		44 ^d
Ъ		B, 32 (28)		(Z) - <u>3</u>	(E/Z=1/10)	45 ^d
4	(E)	B, 49 (45)		CI CI OAc	(E/Z = 2 5/1)	35 ^e
5	(E or Z)	B, 20 (16)	CI 🔪	<u>5a</u> Ac0	<u>5</u> <u>b</u> (1.5:1)	51 ^f
		CI~~			: AcO	R
6	R=Ph	B, 21 (14)	6 <u>a</u> (44)	: <u>6b</u> (43)	5 (13)	36 ^f
7	R=COOEt	B, 42 (40)	<u>7a</u> (44)	: <u>75</u> (36)	: <u>Z</u> '(11)	58 ^f
8	(E,E)	A, 24 (18)		CI OAc	(>95% R [*] ,R [*]) (>99% E)	55 ^g
9	(E,Z)	A, 24 (0)		CI CI OAc	(>95% R [*] ,S [*]) (>99% E)	62 ^g
10	\bigcirc	B, 5 (3)		CI - OAc	(>98% cis)	89 ^g
11	\bigcirc	A, 10 (3)			(>97% cis)	55 ⁸
12	\bigcirc	B , 4.5 (0.5)	Ас0 - («СН3//СН3=	C1 C1 C1	0Ac <u>14</u> (43) (4/1)	65 ^g
13	\bigcirc	B, 12 (0)		CI-CI-OAc	(>98% cis)	74 ^g
14	OAc	B, 72 (0)		CI - OAc	(>95% β.β.α)	58 ^g
15	\bigcirc	B, 36 (0) ^h	c1 -	OAc	(3 : 1) (2 : Cl	61

^a The amount of Pd(II) catalyst was 5 mol% for cyclic and 7.5 mol% for acyclic dienes. Method A: a two-phase system (HOAc/pentane) was used. Method B: a one-phase system (HOAc) was used. ^b Contaminated with $CH_2 = CH - CH(Cl)CH_2OAc$ (1') (9%). ^c Contaminated with 2' (4%) and $ClCH_2 - CH = C(CH_3)CH_2OAc$ (20) (8%). ^d Contaminated with $CH_2 = CCH_3 - CCH_3(Cl)CH_2OAc$ (3') (9%). ^e Contaminated with $CH_3CH = CH - CCH_3(Cl)CH_2OAc$ (7%). ^f >95% E configuration. ^g >99% 1,4-addition. ^h Seven equivalents of LiOAc-2H₂O and 1.2 equivalents of LiCl were used, and the reaction was performed at 43 °C.

dation reactions are catalyzed by palladium(II) and proceed via a (π -allyl)palladium intermediate (Scheme I). The addition of the nucleophiles X and Y is stereo- and regioselective, and importantly, the stereochemistry of the attack by the second nucleophile Y can be controlled in some cases to proceed either from the opposite side (trans attack) or from the same side (cis attack) as the metal atom. Three principal reactions, 1,4-diacetoxylation,^{7a,b} 1,4-acetoxychlorination,^{7c} and 1,4-acetoxytrifluoroacetoxylation^{7d} have been reported which all have their specific synthetic advantages. In this paper, we give a complete



account of the 1,4-acetoxychlorination reaction, discuss the mechanism, and demonstrate its synthetic utility in selective organic transformations.

Results

1,4-Acetoxychlorination. Palladium-catalyzed oxidation of conjugated dienes in acetic acid in the presence of LiCl and LiOAc proceeds smoothly, to selectively produce 1-acetoxy-4-chloro-2-alkenes (eq 1, Table I). All reactions were performed at room

^{(7) (}a) Bäckvall, J. E.; Nordberg, R. E. J. Am. Chem. Soc. 1981, 103, 4959.
(b) Bäckvall, J. E.; Byström, S. E.; Nordberg, R. E. J. Org. Chem. 1984, 49, 4619.
(c) Bäckvall, J. E.; Nordberg, R. E.; Nyström, J. E. Tetrahedron Lett. 1982, 23, 1617.
(d) Bäckvall, J. E.; Vågberg, J.; Nordberg, R. E. Tetrahedron Lett. 1984, 25, 2717.

$$-C = C - C =$$

temperature, and the oxidant used was p-benzoquinone which was found to have a remarkable ability to promote a stereo- and regioselective acetoxychlorination. A number of dienes were tested and so far only two dienes, 1,3-cyclopentadiene and 1-acetoxy-2,4-hexadiene, failed to give an oxidation product under the present reaction conditions. Interestingly, 4-methyl-1,3-pentadiene and 2,5-dimethyl-2,4-hexadiene gave no chloroacetate but afforded the chloro alcohol 8 and hydroxyacetate 9, respectively.⁸

The chemoselectivity for chloroacetate over diacetate and dichloride is high and usually exceeds 99%. On prolonged reaction time, however, a few percent of diacetate was observed in some



cases due to a secondary substitution reaction, but the ratio chloroacetate/diacetate was always >20. The amount of dichloride was usually under the detectable limit, i.e., chloroacetate/dichloride >100. In the case of butadiene, we determined the ratio 1-acetoxy-4-chloro-2-butene/1,4-dichloro-2-butene to be 260:1.

The amount of palladium catalyst utilized (usually Pd(OAc)₂ but Li_2PdCl_4 gives identical results) was 5 mol % for cyclic dienes and 7.5 mol % for acyclic dienes. These amounts were used for convenience in the laboratory procedure, but the reaction works in several cases with a much lower amount of catalyst.⁹ For example, the oxidation of 1,3-cyclohexadiene using 0.2 mol % $Pd(OAc)_2$ selectively produced the chloroacetate in an isolated yield of 60% which corresponds to 300 turnovers.

The acetoxychlorination of acyclic dienes can, a priori, give a product with either an E or Z double bond. When the 1,3-diene is unsubstituted in the 2- and 3-position, the 1,4-chloroacetate formed is almost exclusively of the E configuration (90-100%, entries 1, 5–7, 8, and 9). The highest degree of E stereochemistry was obtained for the 2,4-hexadienes which furnished products with >99% E geometry of the double bond (entries 10 and 11). A substituent in the 2- and/or 3-position increased the amount of Z product (entries 2-4). In the series butadiene, isoprene, (E)-2-methylpentadiene, and 2,3-dimethylbutadiene, the E/Z ratio decreased and was 9/1, 3.6/1, 2.5/1, and 1/10, respectively (entries 1, 2, 4, and 3).

The acetoxychlorination is highly stereospecific, and cyclic dienes afforded products with >97-98% cis stereochemistry. The ¹H NMR spectrum of the product **11** from 1,3-cyclohexadiene (entry 10) is consistent with a cis configuration and shows the CH2-CH2 grouping more concentrated than in the corresponding trans isomer.^{7a,b} Conclusive evidence for the cis stereochemistry of 11 follows from its transformation to the known^{7a,b} trans-1,4diacetoxy-2-cyclohexene by reaction with KOAc in Me₂SO. Also the selective transformations of 11 performed previously^{7c} confirms the cis configuration of 11. Interestingly, the stereospecific addition could be extended to acyclic systems. Palladium-catalyzed acetoxychlorination of (E, E)-2,4-hexadiene selectively produced a single diastereoisomer (R^*, R^*) -10 (entry 8) which corresponds to an overall cis addition of Cl and OAc across the S-trans conformation of the diene. In an analogous manner (E,Z)-2,4hexadiene selectively afforded (R^*, S^*) -10 (entry 9). The diastereomeric purity (>95%) of (R^*,R^*) - and (R^*,S^*) -10 was determined by ¹H NMR. The configurational assignments of (R^*,R^*) - and (R^*,S^*) -10 were made in analogy with the products



from the corresponding palladium-catalyzed diacetoxylation,7b and it was confirmed by chemical transformations (vide infra).

The 1,4-regioselectivity is usually high. For the 2,4-hexadienes and the cyclic dienes, except 1,3-cyclooctadiene, the regioselectivity for 1,4-addition is complete and no 1,2-adduct could be detected in the crude product (>99% 1,4-addition). 1,3-Cyclooctadiene was the only case of those studied where the use of p-benzoquinone as the oxidant gave a considerable amount of 1,2-isomer. This diene afforded 1,4-chloroacetate 17 and 1,2-chloroacetate 17' in a ratio of 3:1, with a cis configuration of both isomers (entry 15). The relative amount of 1,2-adduct among the acyclic dienes having at least one terminal double bond varied between 4% and 13% (entries 1, 2, and 3-7). When p-benzoquinone was replaced by MnO₂ as the oxidant, the relative amount of 1,2-addition increased to 25-40%.

The regioselectivity in the oxidation of an unsymmetrically substituted diene was investigated for both cyclic and acyclic dienes. A methyl group in the 2-position directed the chloro group to the 1-position, and a 4-acetoxy-1-chloro-2-methyl-2-alkene was selectively formed (entries 2, 4, and 11). This regioselectivity is of great use in the synthetic transformations of the chloroacetates (vide infra). On the other hand, the regioselectivity obtained with substituents in the 1-position was less general. Thus, 1,3-pentadiene and related 1,3-dienes showed a slight preference for the chloroacetate in which acetate has attacked the most substituted carbon (entry 5-7). 4-Methyl-1,3-pentadiene, on the other hand, gave a highly regioselective reaction, but in this case, the product was the 1,4-chloro alcohol 8.

To determine the diastereoselectivity of the reaction toward a chirality in the diene, cyclic dienes with a substituent outside the diene unit were studied. In particular the directing effect of an allylic or homoallylic substituent is of interest. Acetoxychlorination of 6-acetoxy-1,3-cycloheptadiene (entry 14) afforded only one isomer according to ¹H NMR, GLC, and HPLC. In analogy with the palladium-catalyzed diacetoxylation of the same diene,^{7b} we have assigned it as the 1β , 4β , 6α -isomer 16. The ¹H NMR spectrum of 16 is consistent with this configuration. The selective functionalization reactions shown in Scheme II, where the chloro group is substituted with either retention or inversion, confirms the stereochemistry of 16.

Palladium-catalyzed 1,4-acetoxychlorination of 5-methyl-1,3cyclohexadiene was less selective (entry 12). Two regioisomers were formed in a ratio of 57:43, and each regioisomer was found to be a 4:1 mixture of diastereoisomers due to the 5-methyl group.

One disturbing side reaction in the 1,4-acetoxychlorination reaction is Diels-Alder addition between the diene and pbenzoquinone. This reaction could be depressed by keeping the diene concentration low in the reaction mixture. This was achieved by slow addition of the diene and/or by the use of a two-phase system (acetic acid-pentane).7b For the cyclic dienes (entries 10-15), the relative yield of Diels-Alder product thus was usually <5% in the crude product. For the acyclic dienes, the relative yield of Diels-Alder product varied from 5% to 20%. The Diels-Alder side product was readily removed from the product by several methods (see Experimental Section).

Because of the above mentioned side reaction, we tried to replace *p*-benzoquinone by other oxidants. Attempts to use chloranil or methyl-substituted benzoquinones failed and gave no reaction or a very slow reaction. When p-benzoquinone was replaced by MnO₂ or urea hydroperoxide, a much lower regio-

⁽⁸⁾ Due to the use of LiOAc·2H₂O, the acetic acid phase contains $\approx 2\%$ of water (by weight) which apparently has attacked in place of acetate: Nyström, J. E., unpublished results.
(9) Byström, S. E.; Bäckvall, J. E.; Nordberg, R. E.; Nyström, J. E.

Swedish Patent 8201 911-8, 1982.



Figure 1. Relative initial rate for the palladium-catalyzed 1,4-acetoxychlorination of 1,3-cyclohexadiene as a function of the amount of pbenzoquinone using 5 mol % Pd(OAc)₂.

Table II. (a) Competitive Experiments and (b) Absolute Relative Rates

а		b		
diene	rate	diene	rate	
(E,Z)-2,4-hexadiene	1	(Z)-1,3-pentadiene	1	
(E,E)-2,4-hexadiene	1.9	(E)-1,3-pentadiene	2.3	
isoprene	3.8	(E,Z)-2,4-hexadiene	4.4	
1,3-cyclohexadiene	50	(E,E)-2,4-hexadiene	8.3	

and stereospecificity was obtained. For example, palladiumcatalyzed acetoxychlorination of isoprene using MnO_2 as the oxidant (at 40 °C) produced the 1,4-isomer (E)-2 and the 1,2isomer 4-acetoxy-3-chloro-2-methyl-1-butene (2') in a ratio of 1.5:1. Interestingly, the 1,4-isomer was exclusively of E configuration (>95% E). This is in contrast to the 1,4-isomer obtained by using p-benzoquinone as the oxidant, which gave an E/Z ratio of 3.6/1 (entries 2a and 2b). Also, 1,3-cyclohexadiene, when oxidized with MnO₂ (at 40 °C), afforded a considerable amount of 1,2-isomer (1,2:1,4 = 15:85), and furthermore the 1,4-isomer was a cis and trans mixture (cis/trans = 1:3).¹⁰ We also tried $CuCl_2$ as the oxidant, since it is known that olefins are oxidized to vicinal chloroacetates in acetic acid by a Pd(II)/CuCl₂ system, however, with low product selectivity.^{11,12} Palladium-catalyzed oxidation of butadiene in acetic acid using $CuCl_2/O_2$ as the oxidant $(pO_2 = 5 \text{ kg/cm}^2)$ proceeded very slowly and gave only two turnovers after 22 h. The 1,4/1,2 ratio was approximately 4/1. Finally, the use of isoamyl nitrite in the oxidation of butadiene afforded 1,4-dichloro-2-butene as the only oxidation product.

The use of *p*-benzoquinone therefore seems to be essential for an efficient and selective reaction. We previously found^{7b} that in the related palladium-catalyzed 1,4-diacetoxylation, the dependence of the benzoquinone concentration on the reaction rate could be interpreted as benzoquinone acting also as a ligand. The dependence of the *p*-benzoquinone concentration on the rate of the palladium-catalyzed 1,4-acetoxychlorination of 1,3-cyclohexadiene is shown in Figure 1. The figure reveals that the rate of the oxidation increases linearly with the concentration of pScheme III







benzoquinone up to 120 mol % and then the increase becomes smaller.

The relative rates of the oxidation of some dienes were investigated and are given in Table II. The relative rates were determined ei her via competitive experiments (Table IIa) or via direct measu ements of the rate in individual experiments (Table IIb). The conpetitive experiments may give wrong relative rates if a competition occurs in the coordination step and if one diene coordinates strongly and completely blocks the coordination of the other diene. The relative rates obtained from the absolute rate determinations, however, indicate that this is not the case. Thus, both methods give a ratio of relative rates between (E,E)and (E,Z)-2,4-hexadiene of 1.9:1. This increase of the rate by approximately a factor of 2 on changing one double bond from Z to E was also observed for 1,3-pentadiene (Table IIb). The competitive experiments show that 1,3-cyclohexadiene reacts considerably faster than the acyclic dienes (Table IIa).

Synthetic Applications. An important aspect of the palladium-catalyzed acetoxychlorination reaction described in this paper is that the chloro and acetoxy groups can be sequentially substituted by two different nucleophiles (Scheme III). After a completely chemoselective nucleophilic substitution of the chloro group,^{7c,13} the acetoxy group can be substituted in a transitionmetal-catalyzed (Pd, Cu, Ni, Mo, and Fe) reaction.^{7c,14,15} This allows a regiochemical choice, and the principle was applied to the synthesis of the Monarch butterfly pheromone by using the chloroacetate 2 from isoprene.¹³ The chloroacetate 2 in this way becomes a useful building block for terpenoid synthesis. More generally the chloroacetate products can be considered as useful multiple coupling reagents (MCR),¹⁶ with two electrophilic sites.

Importantly, the substitution reactions in Scheme III are stereospecific, which allows the creation of new carbon-carbon bonds

⁽¹⁰⁾ It is remarkable that MnO_2 works as an oxidant in the palladiumcatalyzed acetoxychlorination, since it is extremely slow in the corresponding palladium-catalyzed diacetoxylation. We believe that this may be explained by an increase in oxidation potential of Mn(IV) in acetoxychlorination due to the presence of chloride ions.

 ^{(11) (}a) Henry, P. M. "Palladium-Catalyzed Oxidation of Hydrocarbons";
 D. Reidel: Dordrecht, 1980. (b) Henry, P. M. J. Org. Chem. 1974, 39, 3871; J. Am. Chem. Soc. 1972, 94, 7305. (c) Bäckvall, J. E. Tetrahedron Lett. 1977, 467

⁽¹²⁾ For oxidation of nonconjugated dienes using the Pd^{II}-CuCl₂ system in acetic acid, see: Heuman, A.; Waegell, B. Nouv. J. Chem. 1977, 1, 275. Heumann, A. Reglier, M., Waegell, B. Angew. Chem., Int. Ed. Engl. 1979, 18, 866, 867; Tetrahedron Lett. 1983, 24, 1971.

⁽¹³⁾ Nyström, J. E.; Bäckvall, J. E. J. Org. Chem. 1983, 48, 3947.

⁽¹³⁾ Nyström, J. E.; Backvali, J. E. J. O'g. Chem. 1983, 46, 3947.
(14) (a) Trost, B. M. Acc. Chem. Res. 1980, 13, 385. (b) Tsuji, J.
"Organic Synthesis with Palladium Compounds"; Springer-Verlag: Berlin, 1980. (c) Trost, B. M. Tetrahedron 1977, 33, 2615. (d) Trost, B. M.; Verhoeven, T. R. In "Comprehensive Organometallic Chemistry"; Wilkinson, G., Ed.; Pergamon Press: Oxford, 1982; Vol. 8, p 799.
(15) (a) Magid, R. M. Tetrahedron 1980, 36, 1901. (b) Yamamoto, Y.;
Verhoeven S. Vatarai, H.: Marwara, V. J. Am. Chem. Soc. 1980, 102.

Yamamoto, S.; Yatagai, H.; Maruyama, K. J. Am. Chem. Soc. 1980, 102, 2318. (c) Yamamoto, T.; Ishizu, J.; Yamamoto, A. Ibid. 1981, 103, 6863. (d) Trost, B. M.; Lautens, M. Ibid. 1981, 103, 5543. Roustan, J. L.; Marour, Y.; Houlihan, F. Tetrahedron Lett. 1979, 3721. (f) Goering, H. L.; Seitz, E. P.; Tseng, C. C. J. Org. Chem. 1981, 46, 5304.

⁽¹⁶⁾ Seebach, D.; Knochel, P. Helv. Chim. Acta 1984, 67, 261.

with control of the 1,4-relative stereochemistry. Other methods for obtaining a similar control of the 1,4-relative stereochemistry at carbon include transition-metal-catalyzed nucleophilic additions to 1,3-diene monoepoxides.^{17,18} An advantage of the acetoxychlorination approach over the metal-catalyzed nucleophilic addition to 1,3-diene monoepoxide is that it offers a stereochemical choice (Scheme III). By applying either a metal-catalyzed reaction or a classical $S_N 2$ reaction, the chloro group can be displaced with either retention or inversion at carbon. This was previously demonstrated on the chloroacetate 11 from 1,3-cyclohexadiene,⁷c and the principle was also shown on the chloroacetate 16 from 6-acetoxy-1,3-cycloheptadiene (Scheme II). In this way, a great number of 1,4-disubstituted 2-cycloalkenes with either cis or trans stereochemistry are available by choice. One example (Scheme IV), which illustrates the unique ability of the methodology, is the formal cis or trans 1,4-addition of the groups AcO- and (MeOOC)₂CH- to 1,3-cyclohexadiene via 11 in an overall yield of 84-85% and with a diastereoselectivity >98%!

We now also demonstrate how one may obtain a similar stereocontrol in an acyclic system (Scheme V). Reaction of (R^*, R^*) -10 with diethylamine in the presence of Pd(PPh₃)₄ afforded (R^*, R^*) -21a (>92% R^*, R^*) in 70% yield. When $(R^*, R^*$)-10 was allowed to react with diethylamine in acetonitrile in the absence of catalyst, (R^*, S^*) -21a (>90% R^*, S^*) was formed in 82% yield. Analogously, (R^*, S^*) - and (R^*, R^*) -21a were prepared by carrying out the same reactions on (R^*, S^*) -10. The two possible diastereoisomers (R^*, S^*) -21 and (R^*, R^*) -21 were prepared for three other nucleophiles (21b, c, and d) in a diastereomeric purity of 94-96%. Since the allylic acetoxy group can be stereospecifically replaced by carbon or nitrogen nucleophiles using either palladium¹⁴ or copper^{15b,f} catalysis, a great number of acyclic derivatives with a defined 1,4-relative stereochemistry between carbons are available by this method. Such a control of the relative stereochemistry between distant carbons in acyclic systems is a difficult and challenging problem in organic synthesis.¹⁹ General approaches to the construction of remote asymmetric relationships in acyclic systems are rare,^{19d,e} and none of the previous methods allow the choice of preparing both diastereoisomers. We recently applied the methodology illustrated in Scheme V to the stereocontrolled synthesis of the (R^*, R^*) - and (R^*,S^*) -5-hydroxy-2-methylhexanoic acid lactones (pheromone of the carpenter bee) from (E,E)- and (E,Z)-2,4-hexadiene via the synthetic intermediates (R^*, R^*) -21b and (R^*, S^*) -21b, respectively.²⁰ The group $CH(NO_2)(SO_2Ph)$ served as a carboxy anion equivalent in this synthesis.²¹

We were pleased to find that the phenyl sulfinate anion (as NaSO₂Ph) readily displaces the allylic chloro group of the chloroacetates under mild conditions in good yield (71-98%) by using a palladium-catalyzed reaction (eq 2) (cf. (R^*, S^*) - and (R^*, R^*) -21d).²² The chloroacetate 2 from isoprene was trans-



(17) (a) Tsuji, J.; Kataoka, H.; Kobayashi, Y. Tetrahedron Lett. 1981, 22, 2575. (b) Trost, B. M.; Molander, G. A. J. Am. Chem. Soc. 1981, 103, 5969.
(c) Takahashi, T.; Kataoka, H.; Tsuji, J. J. Am. Chem. Soc. 1983, 105, 147. (18) (a) Marino, J. P.; Floyd, D. M. Tetrahedron Lett. 1979, 675. (b)

Marino, J. P.; Hatonaka, H. J. Org. Chem. 1979, 44, 4467. (c) Marino, J. P.; Abe, H. J. Am. Chem. Soc. 1981, 103, 2907. (d) Marino, J. P.; Jaen, J Tetrahedron Lett. 1983, 24, 441. (e) Anderson, R. J. J. Am. Chem. Soc. 1970, 92, 4978

(19) (a) Bartlett, P. A. Tetrahedron 1980, 36, 2. (b) Tetrahedron 1984,
40, 2197–2343 (Symp. No. 16, T. Mukayama, Ed.). (c) Tetrahedron 1984,
40, 1213–1417 (Symp. No. 15, A. I. Meyers, Ed.). (d) Still, W. C.; Darst,
K. P., J. Am. Chem. Soc. 1980, 102, 7387. (e) Fukuyama, T.; Vranesic, B.; Negri, D. P.; Kishi, Y. Tetrahedron Lett. 1978, 2741.

(20) Bäckvall, J. E. Byström, S. E.; Nyström, J. E. *Tetrahedron*, in press. (21) Wade, P. A.; Hinney, H. R.; Amin, N. W.; Vail, P. D.; Morrow, S. D.; Hardinger, S. A.; Saft, M. S. J. Org. Chem. 1981, 46, 765.



formed to the sulfone 22b. The sulfone 22b is another versatile isoprene synthon^{23,24} with one electrophilic and one nucleophilic center and the hydrolized alcohol of 22b has been used in a terpenoid synthesis.²³ The 1-acetoxy-4-(phenylsulfonyl)-2-alkenes and the corresponding 1-acetoxy-4-phosphoryl-2-alkenes, readily available from the corresponding chloroacetates via an Arbuzov reaction, were used for the preparation of the synthetically important 1,3-dienes 23.25



By amination of the chloro group either in a palladium-catalyzed substitution or a classical nucleophilic substitution, a great number of 4-amino-alk-2-envl acetates were prepared.^{26,27} Some of these aminoacetates have been cyclized to pyrrole derivatives.^{26,27}

Discussion

A likely mechanism for the catalytic cycle of the palladiumcatalyzed 1,4-acetoxychlorination is shown for 1,3-cyclohexadiene in Scheme VI). The major inorganic palladium species in solution is most likely monomeric $PdCl_4^{2-,28,29}$ Coordination of the diene followed by a trans acetoxypalladation³⁰ of one of the double bonds gives a $(\pi$ -allyl)palladium intermediate 25. Although monodentate complexes of conjugated dienes with palladium can form at low temperature,³¹ our results are best explained by formation of a cisoid (S)- η^4 -diene complex 24 before the acetoxypalladation takes place (vide infra).³² The regioselectivity for acetate attack at C-1 (terminal) could be a result of a thermodynamic control, since it has been shown that nucleophiles initially attack (η^4 -1,3-di-

(25) Åkermark, B.; Nyström, J. E.; Rein, T.; Bäckvall, J. E.; Helquist, P.; Aslanian, R. Tetrahedron Lett. 1984, 25, 5719.

(26) Genêt, J. P.; Balabane, M.; Bäckvall, J. E.; Nyström, J. E. Tetrahedron Lett. 1983, 24, 2745.

(27) Nyström, J. E.; Bäckvall, J. E., unpublished results.
(28) (a) Goldberg, R. N.; Hepler, L. G. Chem. Rev. 1968, 68, 229. (b) Aguilo, A. Adv. Organometal. Chem. 1967, 5, 327.

(29) (a) Although an ultraviolet study of PdCl₂-LiCl solutions in anhydrous acetic acid (<0.02% water) indicated the presence of dimeric Li₂Pd₂Cle at low chloride concentration, our major species is most likely the monomeric complex for two reasons: (i) the concentration of LiCl is usually 40 times higher then the Pd(II) concentration; (ii) we are working in acetic acid containing 2% water (it is known that Pd(II) exists exclusively as $PdCl_4^{2-}$ in water above a chloride concentration of 0.1 M²⁸). (b) Henry, P. M.; Marks, O. W. *Inorg. Chem.* 1971, *10*, 373.

(30) (a) Henry, P. M.; Ward, G. A. J. Am. Chem. Soc. 1971, 93, 1494.
(b) Andell, O. S.; Bäckvall, J. E. J. Organomet. Chem. 1983, 244, 401.
(31) Donati, F.; Conti, F. Tetrahedron Lett. 1966, 1219.

(32) Treatment of (4-chloro-1,2,3- η^3 -alkenyl)palladium chloride with superacids has in fact been shown to form η^4 complexes related to 24 (L = SbF₅Cl): Lukas, J.; Kramer, P. A. J. Organomet. Chem. 1971, 31, 111.

⁽²²⁾ Attempts to substitute the chloro group in the chloroacetates with the phenyl sulfinate anion in an ordinary S_N2 reaction was less convenient and required higher temperature. Substitution of 2 (NaSO₂Ph, DMF, 60 °C) has been reported.23

⁽²³⁾ Olson, G. L.; Cheung, H. C.; Morgan, K. D.; Neukom, C.; Saucy, G. J. Org. Chem. 1976, 41, 3287.

^{(24) (}a) Cainelli, G.; Cardillo, G. Acc. Chem. Res. 1981, 14, 89. (b) Julia, M.; Verpeux, J. N. Tetrahedron 1983, 39, 3289

Scheme VII



ene)iron complexes at C-2 (internal), followed by rearrangement to the C-1 adduct.³³ Coordination of *p*-benzoquinone followed by an external trans attack by a chloride ion accounts for the overall cis stereochemistry observed. The high stereospecificity shows that no cis migration of chloride from palladium to carbon takes place. Thus, it seems that migration of coordinated chloride from palladium to a coordinated carbon (cf. reductive elimination) is an unfavored process as suggested previously.³⁴

It is likely that *p*-benzoquinone coordinates to palladium and in this way facilitates the electron-transfer process. Moiseev and co-workers³⁵ reported that treatment of $(\pi$ -allyl)palladium chloride with *p*-benzoquinone and aqueous HCl gave allyl chloride, and the kinetics of the reaction were consistent with a coordination of the benzoquinone. Stable *p*-benzoquinone complexes of nickel(0), palladium(0), and platinum(0) have been described,³⁶ and protonation of a (*p*-benzoquinone)nickel(0) complex was shown to yield hydroquinone and nickel(II).^{36c}

The first-order dependence of *p*-benzoquinone on the reaction rate (Figure 1) supports a coordination of the quinone. Because of the strongly coordinated chloride ligands, a large excess of *p*-benzoquinone (30-fold to palladium) is required before the linear increase of the rate is declining.^{7b}

The high product selectivity for chloroacetate over diacetate and dichloride is remarkable. From a statistical point of view, one would expect the latter products to form to some extent. The regiochemistry observed for isoprene (Table I, entry 2) supports that the first nucleophile introduced is acetate, since isoprene is known³⁷ to react with palladium chloride in acetic acid to give (4-acetoxy-2-methyl-1,2,3,- η^3 -butenyl)palladium chloride dimer. The question is, why does acetate win in the first step when chloride wins in the second step. An explanation for this unusual selectivity is offered in Scheme VII. The chloro complex 26, if initially formed, is expected to solvolyze rapidly in acetic acid (via diene complex) to afford the more stable complex 27.33,38 External chloride attack on 27 would give the observed chloroacetate 1 as the sole product. Further support for a chloro complex 26 as a kinetic intermediate is provided by the formation of 1,4-dichloro-2-butene (28) when p-benzoquinone was replaced by the more rapid oxidant isoamyl nitrite.^{27,39} It is interesting to note that a chlorometalation adduct was detected as the initial intermediate in oxymetalation of ethene when using platinum chloride.40

D.; Shaw, B. L. J. Chem. Soc. 1963, 4806. (b) Åkermark, B.; Bäckvall, J.
E.; Löwenborg, A.; Zetterberg, K. J. Organomet. Chem. 1979, 166, C33. (39) The dichloride 28 was a mixture of the E and Z isomer in a ratio of

1:1.4, which reflects a competition between syn-anti isomerization and nucleophilic attack similar to that shown in Scheme VIII. Scheme VIII



The variation of the E/Z ratio of the acyclic chloroacetates with the substrates and reaction conditions can be explained if one assumes that a (S)- η^4 -coordinated cisoid diene complex 29 is formed initially (Scheme VIII). Acetate addition to the complex 29 would give mainly complex 30. A syn-anti isomerization would produce complex 31, which is considerably more stable than 30 if $R^2 = H$. When $R^1 = R^2 = H$, the isomerization of 30 to 31 should be rapid and the E product is expected as the major isomer, which is observed for butadiene (E/Z = 9/1). On the other hand, for 2,3-dimethylbutadiene, the isomerization of complex 30 ($R^1 = R^2 = CH_3$) to 31 is small. This explains the predominant formation of Z product from this diene (E/Z =1/10). For isoprene the rate of isomerization of 30 (R¹ = CH₃, $R^2 = H$) to 31 falls in between, which would explain the observed E/Z ratio of = 3.6/1. Also, the observed increase of the acetoxychlorination rate by a factor of 2 on changing one double bond from Z to E is consistent with a cisoid η^4 -coordinated diene (cf. Table IIa and b). Such a coordination would be more difficult for a (Z)-diene than for an (E)-diene.

The regioselectivity obtained for unsymmetrically substituted dienes is nicely explained by considering the π -allyl intermediates. Thus, a substituent in the 2-position of the 1,3-diene will lead almost exclusively to the more stable (π -allyl)palladium complex.³⁷ Furthermore, the low regioselectivity obtained for 1-substituted 1,3-dienes is consistent with the observation that reaction of 1,3-pentadiene with Pd(II) in methanol produces two isomeric π -allyl complexes.³⁷a

The highly stereospecific palladium(0)-catalyzed substitution reactions of the allylic chloride that take place with complete retention proceed via the mechanism outlined in eq 3 in analogy with the palladium(0)-catalyzed nucleophilic substitution reactions of allylic acetate. The reason for the high stereospecificity

compared to the reaction of allylic acetate is that the coordinated chloro group has no tendecy to migrate to the allyl group (vide supra), which is the case for a coordinated acetoxy group.^{7a,b,41,42} Such a cis migration will lead to isomerization of the starting allylic substrate.^{41,43}

It is interesting to note that the acyclic (π -allyl)palladium complexes 32, which occur as intermediates in the oxidation of 2,4-hexadienes as well as in the palladium(0)-catalyzed substitutions, are configurationally stable. Diastereomeric complexes related to 32 have been shown to be configurationally stable,⁴⁴ and recently, the optically active (π -allyl)palladium complex 33 was prepared and shown not to racemize.^{45,46}

⁽³³⁾ Semmelhack, M. F.; Le, H. T. M. J. Am. Chem. Soc. 1984, 106, 2715.

^{(34) (}a) Bäckvall, J. E.; Björkman, E. E. J. Chem. Soc., Chem. Commun. 1982, 693. (b) Bäckvall, J. E.; Björkman, E. E.; Petterson, L.; Siegbahn, P. J. Am. Chem. Soc. 1984, 106, 4369.

J. Am. Chem. Soc. 1984, 106, 4369.
 (35) Satsko, N. G.; Below, A. P.; Moiseev, I. I. Kinet. Catal. 1972, 13, 892;
 Chem. Abstr. 1973, 78, 83584c.

^{(36) (}a) Hiramatsu, M.; Nakano, H.; Fujinami, T.; Sakai, S. J. Organomet. Chem. 1982, 236, 131. Hiramatsu, M.; Shiozaki, K.; Fujinami, T.; Sakai, S. Ibid. 1983, 246, 303. (b) Chetcuti, M. J.; Howard, J. A. K.; Pfeffer, M.; Spencer, J. L.; Stone, F. G. A. J. Chem. Soc., Dalton Trans. 1981, 276. Chetcuti, M. J.; Herbert, J. A.; Howard, J. A. K. Pfeffer, M.; Spencer, J. L.; Stone, F. G. A., J. Chem. 1981, 284. (c) Uhlig, E.; Fischer, R.; Kirmse, R. J. Organomet. Chem. 1982, 239, 385.

^{(37) (}a) Rowe, J. M.; White, D. A. J. Chem. Soc. A 1967, 1451. (b)
Levanda, O. G.; Pek, G. I.; Moiseev, I. I. Zh. Org. Khim. 1971, 7, 217.
(38) Such exchange reactions of the chloro group in (4-chloro-1,2,3-n³-alkenyl)palladium complexes related to 26 are well-known: (a) Robinson, S.

⁽⁴⁰⁾ Halpern, J.; Jewsbury, R. A. J. Organomet. Chem. 1979, 181, 223.
(41) (a) Trost, B. M.; Verhoeven, T. R.; Fortunak, J. M. Tetrahedron Lett.
1979, 2301. (b) Trost, B. M.; Verhoeven, T. R. J. Am. Chem. Soc. 1980, 102, 4730.

⁽⁴²⁾ Bäckvall, J. E.; Nordberg, R. E.; Björkman, E. E.; Moberg, C. J. Chem. Soc., Chem. Commun. 1980, 943.

⁽⁴³⁾ Nordberg, R. E.; Bäckvall, J. E. J. Organomet. Chem., in press.
(44) Lukas, J.; van Leeuwen, P. W. N. M.; Volger, H. C.; Kouwenhoven,
A. P. J. Organomet. Chem. 1973, 47, 153.

 ^{(45) (}a) Hayashi, T.; Hagihara, T.; Konishi, M.; Kumada, M. J. Am.
 Chem. Soc. 1983, 105, 7767. (b) Hayashi, T.; Konishi, M.; Kumada, M. J.
 Chem. Soc., Chem. Commun. 1984, 107. (c) Ibid. 1983, 736.



It should be mentioned that the sequential nucleophilic substitution of the chloroacetates outlined in Scheme III can be performed in one pot. This was used for a highly selective sequential dialkylation of chloroacetate 2 by using two different carbon nucleophiles.¹³ Related sequential palladium-catalyzed nucleophilic substitutions were recently performed on (Z)-4acetoxybut-2-enyl diethyl phosphate47 and 2-cyclopentene-1,4-diol dicarboxylates.⁴⁸ A discrete two step alkylation has also been used to prepare vinylcyclopropanes from 2-alkene-1,4-diol derivatives.49

Concluding Remark

The present work has shown that conjugated dienes can be selectively oxidized to the synthetically useful 1-acetoxy-4chloro-2-alkenes under mild conditions (room temperature) by using a Pd(II) salt as the catalyst (0.2-7.5 mol %). The mechanism of the acetoxychlorination is now well understood. It proceeds via a (4-acetoxy-1,2,3- η^3 -allyl)palladium intermediate, followed by an oxidation-induced attack by chloride.

The reaction is 1,4-regioselective in most cases. Furthermore, the acetoxychlorination is highly stereospecific in both cyclic and acyclic systems. This constitutes a useful internal 1,4-asymmetric induction. Combined with the fact that the chloro group can be substituted with either retention or inversion, the acetoxychlorination approach allows a complete control of the 1.4-relative stereochemistry. Such a dual control of the 1,4-relative stereochemistry with the unique choice of 1,4-functionality is previously unprecedented in organic chemistry.

Experimental Section

Infrared spectra were recorded on a Perkin-Elmer 257 spectrophotometer. NMR spectra were obtained with a Bruker WP 200 FT spectrometer, ¹H NMR at 200 MHz, and ¹³C NMR at 50.3 MHz. ¹³C NMR multiplicities were obtained by proton off-resonance decoupling at 1650 Hz. Chemical shifts are reported in δ units, parts per million (ppm) downfield from tetramethylsilane, Me₄Si. High-pressure liquid chromatography (HPLC) was performed on a Waters M-45 instrument with a micro-Porasil column (silica, 10- μ m packing, 0.4 × 30 cm) and a differential refractometer as detector. The slow diene additions were performed with a Sage Instruments Model 355 syringe pump. Bulb-tobulb distillations were performed with a Büchi Kugelrhor apparatus.

Butadiene, isoprene, 2,3-dimethylbutadiene, (E)-2-methyl-1,3-pentadiene, (E)- and (Z)-1,3-pentadiene, 4-methyl-1,3-pentadiene, 2,5-dimethyl-2,4-hexadiene, (E,E)- and (E,Z)-2,4-hexadiene, 1,3-cyclohexadiene, 1,3-cycloheptadiene, and 1,3-cyclooctadiene were purchased from FLUKA AG and were distilled before use. (E)-6-Phenyl-1,3-hexadiene,⁵⁰ ethyl (E)-4,6-heptadienoate,⁵¹ and 2-methyl- and 5-methyl-1,3-cyclohexadiene^{7b,52} were prepared according to literature procedures. 6-Acetoxy-1,3-cycloheptadiene was prepared by acetylation (Ac₂O, 4-(dimethylamino)pyridine, and Et₃N) of 6-hydroxy-1,3-cycloheptadieneъ,53 Palladium acetate was purchased from Engelhard Industries. Lithium acetate dihydrate, lithium chloride, p-benzoquinone, and sodium benzenesulfinate were 99% grade and were used without further purification. Tetrahydrofuran (THF) was distilled from a deep blue solution of potassium/benzophenone. Acetonitrile was distilled over CaH2 and stored over molecular sieves (4 Å).

Tetrakis(triphenylphosphine)palladium (Pd(PPh₃)₄). To a homogeneous, brown, stirred solution of Pd(PhCN)₂Cl₂ (3.07 g, 8.0 mmol) in

 (49) Genêt, J. P.; Piau, F. J. Org. Chem. 1981, 46, 2414.
 (50) Reich, H. J.; Wollowitz, S. J. Am. Chem. Soc. 1982, 104, 7051. (51) Ethyl (E)-4,6-heptadienoate was a gift from Dr. B. Nilsson, Organic Chemistry, Chemical Center, S-220 07 Lund, Sweden.

degassed acetone (80 mL) under nitrogen at 0 °C was added butadiene (1.8 g, 33 mmol), which resulted in formation of a yellow precipitate. The cooling bath was removed, and the reaction mixture was stirred at room temperature for 2 h. The mixture was cooled to 0 °C, and diethylamine (6.7 mL, 64 mmol) was added. Triphenylphosphine (8.4 g, 32 mmol) was immediately added and a precipitation of the yellow Pd(0) complex was observed. After the mixture was stirred at 0 °C for 1 h, the crystals were filtered off under nitrogen by using a glass filter (Schlenk apparatus). The crystals were washed under nitrogen with ice-cold degassed solvents: acetone (2 \times 20 mL), acetone/water 1/1 mixture (2 × 20 mL), and finally acetone (2 × 20 mL). The complex was dried in vacuo to give 6.41 g (69%) of a light yellow powder. The palladium complex can be stored under nitrogen in a freezer for months.

General Procedures for Palladium(II)-Catalyzed 1,4-Acetoxychlorination of Conjugated Dienes. Unless otherwise noted, all reactions were performed at 25 °C in acetic acid using Pd(OAc)₂ as catalyst (5 mol % for the cyclic dienes and 7.5 mol % for the acyclic dienes). The amount of diene used was 0.25-0.30 mmol/mL of acetic acid. Two equivalents of p-benzoquinone, LiCl, and LiOAc $\cdot 2H_2O$ were used.

Method A. The diene was added to the pentane phase of a biphasic pentane/acetic acid reaction mixture (1.5-2.0/1, pentane/HOAc, v/v).

Method B. The diene was added to a monophasic acetic acid reaction mixture. The total reaction time and the diene addition time are shown in Table I. The longer addition times (>3 h) of the diene were accomplished by using a syringe pump.

The results are shown in Table I. The yields refer to isolated yield of distilled product unless otherwise stated. Complete, representative experimental procedures are described for the preparation of 1 (method A and B), 11 (method B), and 12 (method A). For a purification procedure, see under 1, 6, or 7.

(E)-1-Acetoxy-4-chloro-2-butene (1) was prepared by either method A (81%) or method B (78%).

Method A. To butadiene (5.40 g, 100 mmol) dissolved in pentane (800 mL) at 0 °C was added an acetic acid (400 mL) solution of Pd(OAc)₂ (1.68 g, 7.5 mmol), LiCl (8.4 g, 200 mmol), LiOAc·2H₂O (20.4 g, 200 mmol), and p-benzoquinone (21.6 g, 200 mmol). After the solution was stirred for 26 h at 25 °C, brine (300 mL) and ether (160 mL) were added, and stirring was continued for another 5 min. The organic phase was separated and the aqueous phase filtered and extracted with pentane/ether (3 \times 300 mL, 80/20). The combined organic phases were washed with water (100 + 50 mL), saturated aqueous K_2CO_3 (3 × 100 mL), and 2 M NaOH (3 \times 100 mL). The alkaline aqueous phases were back-extracted with pentane/ether $(2 \times 100 \text{ mL}, 80/20)$. The combined organic phases were washed with brine and dried (MgSO₄). The solvent was distilled off at ambient pressure and finally rotary-evaporated at reduced pressure, giving a crude product (≈ 15 g) which was then distilled (10 mmHg, 70-90 °C), yielding 12.0 g (81%) of 1 as a light yellow liquid consisting of (E)-1 (82%), (Z)-1 (9%), and 4-acetoxy-3-chloro-1-butene (1') (9%) according to ¹H NMR. The chloroacetate was contaminated with $\approx 1\%$ of 5,8-dihydronaphthoquinone: IR (neat) 2940, 1740, 1380, 1360, 1230, 1025, 970 cm⁻¹. Anal. Calcd for $C_6H_9ClO_2$: C, 48.50; H, 6.11. Found: C, 48.68; H, 5.99.

(E)-1: ¹H NMR (CDCl₃) δ 5.9 (br s, 2 H, CH=CH), 4.59 (br s, 2 H, CH_2OAc), 4.08 (br s, 2 H, CH_2Cl), 2.09 (s, 3 H, AcO); ¹³C NMR (CDCl₃) & 170.4 (s, CH₃COO), 129.7 (d, one of HC=CH), 128.5 (d, one of HC==CH), 63.56 (t, CH2OAc), 43.82 (t, CH2Cl), 20.79 (q, $CH_1COO)$

(Z)-1 (distinguishable peaks in mixture with (E)-1 and 1'): ¹H NMR $(CDCl_3) \delta 4.68$ (br d, J = 6 Hz, 2 H, CH_2OAc), 4.15 (br d, J = 6 Hz, 2 H, CH₂Cl).

1' (distinguishable peaks in mixture with (E)- and (Z)-1): ¹H NMR $(CDCl_3) \delta 5.4 (d, J = 17 Hz, 1 H, trans in CH_2=CH), 5.28 (d, J = 10$ Hz, 1 H, cis in CH2=CH), 4.45-4.20 (AB part of ABX, 2 H, CH-CH₂OAc).

5,8-Dihydronaphthoquinone (distinguishable peaks in mixture with 1): ¹H NMR (CDCl₃) δ 6.7 (s, 2 H, C—CH=CH-C=O), 3.08 (s, 4 H, CH₂).

Purification of 1. 1 (9 g containing ≈ 0.2 g of 5,8-dihydronaphthoquinone) dissolved in ether (150 mL) was stirred with an aqueous 2 M NaOH solution saturated with NaBH₄ (10 mL) until the yellow color had disappeared (15 min). The aqueous phase (brown) was discarded and the etheral phase was washed with 2 M NaOH (5 mL) and dried (MgSO₄). Concentration at ambient pressure and finally rotary evaporation afforded 8.5 g of 1 as a colorless liquid which was free from 5,8-dihydronaphthoquinone (<0.1%) according to ¹H NMR.

Method B. To a well-stirred solution of Pd(OAc)₂ (90 mg, 0.4 mmol), LiCl (336 mg, 8 mmol), LiOAc·2H₂O (816 mg, 8 mmol), and p-benzoquinone (950 mg, 8.8 mmol) in acetic acid (14 mL) at 25 °C was slowly

^{(46) (}a) The configuration of 1,3-disubstituted (π -allyl)palladium complexes such as 32 and 33 cannot change due to repeated syn-anti isomeriza-tions (via σ -allyl).^{7b,46b} (b) Faller, J. W.; Thomsen, M. E.; Mattina, M. J. J. Am. Chem. Soc. 1971, 93, 2642.

⁽⁴⁷⁾ Tanigawa, Y.; Nishimura, K.; Kawasaki, A.; Murahashi, S. I. Tet-rahedron Lett. 1982, 23, 5549.

⁽⁴⁸⁾ Valpey, R. S.; Miller, D. J.; Estes, J. M.; Godleski, S. A. J. Org. Chem. 1982, 47, 4717.

⁽⁵²⁾ Zelinski, N.; Gorsky, A. Chem. Ber. 1908, 41, 2630.

⁽⁵³⁾ Schuster, D. J.; Palmer, J. M.; Dickerson, S. C. J. Org. Chem. 1966, 31, 4281.

added butadiene dissolved in acetic acid ($1.94 \text{ g} \times 1.88 \text{ m} = 3.64 \text{ mmol}$) with a syringe (using a syringe pump) over 16 h. After additional stirring for 4 h, the reaction mixture was diluted with water (8 mL) and pentane/ether (20 mL, 80/20). The workup was the same as in method A. Bulb-to-bulb distillation afforded 422 mg (78%) of 1 with the same composition as that obtained from method A.

1-Acetoxy-4-chloro-3-methyl-2-butene (2) was prepared from isoprene by method A (76%), 22 (0) h, or method B (74%), 26 (16) h.¹³ The product consisted of (E)-2 (69%), (Z)-2 (19%), 4-acetoxy-3-chloro-2methyl-1-butene (2') (4%), and 1-acetoxy-4-chloro-2-methyl-2-butene (20) (8%).

Mixture of (E)-2 and 2' was prepared from isoprene by method A (65%), 14 (0) h, but benzoquinone was replaced by MnO_2 . The product consisted of 2 (>95% E) (53%), 2' (34%), 20 (7%), and 4-acetoxy-3-chloro-3-methyl-1-butene (20') (6%) according to ¹H NMR. Compounds (E)-2, (Z)-2 and 2' have previously been fully characterized.¹³

20' (distinguishable peaks in mixture with **2**'): ¹H NMR (CDCl₃) δ 6.05 (dd, J = 17.2, 10.7 Hz, 1 H, CH₂=-CH), 5.37 (d, J = 17.2 Hz, 1 H, trans in CH₂=-CH), 5.22 (d, J = 10.7 Hz, 1 H, cis in CH₂=-CH), 5.42 kH, control H, control H,

4.23 (br s, 2 H, CH_2OAc), 2.11 (s, 3 H, OAc), 1.68 (s, 3 H, CH_3 —C). 20: ¹H NMR ($CDCl_3$) δ 5.65 (br t, J = 8 Hz, 1 H, CH=C), 4.63 (s, 2 H, CH_2OAc), 4.15 (d, J = 8 Hz, 2 H, CH_2Cl), 2.08 (s, 3 H, OAc), 1.83 (br s, 3 H, CH_3 —C=C).

(Z)-1-Acetoxy-4-chloro-2,3-dimethyl-2-butene ((Z)-3) was prepared from 2,3-dimethylbutadiene by method A (44%), 20 (0) h, or method B (45%), 32 (28) h. Bulb-to-bulb distillation (100 °C, 1 mmHg) of the crude product afforded 3 consisting of (Z)-3 (83%), (E)-3 (8%),⁵⁴ and 4-acetoxy-3-chloro-2,3-dimethyl-1-butene (3') (9%) according to ¹H NMR analysis.

(Z)-3: ¹H NMR (CDCl₃) δ 4.64 (s, 2 H, CH₂OAc), 4.17 (s, 2 H, CH₂Cl), 2.08 (s, 3 H, OAc), 1.84 (s, 3 H, CH₃ β to Cl), 1.78 (s, 3 H, CH₃ β to OAc); (C₆D₆) δ 4.45 (s, 2 H, CH₂OAc), 3.87 (s, 2 H, CH₂Cl), 1.64 (s, 3 H, OAc), 1.50 (s, 3 H, CH₃ β to Cl), 1.46 (s, 3 H, CH₃ β to OAc); ¹³C NMR (CDCl₃) δ 170.9 (s, CH₃COO), 131.6 (s), 130.3 (s), 64.30 (t, CH₂OAc), 45.45 (t, CH₂Cl), 20.92 (q, CH₃COO), 17.59 (q), 17.53 (q).

(*E*)-3 (distinguishable peaks in mixture with (*Z*)-3): ¹H NMR (CD-Cl₃) δ 4.61 (s, 2 H, CH₂OAc), 4.10 (s, 2 H, CH₂Cl); (C₆D₆) δ 4.39 (s, 2 H, CH₂OAc), 3.62 (s, 2 H, CH₂Cl).

Treatment of 3 with the shift reagent $Eu(fod)_3$ revealed information about the double bond configuration. Thus, the methylene protons are more shifted for the major isomer than for the minor isomer, while the reversed is true for the vinylic methyls.

3' (distinguishable peaks in mixture with (*E*)- and (*Z*)-3): ¹H NMR (CDCl₃) δ 5.12 (m, 1 H, one of CH₂==C), 5.02 (m, 1 H, one of CH₂== C), 4.4-4.25 (AB spectrum, J_{AB} = 11.5 Hz, 2 H, CH₂OAc), 1.91 (d, J = 1.5 Hz, 3 H, CH₃-C==C), 1.60 (s, 3 H, CH₃-C=Cl); ¹³C NMR (CDCl₃) δ 114.1 (t, CH₂=C).

For analytical purpose, 3 (0.5 mmol) was converted to 1,4-diacetoxy-2,3-dimethyl-2-butene (34) by reaction with KOAc (196 mg, 2 mmol) in Me₂SO (2 mL) at 70 °C for 2 h. After the solution cooled, water (2 mL) was added and the aqueous phase was extracted with pentane (4×4 mL). The combined organic phases were washed with water (1 mL) and dried (MgSO₄). Evaporation of the solvent afforded 94 mg (89%) of 34 as a mixture of Z/E isomers (10/1).⁵⁴

(Z)-34: ¹H NMR (CDCl₃) δ 4.65 (s, 4 H, CH₂OAc), 2.06 (s, 6 H, OAc), 1.77 (s, 6 H, CH₃).

(É)-34: ¹H NMR (CDCl₃) δ 4.62 (s, 4 H, CH₂OAc), 2.08 (s, 6 H, OAc), 1.79 (s, 6 H, CH₃).

4-Acetoxy-1-chloro-2-methyl-2-pentene (4) was prepared from (*E*)-2methyl-1,3-pentadiene by method B (35%, E/Z = 2.5), 49 (45) h, containing $\approx 7\%$ of the regioisomer 5-acetoxy-4-chloro-4-methyl-2-pentene (35): IR (neat) 2980, 1770, 1450, 1370, 1240, 1050 cm⁻¹. Anal. Calcd for C₈H₁₃ClO₂: C, 54.40; H, 7.42. Found: C, 54.40; H, 7.42.

(E)-4: ¹H NMR (CDCl₃) δ 5.55 (dq, A part of ABX₃ spectrum, $J_{AB} = 8$, $J_{AX} = 6.1$ Hz, 1 H, CH₃—CH(OAc)—CH=C), 5.51 (br d, B part of ABX₃ spectrum, $J_{AB} = 8$, $J_{BX} = 0$ Hz, CH₃CH(OAc)—CH=C), 3.99 (s, 2 H, CH₂Cl), 2.03 (s, 3 H, OAc), 1.83 (d, J = 1.2 Hz, 3 H, CH₃—C=C), 1.29 (d, $J_{AX} = 6.1$ Hz, 3 H, CH₃CHO); ¹³C NMR (CDCl₃) δ 170.2 (s, CH₃COO), 134.8 (s, C=CH), 129.5 (d, C=CH), 67.52 (d, CHOAc), 50.99 (t, CH₂Cl), 21.27 (q, CH₃C), 20.36 (q, CH₃COO), 14.73 (q, CH₃—C=C); HPLC, $V_t/V_0 = 2.6$ (EtOAc/hexane = 4/96).

(Z)-4: ¹H NMR (CDCl₃) δ 5.61 (dq, A part of ABX₃ spectrum, $J_{AB} = 9$, $J_{AX} = 6.2$ Hz, 1 H, CH₃—CH(OAc)—CH=C), 5.35 (br d, B part of ABX₃ spectrum, $J_{AB} = 9$, $J_{BX} = 0$ Hz, 1 H, CH₃CH(OAc)—CH=C), 4.36 (d, J = 11 Hz, 1 H, one of CH₂Cl), 3.94 (d, J = 11 Hz, 1 H, one of CH₂Cl), 2.02 (s, 3 H, OAc), 1.86 (d, J = 1.4 Hz, 3 H, CH₃—C=C), 1.31 (d, J = 6.2 Hz, 3 H); ¹³C NMR (CDCl₃) δ 170.2 (s, CH₃COO),

135 (s, C=CH), 130.1 (d, C=CH), 66.95 (d, CHOAc), 43.12 (t, CH₂Cl), 21.65 (q, CH₃-C=C), 20.78 (q, CH₃COO), 20.55 (q, CH₃CH(OAc)); HPLC, $V_r/V_o = 2.3$ (EtOAc/hexane = 4/96).

35 (distinguishable peaks in mixture with 4): ¹H NMR (CDCl₃) δ 4.56 (s, 2 H, CH₂OAc).

Mixture of 5a and 5b was prepared from (E)-1,3-pentadiene or (Z)-1,3-pentadiene by method B (51%), 20 (16) h. The ratio 5a/5b = 1.5 was established by ¹H NMR.

5a: ¹H NMR (CDCl₃) δ 5.9–5.8 (m, J = 15.6 Hz, 2 H, CH=CH), 5.38 (quin, J = 6 Hz, 1 H, CHOAc), 4.05 (d, J = 6 Hz, 2 H, CH₂Cl), 2.05 (s, 3 H, OAc), 1.33 (d, J = 6.5 Hz, 3 H, CH₃CHOAc); ¹³C NMR (CDCl₃) δ 170.1 (s, OCOCH₃), 133.9 (d, ClCH₂—CH=CH), 127.8 (d, CH=CH—CHOAc), 69.62 (d, CHOAc), 44.05 (t, CH₂Cl), 21.24 (q, OCOCH₃), 19.99 (q, CH₃CHOAc).

5b: ¹H NMR (CDCl₃) δ 5.9–5.8 (m, 2 H, HC=CH), 4.65 (quin, J = 6.5 Hz, 1 H, CHCl), 4.58 (d, J = 4.8 Hz, 2 H, CH₂OAc), 2.09 (s, 3 H, OAc), 1.61 (d, J = 6.7 Hz, 3 H, CH₃CHCl); ¹³C NMR (CDCl₃) 170.1 (s, OCOCH₃), 135.7 (d, ClCH-CH=CH), 127.4 (d, AcOCH₂-CH=CH), 63.65 (t, CH₂OAc), 56.54 (d, CHCl), 24.90 (q, CH₃CHCl), 20.87 (q, OCOCH₃).

Anal. Calcd for $C_7H_{11}ClO_2$: C, 51.70; H, 6.82. Found (for the mixture **5a** and **5b**): C, 51.78; H, 6.68. IR (neat) 2980, 1730, 1440, 1370, 1230, 1140, 1040, 960, 910 cm⁻¹.

Mixture of 6a and 6b was prepared from (E)-6-phenyl-1,3-hexadiene (1.7 mmol) by method B (36%), 21 (14) h. After the usual extraction procedure, the combined organic phases were washed with saturated aqueous Na₂CO₃ (2 × 5 mL) and 2 M NaOH (5 mL) and then stirred for 1.5 h with 2 M NaOH (5 mL) which was saturated with NaBH₄. In this way the quinoid Diels-Alder adduct was reduced according to HPLC, $V_r/V_o = 2.6$ (EtOAc/hexane = 2.5/97.5). The organic phase was washed with brine (3 mL), dried (MgSO₄), and eluted through a silica column (5 × 2 cm) with EtOAc/hexane (1/1). Evaporation of the solvent afforded 155 mg (36%) of 6 consisting of 6a (44%), 6b (43%), and 1-acetoxy-2-chloro-6-phenyl-3-hexene (6') (13%). For analytical purposes, 6a and 6b were separated by preparative HPLC.

6a: HPLC, $V_r/V_o = 2.8$ (EtOAc/hexane = 2.5/97.5); ¹H NMR (CDCl₃) δ 7.35-7.1 (m, 5 H, Ph), 5.86 (dt, J = 15.2 and 5.4 Hz, 1 H, CICH₂—CH=CH), 5.77 (dd, J = 15.2 and 5.4 Hz, 1 H, CH=CH—CHOAc), 5.29 (br q, J = 6 Hz, 1 H, CHOAc), 4.05 (d, J = 5.4 Hz, 2 H, CH₂Cl), 2.65 (t, J = 7.3 Hz, 2 H, CH₂Ph), 2.06 (s, 3 H, OAc), 1.97 (br q, J = 8 Hz, 2 H, CH₂CH₂Ph); ¹³C NMR (CDCl₃) (assigned peaks in mixture with **6b**) δ 72.85 (d, CHOAc), 43.96 (t, CH₂Cl).

6b: HPLC, $V_r/V_o = 3.0$ (EtOAc/hexane = 2.5/97.5); ¹H NMR (CDCl₃) δ 7.35-7.1 (m, 5 H, Ph), 5.95-5.7 (m, 2 H, CH=CH), 4.57 (d, J = 4.5 Hz, 2 H, CH₂OAc), 4.35 (br q, J = 6.7 Hz, 1 H, CHCl), 2.80-2.75 (AB part of ABX₂, $J_{AB} = 14$, $J_{AX} = 7$ Hz, 2 H, CH₂CH₂Ph), 2.12 (br q, J = 7 Hz, 2 H CH₂Ph), 2.08 (s, 3 H, OAc), ¹³C NMR (CDCl₃) (assigned peaks in mixture with **6a**) δ 60.63 (t, CH₂OAc), 60.76 (d, CHCl).

6': HPLC, $V_r/V_0 \approx 2.4$ (EtOAc/hexane $\approx 2.5/97.5$); ¹H NMR (CDCl₃) (distinguishable peaks in mixture with **6a** and **6b**) δ 4.2 (AB part of ABX, 2 H, CH₂OAc), 2.45–2.30 (m, 2 H, CH=CH-CH₂).

Anal. Calcd for $C_{14}H_{17}ClO_2$: C, 66.53; H, 6.78. Found (for the mixture **6a**, **6b**, and **6'**): C, 66.65; H, 6.76. IR (neat) 3020, 2920, 1740, 1370, 1230, 1025, 965, 750, 700 cm⁻¹.

Mixture of 7a and 7b was prepared from ethyl (E)-4,6-heptadienoate (2 mmol) by method B (58%), 42 (40) h. A slightly modified workup procedure was used. After addition of brine (6 mL) to the reaction mixture, the aqueous phase was extracted with ether/pentane = 40/60 (4 × 20 mL). The combined organic phase was washed with H₂O (2 mL which was saturated with solid K₂CO₃ in the separating funnel) and 2 M NaOH (2 mL). The organic phase was stirred with 2 M NaOH (2 mL). The organic phase was stirred with 2 M NaOH (5 mL saturated with NaBH₄) until the Diels-Alder adduct HPLC peak ($V_r/V_o = 2$, EtOAc/hexane = 10/90) had disappeared (1 h). After washing with 2 M NaOH (2 mL) and brine (5 mL), the organic phase was dried (MgSO₄) and the solvents were evaporated, yielding 278 mg (58%) of 7 consisting of 7a (44%), 7b (36%), 1-acetoxy-2-chloro-6-carbethoxy-3-hexene (7') (11%), and an unidentified compound (9%) (HPLC $V_r/V_o = 2.9$). The composition was determined by HPLC and ¹H NMR of the mixture.

7a: HPLC, $V_r/V_o = 3.3$ (EtOAc/hexane = 10/90); ¹H NMR (CD-Cl₃) δ 5.84 (dt, J = 15.6, 6 Hz, 1 H, ClCH₂—CH=CH), 5.75 (dd, J = 15.6, 6.2 Hz, 1 H, CH=CH—CHOAc), 5.31 (br q, J = 6.2 Hz, 1 H, CH=CH—CHOAc), 5.31 (br q, J = 6.2 Hz, 1 H, CHOAc), 4.34 (q, J = 7.2 Hz, 2 H, OCH₂), 4.04 (d, J = 6 Hz, 2 H, CH₂Cl), 2.35 (t, J = 7.3 Hz, 2 H, CH₂CO), 2.08 (s, 3 H, OAc), 1.98 (d, J = 6-7 Hz, 2 H, ACOCHCH₂), 1.26 (t, J = 7.2 Hz, 3 H, OCH₂CH₃); ¹³C NMR (CDCl₃) δ 172.5 (s, COOEt), 170.0 (s, OCOCH₃), 133.6 (d, ClCH₂—CH=CH), 127.0 (d, CH—CH=CHOAc), 72.30 (d, CHOAc), 60.57 (t, OCH₂CH₃), 43.82 (t, CH₂Cl), 30.98 (t, CH₂CCO), 29.23 (t, CH₂CO), 21.06 (q, OCOCH₃), 14.21 (q, OCH₂CH₃).

^{(54) (}a) The 60 MHz-spectra of 3 and 34 have been published but the shifts differ slightly from ours. (b) Ganem, B.; Small, V. R., Jr. J. Org. Chem. 1974, 39, 3728.

7b: HPLC, $V_r/V_o = 3.4$ (EtOAc/hexane = 10/90); ¹H NMR (CD-Cl₃) $\delta 5.84$ (m, 2 H, CH=CH), 4,58 (d, J = 4 Hz, 2 H, CH₂OAc), 4.46 (m 1 H, CHCl), 4.34 (q, J = 7.2 Hz, 2 H, OCH₂CH₃), 2.49 (t, J = 7.5Hz, 2 H, CH₂CO), 2.1 (m, 2 H, CH₂CH₂CO), 2.08 (s, 3 H, OAc); ¹³C NMR (CDCl₃) δ 172.5 (s, COOEt), 170.0 (s, OCOCH₃), 131.9 (d, ClCH-CH=CH), 128.8 (d, CH=CH-CH₂OAc), 63.51 (t, CH₂OAc), 60.57 (t, OCH₂CH₃), 60.57 (d, CHCl), 33.24 (t, CH₂CH₂CO), 29.90 (t, CH₂CO), 20.85 (q, OCOCH₃), 14.21 (q, OCH₂CH₃).

7': HPLC, $V_r/V_o = 3.1$ (EtOAc/hexane = 10/90); ¹H NMR (CD-Cl₃) (distinguishable peaks in mixture with 7a and 7b) δ 4.22 (AB part of ABX spectrum, $J_{AB} = 11.5$, $J_{AX} = 6$, $J_{BX} = 7$ Hz, 2 H, ClCHCH₂OAc); ¹³C NMR (CDCl₃) (distinguishable peaks in mixture with 7a and 7b) δ 131.4 (d, ClCH—CH=CH), 127.8 (d, CH=CH— CH₂), 66.95 (t, CH₂OAc), 58.5 (d, CHCl), 33.4 (t), 27.35 (t).

Anal. Calcd for $C_{11}H_{17}ClO_4$: C, 53.12; H, 6.89. Found (for a mixture of **7a** and **7b** which was obtained by preparative HPLC): C, 53.30; H, 6.93. IR (neat) 2940, 1735, 1370, 1230, 1025, 970 cm⁻¹.

(E)-(R^* , R^*)-2-Acetoxy-5-chloro-3-hexene ((R^* , R^*)-10) was prepared from (E,E)-2,4-hexadiene (4 mmol) by method A (55%), 24 (18) h, pentane (24 mL)/acetic acid (16 mL). From the ¹H NMR spectrum, the diastereomeric purity was established to be >95% R^* , R^* . On conversion to the diacetate (see under preparation of 3) and by subsequent HPLC analysis, it was found that $E/Z \approx 99/1$.

(R^*, R^*)-10: ¹H NMR (CDCl₃) δ 5.82 (dd, J = 15.4, 6.6 Hz, 1 H, CH=CH-CHCl), 5.72 (dd, J = 15.4, 4.9 Hz, 1 H, CH=CH-CHOAc), 5.36 (dq, J = 6.5, 4.9 Hz, 1 H, CHOAc), 4.53 (quin, J = 6.6Hz, 1 H, CHCl), 2.06 (s, 3 H, OAc), 1.60 (d, J = 6.6 Hz, 3 H, CH₃CHCl), 1.32 (d, J = 6.5 Hz, 3 H, CH₃CHOAc); ¹³C NMR (CDCl₃) δ 170.1 (s, OCOCH₃), 133.2 (d, CICH-CH=CH), 130.8 (d, CH= CH-CHOAc), 69.57 (d, CHOAc), 56.69 (d, CHCl), 24.94 (q, CH₃CHCl), 2.1.27 (q, OCOCH₃), 20.02 (q, CH₃CHOAc). Anal. Caled for C₈H₁₃ClO₂: C, 54.40; H, 7.42. Found: C, 54.86; H, 7.42.

 (R^*, S^*) -10 was prepared from (E, Z)-2,4-hexadiene by method A, 24 (0) h (62%). From the ¹H NMR spectrum, the diastereomeric purity was established to be >95% R^*, S^* . On conversion to the diacetate (see under preparation of 3) and by subsequent HPLC analysis, it was found that $E/Z \approx 99/1$.

(R^*, S^*)-10: ¹H NMR (CDCl₃) δ 5.82 (dd, J = 15.4, 6.6 Hz, 1 H, ClCH--CH=CH), 5.71 (dd, J = 15.4, 5.3 Hz, 1 H, CH=CH-CHOAc), 5.37 (dq, J = 6.5, 5.4 Hz, 1 H, CHOAc), 4.53 (quint, J = 6.6Hz, 1 H, CHCl), 2.06 (s, 3 H, OAc), 1.60 (d, J = 6.6 Hz, 3 H, CH₃CHCl), 1.32 (d, J = 6.5 Hz, 3 H, CH₃CHOAc); ¹³C NMR (CDCl₃) δ 170.0 (s, OCOCH₃), 133.4 (d, ClCH--CH=A), 130.8 (d, CH=CH-CHOAc), 69.68 (d, CHOAc), 56.77 (d, CHCl), 25.05 (q, CH₃CHCl), 21.25 (q, CH₃CO), 20.02 (q, CH₃CHOAc); IR (neat) 2980, 1725, 1370, 1240, 1040, 965 cm⁻¹. Anal. Calcd for C₈H₁₃ClO₂: C, 54.40; H, 7.42; Cl, 20.07. Found: C, 54.50; H, 7.35; Cl, 19.91.

cis-1-Acetoxy-4-chloro-2-cyclohexene (cis-11) was prepared from 1,3-cyclohexadiene by adding small portions (method B) of the diene (0.62 g, 7.7 mmol) over 3 h to an acetic acid (28 mL) solution of Pd- $(OAc)_2$ (87 mg 0.39 mmol), LiCl (0.65 g, 15 mmol), LiOAc-2H₂O (1.58 g, 15 mmol), and p-benzoquinone (1.76 g, 16 mmol). After the solution was stirred for an additional 2 h at room temperature, brine (30 mL) was added. The workup procedure was the same as for cis-12. Bulb-to-bulb distillation gave 1.19 g (89%) of cis-11 as a colorless oil (>98% cis according to HPLC): HPLC, $V_r/V_o = 4.5$ (EtOAc/hexane = 2.5/97.5); ¹H NMR (CDCl₃) δ 5.97 (ddd, J = 10.0, 3.7, and 1.5 Hz, 1 H, CH=CH--CHCl), 5.81 (dd, J = 10, 3 Hz, 1 H, CH=CH-OHOAc), 5.28 (m, 1 H, CHOAc), 4.56 (brq, 1 H, CHCl), 2.2-1.9 (m, 4 H, CH₂-CH₂), 2.08 (s, 3 H, OAc); IR (KBr) 1740, 1378, 1245, 1038, 880 cm⁻¹. Anal. Calcd for C₈H₁₁ClO₂: C, 55.02; H, 6.35; Cl, 20.30. Found: C, 55.00; H, 6.30; Cl, 20.16.

trans-11 was obtained in a mixture of *cis*-11 when *p*-benzoquinone was replaced by MnO₂: (isolated by preparative HPLC, EtOAc/hexane = 2.5/97.5); ¹H NMR (CDCl₃) δ 6.02 (dd, J = 10, 4 Hz, 1 H, CH=CH-CHCl), 5.89 (dd, J = 10, 4 Hz, 1 H, CH=CH-CHOAc), 5.28 (br q, J = 4 Hz, 1 H, CHOAc), 4.60 (br q, 1 H, CHCl), 2.35-1.9 (m, 3 H, three of CH₂-CH₂), 2.05 (s, 3 H, OAc), 1.85-1.7 (m, 1 H, one of CH₂-CH₂).

cis-1-Acetoxy-4-chloro-3-methyl-2-cyclohexane (cis-12) was prepared from 2-methyl-1,3-cyclohexadiene by slow addition of the diene (1.38 g, 14.7 mmol) over 3 h to pentane (70 mL) and an acetic acid (45 mL) solution of Pd(OAc)₂ (153 mg, 0.68 mmol), LiCl (1.15 g, 27.2 mmol), LiOAc-2H₂O (2.77 g, 27.2 mmol), and p-benzoquinone (2.94 g, 27.2 mol) (method A). After the solution was stirred at room temperature for 7 h, brine (25 mL) was added and the mixture stirred (15 min) and then filtered. The organic phase was separated and the aqueous phase extracted with ether/pentane (3 × 100 mL, 10/90). The combined organic phases were washed with H₂O (2 × 50 mL), saturated aqueous Na₂CO₃ (50 mL), 2 M NaOH (2 × 40 mL), and brine (50 mL). After drying (MgSO₄), the solvent was removed, yielding crude *cis*-12 which contained 2-methyl-1,3-cyclohexadiene (\approx 10%) and the Diels–Alder adduct (\approx 5%) (δ 3.2–2.9) according to ¹H NMR. Bulb-to-bulb distillation gave 1.51 g (55%) of *cis*-12 (>97% cis according to HPLC): IR (neat) 2950, 1735, 1440, 1360, 1240, 1030, 980 cm⁻¹; ¹H NMR (CDCl₃) δ 5.53 (br s, 1 H, olefin), 5.29 (br t, J = 7 Hz, 1 H, CHOAc), 4.36 (br t, J = 3-4 Hz, CHCl), 2.25–1.9 (m, 4 H, CH₂—CH₂), 2.07 (s, 3 H, OAc), 1.84 (s, 3 H, CH₃—C=C). Irradiation of CH₂CH₂ afforded $J_{(olefin-CHOAc)} = 3.8$ Hz: ¹³C NMR (CDCl₃) δ 170.7 (s, OCOCH₃), 138.0 (s, C=CH), 125.9 (d, C=CH), 69.40 (d, CHOAc), 58.11 (d, CHCl), 30.47 (t, CH₂CHCl), 2.343 (t, CH₂CHOAc), 21.23 (q, CH₃C=C), 21.14 (q, OCOCH₃); HPLC, $V_r/V_0 = 2.3$ (EtOAc/hexane = 5/95). Anal. Calcd for C₉H₁₃ClO₂: C, 57.30; H, 6.95. Found: C, 57.13; H, 6.91.

trans-12 (isolated by prepatative HPLC, $V_r/V_o = 2.0$ (EtOAc/hexane = 5/95)): ¹H NMR (CDCl₃) δ 5.67 (dt, J = 5, 1.5 Hz, 1 H, olefin), 5.23 (m, 1 H, CHOAc), 4.36 (t, J = 3-4 Hz, 1 H, CHCl), 2.30–1.95 (m, 4 H, CH₂CH₂), 2.03 (s, 3 H, OAc), 1.87 (s, 3 H, CH₃C=C). Irradiation of CH₂CH₂ afforded $J_{(olefin-CHOAc)} = 5.0$ Hz.

Mixture of 1 β -acetoxy-4 β -chloro-6 β -methyl-2-cyclohexene (13) and 1 β -acetoxy-4 α -chloro-5 β -methyl-2-cyclohexene (14) was prepared from 5-methyl-1,3-cyclohexadiene by method B, 4.5 (0.5) h, using 7.5 mol % Pd(OAc)₂. The yield was 65% consisting of 13 (46%), α -CH₃-13 (11%), 14 (35%), and β -CH₃-14 (8%). The composition was determined by integration of CHOAc, CHCl, and CH₃ signals in the ¹H NMR spectrum of the mixture. ¹H NMR (CDCl₃) (overlapping signals) 6.1-5.7 (m, 2 H, olefin), 2.2-1.6 (m, 3 H, CH-CH₂). Distinguishable signals are as follows.

13: ¹H NMR (CDCl₃) δ 5.38 (br t, J = 8 Hz, 1 H, CHOAc), 4.4 (br s, 1 H, CHCl), 2.07 (s, 3 H, OAc), 1.12 (d, J = 6.6 Hz, 3 H, CH₃CH). α -CH₃-13: δ 5.23 (br q, J = 4 Hz, 1 H, CHOAc), 4.11 (br d, J = 10Hz, 1 H, CHCl), 1.16 (d, J = 6.6 Hz, 3 H, CH₃CH).

14: ¹H NMR (CDCl₃) δ 5.04 (br d, J = 10 Hz, 1 H, CHOAc), 4.61 (br s, 1 H, CHCl), 2.11 (s, 3 H, OAc), 1.02 (d, J = 6.6 Hz, 3 H, CH₃CH). β -CH₃-14: δ 5.12 (br s, 1 H, CHOAc), 4.55 (br d, J = 10 Hz, 1 H, CHCl), 0.99 (d, concealed, CH₃CH).

cis-1-Acetoxy-4-chloro-2-cycloheptene (15) was prepared from 1,3cycloheptadiene by addition of the diene (683 mg, 7.3 mmol) in one portion to the acetic acid solution (method B), 74%, 12 (0) h. 15 (>98% cis according to ¹H NMR): IR (KBr) 1742, 1374, 1245, 1032 cm⁻¹. HPLC $V_r/V_o = 2.1$ (EtOAc/hexane = 5/95); ¹H NMR (CDCl₃) δ 5.9-5.6 (m, 2 H, olefinic), 5.38 (m, 1 H, CHOAc), 4.63 (m, 1 H, CHCl), 2.06 (s, 3 H, OAc), 2.2-1.7 (m, 6 H, (CH₂)₃); ¹³C NMR (CDCl₃) δ 170.0 (s, OCOCH₃), 133.8 (d), 133.4 (d), 72.64 (d, CHOAc), 58.54 (d, CHCl), 36.17 (t, CH₂CHCl), 32.15 (t), 22.80, (t), 21.15 (q, OCOCH₃). Anal. Calcd for C₉H₁₃ClO₂: C, 57.30; H, 6.95. Found: C, 58.21; H, 6.48.

1β,6α-Diacetoxy-4β-chloro-2-cycloheptene (16) was prepared from 6-acetoxy-1,3-cycloheptadiene by addition of the diene in one portion to the acetic acid solution (method B), 58%, 72 (0) h. **16** (>95% β,β,α according to ¹H NMR): IR (neat) 2940, 1735, 1435, 1370 1240 cm⁻¹; ¹H NMR (CDCl₃) δ 5.91 (br d, 1 H, J = 18 Hz, ClCH—CH—CH), 5.77 (br d, 1 H, J = 12 Hz, CH—CH—CHOAc), 5.62 (br d, 1 H, J = 11 Hz, CH—OAc allylic), 5.36 (dq, 1 H, J = 7, 4.5 Hz, 1 H, CHOAc), 4.83 (m, 1 H, CHCl), 4–2.0 (m, 4 H, CH₂), 2.11 (s, 3 H, OAc), 2.07 (s, 3 H, OAc allylic). Anal. Calcd for C₁₁H₁₅ClO₄: C, 53.55; H, 6.13. Found: C, 53.62; H, 6.02.

Mixture of 17 and 17' was prepared from 1.3-cyclooctadiene by addition of the diene (174 mg, 1.6 mmol) in one portion, 36 (0) h, to an acetic acid (6.5 mL) solution of $Pd(OAc)_2$ (30 mg, 0.16 mmol), LiO-Ac-2H₂O (1.14 g, 11 mmol), LiCl (74 mg, 1.8 mmol), and *p*-benzo-quinone (371 mg, 3.4 mmol) (method B). The mixture was stirred for 36 h at 43 °C. Workup gave 256 mg of a crude product, which on bulb-to-bulb distillation afforded 198 mg (61%) of 17 (75%) and 17' (25%): IR (KBr) 1742, 1372, 1248, 1032 cm⁻¹.

17: ¹H NMR (CDCl₃) δ 5.8–5.5 (m, 2 H, CH=CH--CHOAc), 4.76 (ddd, J = 12.0, 7.5, 4.5 Hz, 1 H, CHCl), 2.05 (s, 3 H, OAc), 2.2–1.5 (m, 8 H, (CH₂)₄).

17': ¹H (CDCl₃) δ 5.8-5.5 (m, 2 H, CH=CH), 5.27 (m, 1 H, CHOAc), 4.93 (ddd, J = 8, 3, 1 Hz, 1 H, CHCl), 2.09 (s, 3 H, OAc), 2.1-1.5 (m, 8 H, (CH₂)₄).

1β,6α-Diacetoxy-4β-(dimethylamino)-2-cycloheptene (18). 16 (50 mg, 0.2 mmol) in THF (1 mL) was added to Pd(acac)₂ (4 mg, 0.013 mmol) and PPh₃ (13.6 mg, 0.052 mmol) in THF (1 mL) under an N₂ atmosphere. Dimethylamine in THF (0.72 mL, 2 M, 1.43 mmol) was added, and the reaction mixture was stirred at 20 °C for 2.5 h. The same workup procedure as described for (R^*,R^*) -21a from (R^*,R^*) -10 afforded 41 mg (81%) of 18: ¹H NMR (CDCl₃) δ 5.94 (br d, J = 11.5 Hz, 1 H, NCH—CH=CH), 5.76 (br d, J = 11.5 Hz, 1 H, AcOCH—CH=CH), 5.65 (br d, J = 10 Hz, 1 H, CHOAc allylic), 5.35 (tt, J =

5.5, 2.5 Hz, 1 H, CHOAc), 3.55 (br d, J = 10 Hz, 1 H, CHN), 2.25 (s, 6 H, $(CH_3)_2N$), 2.10 (s, 3 H, OAc) 2.05 (s, 3 H, OAc allylic), 2.1–1.8 (m, 4 H, CH₂).

1 β ,6 α -Diacetoxy-4 α -(dimethylamino)-2-cycloheptene (19). 16 (50 mg, 0.2 mmol), Na₂CO₃ (76 mg, 0.72 mmol), dimethylamine (180 mg, 4 mmol), and acetonitrile (5 mL) were heated under reflux (2 days). After filtration and concentration, saturated aqueous Na₂CO₃ (3 mL) was added and the resulting mixture extracted with ether (4 × 4 mL). Drying (K₂CO₃) and evaporation in vacuo (aspirator and then oil pump) afforded 35 mg (69%) of 19. ¹H NMR (CDCl₃) δ 5.85 (br d, J = 12 Hz, 1 H, NCH—CH=CH), 5.75 (br d, J = 12 Hz, 1 H, AcOCH—CH=CH), 5.55 (br d, J = 11.2 Hz, 1 H, CHOAc allylic), 5.17 (dq, J = 11, 5.5 Hz, 1 H, CHOAc), 3.29 (br d, 11.6 Hz, 1 H, CHN), 2.25 (s, 6 H, (CH₃)₂N), 2.07 (s, 3 H, OAc), 2.065 (s, 3 H, OAc), 2.2–1.9 (m, 4 H, two-CH₂).

Dimethyl (cis-4-Acetoxycyclohex-2-en-1-yl)malonate. A solution of sodium dimethyl malonate in THF [5.3 mL of a 0.125 M solution (0.66 mmol) prepared from equimolar amounts of dimethyl malonate and sodium hydride (80% in oil)] was added to a mixture of palladium acetate (3.0 mg, 0.013 mmol), triphenyl phosphine (15.8 mg, 0.060 mmol), and cis-1-acetoxy-4-chloro-2-cyclohexene (11) (105 mg, 0.603 mmol) under nitrogen at room temperature. After the solution was stirred for 0.5 h at 25 °C, saturated aqueous sodium hydrogen carbonate (6 mL), water (3 mL), and ether (8 mL) were added. The two layers were separated, and the aqueous layer was extracted with ether $(4 \times 5 \text{ mL})$. The combined organic layers were washed with brine (5 mL), dried (MgSO₄), concentrated to approximately 2 mL, and filtered through a silica gel column. Eluation with ether followed by evaporation of the solvent and removal of the excess dimethyl malonate by bulb-to-bulb distillation furnished 155 mg (95%) of essentially pure dimethyl (cis-1-acetoxy-2cyclohexenyl)malonate (>98% cis): ¹H NMR (CDCl₃) δ. 5.85 (br s, 2 H, CH=CH), 5.19 (m, 1 H, CHO), 3.76 (s, 6 H, MeOOC), 3.35 (d, $J = 9.5 \text{ Hz}, 1 \text{ H}, CH - COOMe), 2.88 (m, 1 \text{ H}, CH - CH(COOMe)_2),$ 2.04 (s, 3 H, OAc), 1.9-1.5 (m, 4 H, CH₂); IR (neat) 2950, 1740, 1435, 1370, 1330, 1230, 1150, 1120 cm^{-1}

Dimethyl (trans-4-Acetoxycyclohex-2-en-1-yl)malonate. cis-1-Acetoxy-4-chloro-2-cyclohexene (11) (166 mg, 0.95 mmol) was added under nitrogen to a solution of sodium dimethyl malonate in dry acetonitrile (10 mL of a 0.113 M solution (1.13 mmol) prepared as above followed by removal of the THF in vacuo and replacing it with CH₃CN). The mixture was heated at reflux for 12 h. Sodium hydrogen carbonate (100 mg, 1.2 mmol) was added and after stirring the mixture at room temperature for 2 h, ether (30 mL) was added followed by filtration. The solvent was removed in vacuo to give a light brown oil (270 mg) containing the product together with dimethyl malonate. The excess dimethyl malonate was removed by bulb-to-bulb distillation to give a residue oil, 242 mg (94%) of essentially pure dimethyl (trans-4-acetoxy-2cyclohexenyl)malonate (>98% trans): ¹H NMR (CDCl₃) δ 5.75 (br s, 2 H, CH=CH), 5.28 (m, 1 H, CHO), 3.75 (s, 6 H, MeOCO), 3.29 (d, $J = 9.0 \text{ Hz}, 1 \text{ H}, CH-COOMe), 2.98 (m, 1 \text{ H}, CH-CH(COOMe)_2),$ 2.06 (s, 3 H, OAc), 2.1-1.5 (m, 4 H, CH₂): IR (neat) 2960, 1740, 1440, 1375, 1250, 1160, 1030 cm⁻¹

(E)-(R*,R*)-2-Acetoxy-5-(diethylamino)-3-hexene ((R*,R*)-21a) from (R^*, R^*) -10. To a stirred solution of Pd(PPh₃)₄ (30 mg, 0.026 mmol) in THF (2 mL) under a N₂ atmosphere was added (R^*, R^*)-10 (70 mg, 0.4 mmol) and diethylamine (200 μ L, 2 mmol). After the solution was stirred for 2 h at 20 °C, ether (5 mL) and 2 M HCl (4 mL) were added. The organic phase was then separated and extracted with aqueous 0.1 M HCl (4×3 mL). After washing with ether (2 mL), the water phase was saturated with solid K₂CO₃ and extracted with ether (4 \times 10 mL). The combined organic phases were dried (K₂CO₃) and the solvent was removed to yield 60 mg (70%) of (R*,R*)-21a (>92% R*,R* according to ¹H NMR): ¹H NMR (CDCl₃) δ 5.69 (dd, J = 15.5, 6.9 Hz, 1 H, NCH-CH=CH), 5.55 (dd, J = 15.6, 5.8 Hz, 1 H, AcOCH-CH=CH), 5.34 (quin, J = 6.3 Hz, 1 H, CHOAc), 3.30 (quin, J = 6.7 Hz, 1 H, CNH), 2.65–2.37 (m, 4 H, CH₂N), 2.04 (s, 3 H, OAc), 1.30 (d, J = 6.4 Hz, 3 H, CH_3 CHOAc), 1.11 (d, J = 6.7 Hz, 3 H, CH_3CHN), 1.01 (t, J = 7.1 Hz, 6 H, CH_3CH_2N); IR (KBr) 2970, 2930, 2800, 1740, 1450, 1370, 1240, 1045 cm⁻¹

 (R^*,R^*) -21a from (R^*,S^*) -10. (R^*,S^*) -10 (70 mg, 0.4 mmol), K₂CO₃ (138 mg, 1 mmol), diethylamine (334 μ L, 3.2 mmol), and CH₃CN (1 mL) were heated under reflux for 25 h. Saturated aqueous Na₂CO₃ (1 mL) was added and the mixture extracted with ether (3 × 5 mL). After drying (K₂CO₃) and rotary evaporation, the crude prouct (75 mg) was bulb-to-bulb distilled, affording 70 mg (80%) of (R^*, R^*) -21a (>90% R^*, R^*) according to ¹H NMR (CDCl₃).

(E)-(R^* , S^*)-2-Acetoxy-5-(diethylamino)-3-hexane ((R^* , S^*)-21a) from (R^* , S^*)-10. The same procedure as for the preparation of (R^* , R^*)-21c from (R^* , R^*)-10 was applied. (R^* , S^*)-10 (70 mg, 0.4 mmol) furnished 60 mg (70%) of (R^* , S^*)-21a (>92% R^* , S^*) according to ¹H NMR (vide infra).

Table III.

AcOCH	$Cr^{1} =$	CR ² C	H ₂ Cl	→
-------	------------	-------------------	-------------------	---

<i>22</i>			50			
chloroacetate (E/Z)	R ¹	R ²	reaction time, h	$\begin{array}{c} \textbf{22, yield} \\ (E/Z) \end{array}$	36 , yield	
$ \begin{array}{r} 1 (9/1) \\ 2 (3.6/1)^a \\ 3 (1/10) \end{array} $	H H CH3	H CH ₃ CH ₃	0.5 3 ^b 8	a , 87% (9/1) b , 92% (3/1) ^c c , 82% (1/10)	a , 9% b , 6% ^d c , 12%	

^aContains 8% of the regioisomer 20. ^b 1.6 mol % Pd(PPh₃)₄ was used. ^cContaminated with 5% of 1-acetoxy-2-methyl-4-(phenyl-sulfonyl)-2-butene. For (E)-22b, see ref 23. ^dContaminated with 4-acetoxy-3-methyl-3-(phenylsulfonyl)-1-butene.

(*R**,*S**)-21a from (*R**,*R**)-10. The same procedure as for the preparation of (*R**,*R**)-21a from (*R**,*S**)-10 was applied. Bulb-to-bulb distillation of the crude product (80 mg) afforded 72 mg (82%) of (*R**,*S**)-21a (>90% *R**,*S** according to ¹H NMR): ¹H NMR (CDCl₃) δ 5.70 (dd, *J* = 15.7, 6.6 Hz, 1 H, NCH—CH=CH), 5.55 (dd, *J* = 15.6, 5.6 Hz, 1 H, ACOCH—CH=CH), 5.35 (quin, *J* = 6.4 Hz, 1 H, CHOAc), 3.29 (quin, *J* = 6.8 Hz, 1 H, CHN), 2.65–2.37 (m, 4 H, CH₂N), 2.04 (s, 3 H, OAc), 1.31 (d, *J* = 6.4 Hz, 3 H, CH₃CHOAc), 1.12 (d *J* = 6.7 Hz, 3 H, CH₃CHN), 1.01 (t, *J* = 7.1 Hz, 6 H, CH₃CH₂N); IR (KBr) 2970, 2930, 2800, 1740, 1450, 1370, 1240, 1040 cm⁻¹.

Dimethyl $((E) - (R^*, S^*)$ -5-Acetoxyhex-3-en-2-yl)malonate $((R^*, S^*)$ -21c) from (R^*, S^*) -10. The same procedure as for the preparation of dimethyl (*cis*-4-acetoxy-2-cyclohexenyl)malonate was applied starting with (R^*, S^*) -10 (88 mg, 0.5 mmol). The excess dimethyl malonate was removed by bulb-to-bulb distillation (90 °C, 1 mmHg), and further distillation (150 °C, 1 mmHg) of the residue afforded 108 mg (79%) of (R^*, S^*) -21c (>95% R^*, S^*) as a colorless oil: ¹H NMR (CDCl₃) δ 5.61 (dd, J = 16, 7 Hz, 1 H, olefin), 5.55 (dd, J = 16, 5 Hz, olefin), 5.28 (quin, J = 4.8 Hz, 1 H, CHOAc) 3.74 (s, 3 H, one of CH_3O), 3.70 (s, 3 H, one of CH_3O), 3.28 (d, J = 9 Hz, 1 H, (CH₃OOC)₂CH), 2.94 (d quin, J = 9, 7 Hz, 1 H, CH_3CHOAc), 1.10 (d, J = 6.6 Hz, 3 H, CH_3CHOAc), 1.10 (d, J = 6.6 Hz, 3 H, CH_3CHOAc).

 (R^*, R^*) -21c from (R^*, R^*) -10. The same procedure as for the preparation of (R^*, S^*) -21c from (R^*, S^*) -10 was applied. (R^*, R^*) -10 (88 mg, 0.5 mmol) furnished 135 mg (99%) of (R^*, R^*) -21c (>95% R^*, R^*): ¹H NMR (CDCl₃) δ 5.62 (dd, J = 16, 7 Hz, 1 H, olefin), 5.55 (dd, J = 16, 5 Hz, olefin), 5.28 (quin, J = 4.8 Hz, 1 H, CHOAc), 3.74 (s, 3 H, one of CH₃O), 3.69 (s, 3 H, one of CH₃O), 3.28 (d, J = 9 Hz, 1 H, (CH₃OOC)₂CH), 2.94 (d quin, J = 9, 7 Hz, 1 H, CHCH-(COOMe)₂), 2.03 (s, 3 H, OAc), 1.27 (d, J = 6.6 Hz, 3 H, CH₃CHOAc), 1.09 (d, J = 7 Hz, 3 H, CH₃CHCH).

(E)-(R*,S*)-2-Acetoxy-5-(phenylsulfonyl)-3-hexene ((R*,S*)-21d) from (R^*, S^*) -10. To a stirred solution of sodium benzene sulfinate (2.62 g, 16 mmol) and Pd(PPh₃)₄ (232 mg, 0.2 mmol) in THF (68 mL) under a N₂ atmosphere at 25 °C was added Me₂SO (12 mL, degassed) and (R^*, S^*) -10 (704 mg, 4 mmol). The mixture was stirred at 50 °C for 2 h and allowed to cool. Water (70 mL) was then added and the mixture extracted with ether (4 \times 60 mL). The combined organic phases were washed with water (30 mL) and brine (30 mL) and finally dried (Mg-SO₄). The solvent was removed and the crude product was purified by flash chromatography, yielding 857 mg (76%) of (R^*, S^*)-21d (>95% R^*, S^*) as a yellow oil: ¹H NMR (CDCl₃) δ 7.80 (d, 2 H, ortho in Ph), 7.65-7.45 (m, 3 H, para and meta in Ph), 5.61 (dd, J = 16.0, 8 Hz, 1 H, SCH--CH=CH), 5.44 (dd, J = 16.0, 6 Hz, 1 H, AcOCH--CH= CH), 5.25 (quin, J = 6 Hz, 1 H, CHOAc), 3.68 (quin, J = 7-8 Hz, 1 H, CHS), 2.02 (s, 3 H, OAc), 1.46 (d, J = 7, 3 Hz, CH₃CHS), 1.17 (d, J = 6.5 Hz, 3 H, CH₃CHOAc).

(E)-(R^* , R^*)-2-Acetoxy-5-(phenylsulfonyl)-3-hexene ((R^* , R^*)-21d) from (R^* , R^*)-10. Applying the same procedure as for the preparation of (R^* , S^*)-21d gave a 71% yield of (R^* , R^*)-21d (>94% R^* , R^*): ¹H NMR (CDCl₃) δ 7.80 (d, 2 H, ortho in Ph), 7.65–7.45 (m, 3 H, para and meta in Ph), 5.64 (dd, J = 16.0, 8 Hz, 1 H, S-CH-CH=CH), 5.46 (dd, J = 16.0, 6 Hz, 1 H, AcOCH-CH=CH), 5.25 (quin, J = 6 Hz, 1 H, CHOAc), 3.69 (quin, J = 7.3 Hz, 1 H, SCH), 2.03 (s, 3 H, OAc), 1.46 (d, J = 6.9 Hz, 3 H, CH₃CHS), 1.19 (d, J = 6.5 Hz, 3 H, CH₃CHOAc); IR (neat) 2970, 1730, 1445, 1370, 1300, 1240, 1140, 1040, 730, 690 cm⁻¹. Anal. Calcd for C₁₄H₁₈SO₄: C, 59.55; H, 6.43. Found: C, 59.64; H, 6.45.

Preparation of 1-Acetoxy-4-(phenylsulfonyl)-2-alkenes (22a-c). Applying the same procedure as for preparation of (R^*,S^*) -21d, but at 20 °C using 5 mmol of chloroacetate in THF (27 mL)-Me₂SO (3 mL), gave 22a-c (Table III).

Acknowledgment. We are grateful to the Swedish Board of Technical Development and the Swedish Natural Science Research Council for financial support and to Thomas Antonsson for performing the syntheses of 18 and 19 during his undergraduate work. We also thank Dr. B. Nilsson (Chemical Center, Lund, Sweden) for a gift of ethyl (E)-4,6-heptadienoate.

Registry No. (E)-1, 34414-28-3; (Z)-1, 55613-61-1; 1', 96039-67-7; (E)-2, 24529-80-4; (Z)-2, 24529-81-5; 2', 38872-51-4; (E)-3, 53060-20-1; (Z)-3, 95177-48-3; 3', 96039-68-8; (E)-4, 76355-81-2; (Z)-4, 96039-69-9; **5a**, 82736-36-5; **5b**, 82736-37-6; **6a**, 96039-70-2; **6b**, 96039-71-3; 6', 96039-72-4; 7**a**, 96039-73-5; 7**b**, 96039-74-6; 7', 96039-75-7; (R*,R*)-10, 95177-49-4; (R*,S*)-10, 95177-50-7; cis-11, 82736-39-8; trans-11, 96039-76-8; cis-12, 96039-77-9; trans-12, 96039-78-0; 13, 96039-82-6; **15**, 82736-40-1; **16**, 96039-83-7; **17**, 96039-84-8; **17**', 96039-85-9; **18**, 96039-86-0; **19**, 96094-31-4; **20**, 58511-44-7; **20**', 96039-87-1; (R*, R*)-**21a**, 96039-88-2; (R*,S*)-**21a**, 96039-89-3; (R*,S*)-**21c**, 96039-91-7; (R*,R*)-**21d**, 95177-61-0; (R*,S*)-**21c**, 96039-91-7; (R*,R*)-**21d**, 95177-61-0; (R*,S*)-**21c**, 96039-91-7; (Z)-**22a**, 95177-57-4; (Z)-**22a**, 95177-58-5; (E)-**22b**, 59830-31-8; (Z)-**22b**, 89345-64-2; (E)-**22c**, 95177-60-9; (E)-**34**, 3780-51-6; (Z)-**34**,

96039-92-8; **35**, 96039-93-9; **36** ($R^1 = CH_3$; $R^2 = H$), 96055-48-0; **36a**, 95199-57-8; 36b, 95199-58-9; 36c, 95199-59-0; Pd(PPh₃)₄, 14221-01-3; Pd(PhCN)₂Cl₂, 14220-64-5; CH₂=CHCH=CH₂, 106-99-0; PPh₃, 603-35-0; Pd(OAc)₂, 3375-31-3; LiCl, 7447-41-8; LiOAc, 546-89-4; $CH_2 = C(CH_3)C(CH_3) = CH_2$, 513-81-5; (E)- $CH_2 = C(CH_3)CH = CHCH_3$, 926-54-5; (E)- $CH_2 = CHCH = CH(CH_2)_2Ph$, 77605-16-4; (E)-CH₂=CHCH=CH(CH₂)₂COOEt, 71779-51-6; (E,Z)-CH₃CH= CHCH=CHCH₃, 5194-50-3; (E,E)-CH₃CH=CHCH=CHCH₃, 5194-51-4; CH₂=C(CH₃), CH=CH₂, 78-79-5; (Z)-CH₂=CHCH= CHCH₃, 1574-41-0; (E)-CH₂=CHCH=CHCH₃, 2004-70-8; Pd(acac)₂, 14024-61-4; Me₂NH, 124-40-3; Et₂NH, 109-89-7; CH₂(COOMe)₂, 108-59-8; PhSO₂Na, 873-55-2; 6-acetoxy-1,3-cycloheptadiene, 29207-42-9; 6-hydroxy-1,3-cycloheptadiene, 1121-63-7; 1,3-cyclohexadiene, 592-57-4; 2-methyl-1,3-cyclohexadiene, 1489-57-2; 6-methyl-4a,5,8,8atetrahydro-5,8-ethano-1,4-naphthoquinone, 96039-94-0; 5-methyl-1,3cyclohexadiene, 19656-98-5; 1,3-cycloheptadiene, 4054-38-0; 1,3-cyclooctadiene, 1700-10-3; dimethyl (cis-4-acetoxy-2-cyclohexenyl)malonate, 82736-52-5; dimethyl (trans-4-acetoxy-2-cyclohexenyl)malonate, 82736-53-6.

Supplementary Material Available: Experimental data on compounds 22 and 36 (2 pages). Ordering information given on any current masthead page.

Synthesis, Thermolysis, and Photolysis of the Azoalkane 4,5-Diazatricyclo[4.3.0.0^{3,7}]nona-4,8-diene: Denitrogenation vs. Azirane Formation

Waldemar Adam* and Karlheinz Hill[†]

Contribution from the Institute of Organic Chemistry, University of Würzburg, D-8700 Würzburg, West Germany. Received August 6, 1984

Abstract: The azoalkane 4,5-diazatricyclo[$4.3.0.0^{3.7}$]nona-4,8-diene (4) was prepared by reaction of the keto urazole 1 with *p*-toluenesulfonohydrazide, treatment of the hydrazone with sodium hydride, which lead to the corresponding urazole 3, and subsequent oxidative hydrolysis. Thermolysis of the azoalkane 4 gave mainly (ca. 95%) the retro-Diels-Alder product (Z)-pyrazole 7, which isomerized to the (E)-pyrazole 7 and the dihydroindazole 8. To a small extent (5%) the 1- and 2-vinylcyclopentadienes were produced, presumably via thermolysis of the tricyclo[$3.2.0.0^{2,7}$]hept-3-ene (5). In the direct photolysis the tricycloalkene 5 and the azirane 6 were formed in the ratio of ca. 1:1, together with traces of quadricyclane, whereas in the benzophenone-sensitized photolysis the azirane 6 was obtained exclusively. In terms of a Salem diagram it is proposed that in the $^{1,3}n,\pi^*$ -excited azoalkane C-C bond cleavage occurs leading to a doubly allyl-stabilized D_{π,π} diradical, which subsequently cyclizes into the azirane 6. In the direct photolysis competitive C-N bond cleavage is observed, leading almost exclusively to the tricycloalkene 5 and a little quadricyclane, but no norbornadiene. The 1,3-diradical 12 serves as immediate precursor to the tricycle 5.

A recent paper¹ on the unusual valence isomerization of perfluoronorbornadiene, involving a thermal di- π -methane rearrangement of quadricyclane into tricyclo[3.2.0.0^{2,7}]hept-3-ene presumably via the 1,3-diradical bicyclo[2.2.1]hept-5-ene-2,7-diyl, prompts us to report our results on the latter parent species, generated in the thermolysis and photolysis of 4,5-diazatricyclo-[4.3.0.0^{3,7}]nona-4,8-diene (4). The synthesis of this new azoalkane commenced with the known keto urazole 1² (eq 1); its details and spectral data are given in the Experimental Section.



The products of the thermal (vacuum flash pyrolysis or VFP at 350 °C and 20 torr) and direct photochemical (irradiation with the 334-, 351-, and 364-nm lines of an argon ion laser in C_6D_6 or CD_3CN) and triplet sensitized reactions (0.45 M benzophenone in C_6D_6 , irradiating only with the 364-nm line of an argon ion laser) are given in Scheme I. The product yields were determined by quantitative ¹H NMR (400 MHz) and/or capillary GC (50-m Carbowax 20 M; injector, column, and detector temperatures of 80, 60 and 120 °C; nitrogen carrier gas pressure 0.25–0.30 kg/cm²). Product balance was better than 75%, remainder being intractable high-molecular-weight material. The retention times and spectral data of the known volatile products tricycloalkene 5,³ norbornadiene, quadricyclane, and mixture of vinylcyclopentadienes⁴ were identical with authentic compounds. The

Dailey, W. P.; Lemal, D. M. J. Am. Chem. Soc. 1984, 106, 1169.
 Adam, W.; De Lucchi, O.; Hill, K. J. Am. Chem. Soc. 1982, 104, 2934.
 (a) Story, P. R. J. Am. Chem. Soc. 1961, 83, 3347.
 (b) Brown, H. C.; Bell, H. M. J. Am. Chem. Soc. 1963, 85, 2324.
 (c) Franzus, B.; Snyder, E. I. J. Am. Chem. Soc. 1963, 85, 3902.

[†]Doctoral Dissertation, University of Würzburg, West Germany, July 1984.