

# Iron-Catalyzed Reduction of CO<sub>2</sub> into Methylene: Formation of C-N, C-O, and C-C Bonds


Guanghua Jin, C. Gunnar Werncke, Yannick Escudié, Sylviane Sabo-Etienne\* and Sébastien Bontemps\*

CNRS, LCC (Laboratoire de Chimie de Coordination), 205 route de Narbonne, BP 44099, F-31077 Toulouse Cedex 4, France and Université de Toulouse, UPS, INPT, F-31077 Toulouse Cedex 4, France

**ABSTRACT:** We report herein the use of the (dihydrido)iron catalyst, Fe(H)<sub>2</sub>(dmpe)<sub>2</sub>, for the selective reduction of CO<sub>2</sub> into either bis(boryl)acetal or methoxyborane depending on the hydroborane used as a reductant. In a one-pot two-steps procedure, the in-situ generated bis(boryl)acetal was shown to be a reactive and versatile source of methylene to create new C-N but also C-O and C-C bonds.

Carbon dioxide is a very abundant and rather non-toxic source of carbon when compared to other C1 sources employed by the chemical industry. Its use has thus attracted interest in various fields of chemistry despite its intrinsic high thermodynamic stability.<sup>1</sup> In the most advanced processes of transformation of CO<sub>2</sub>, the O-C-O unit is maintained, like in carbonate or salicylic acid syntheses and / or the carbon atom remains in the +4 oxidation state like in urea derivatives.<sup>2</sup> However, to fully use CO<sub>2</sub> as a source of carbon, one needs to abstract the oxygen atom(s) and to have access to the full range of carbon oxidation states. As a consequence, homogeneous catalyzed CO<sub>2</sub> reduction with concomitant oxygen abstraction was investigated with dihydrogen,<sup>3</sup> silanes<sup>4</sup> and boranes<sup>5</sup> as reducing agents. Regardless the reductant, the scope of products resulting from the reduction of CO<sub>2</sub> is still rather limited.<sup>6</sup> The use of dihydrogen as a reductant gave rise to methanol,<sup>7</sup> whereas the addition of amines, imines or olefins to the hydrogenation process afforded formamides,<sup>8</sup> methylamines<sup>9</sup> and aliphatic alcohols,<sup>10</sup> thus generating new C-N and C-C bonds. While CO<sub>2</sub> hydrogenation appears as the ideal


reaction in term of atom economy, a sustainable source of “carbon free” dihydrogen and milder reaction conditions remain to be found. The use of silanes and boranes as reductants allowed to significantly milder the operating conditions, and to develop various types of catalysts to afford the complete list of C1 compounds (CO, CH<sub>2</sub>O, CH<sub>3</sub>OH and CH<sub>4</sub>).<sup>4-5</sup> Key mechanistic insights were gained and up to four consecutive hydrofunctionalization reactions have been described (Figure 1) yielding formoxy **A**, acetal **B** and / or formaldehyde, methoxy **C**, and ultimately methane, the latter being only obtained in the case of hydrosilanes. Our studies on ruthenium-catalyzed CO<sub>2</sub> hydroboration shed light on the second stage of CO<sub>2</sub> reduction with the first experimental evidence of the formation of a bis(boryl)acetal compound **B**,<sup>11-12</sup> and of formaldehyde.<sup>13</sup> To increase molecular complexity, Cantat et al. and others described the reduction of CO<sub>2</sub> to the formoxy level **A**, with subsequent functionalization with amine to afford formamides, and further reduction to generate methylamines (Figure 1).<sup>14</sup>



**Figure 1:** General scheme for the reduction and functionalization of CO<sub>2</sub>.

In this context, we sought to further enlarge the scope of carbon-containing compounds accessible from CO<sub>2</sub> transformation. We postulated that the selective synthesis of the bis(boryl)acetal **B** would be possible and that such a compound would be a reactive and versatile source of methylene. In addition, we sought for earth abundant based catalytic systems and looked at defined iron complexes featuring hydride ligands to transfer our expertise from ruthenium polyhydride complexes. The use of iron-based catalysts for homogeneous transformation of CO<sub>2</sub> is scarce and recent studies focused on the formation of formic acid and derivatives.<sup>15</sup> It was also reported that in the presence of multidentate phosphine ligands, the iron precursor Fe(acac)<sub>2</sub> was able to catalyze the reductive functionalization of CO<sub>2</sub> into formamide and methylamine.<sup>16</sup> We choose the dihydride iron complex Fe(H)<sub>2</sub>(dmpe)<sub>2</sub> (**1**) as catalyst precursor, because it has been shown by Field et al. to react with CO<sub>2</sub>,<sup>17</sup> and by us to catalyze the dehydrogenative borylation of arenes.<sup>18</sup> We describe herein the iron-catalyzed selective reduction of CO<sub>2</sub> into bis(boryl)acetal and its subsequent use as a versatile and reactive methylene transfer agent not only to form C-N but also, C-O, C-C, C=N, and C=C bonds in a one-pot two-steps strategy.

Catalytic hydroboration of CO<sub>2</sub> was carried out with 5 mol % of **1** with three different hydroboranes: catecholborane (HBCat), pinacolborane (HBpin) and 9-borabicyclo[3.3.1]nonane (9-BBN) (Scheme 1). The reactions were first conducted in C<sub>6</sub>D<sub>6</sub> at room temperature under 1 atm of CO<sub>2</sub> in a NMR tube and the yields of the two main products observed, bis(boryl)acetal and methoxyborane, were determined by <sup>1</sup>H NMR integrations at full conversion of the corresponding hydroborane.<sup>19</sup> With HBCat, full conversion was observed within 3 h and only methoxyborane was obtained in 59 % yield. In the case of HBpin and 9-BBN, complete conversion occurred within 5h and the corresponding bis(boryl)acetal was favored (30 % and 46 %, respectively) over methoxyborane (18 % and 27 %, respectively).




**Scheme 1:** CO<sub>2</sub> hydroboration with HBCat, HBpin and 9-BBN

We thus choose 9-BBN for a selective synthesis of the bis(boryl)acetal and optimized the experimental conditions. Among the different parameters tested, solvent proved to have the most important impact, since the use of THF favored the formation of the bis(boryl)acetal **2** in terms of both rate and selectivity (Table S3). After 47 min, at full 9-BBN conversion, a yield of 8 % of methoxyborane and 85 % in compound **2** was determined by <sup>1</sup>H NMR integration compare to an internal standard. It should be noted that, before complete reduction to methoxyborane, transient accumulation of bis(boryl)acetal was recently reported in a catalyzed system using strong bases as catalyst and also 9-BBN as reductant.<sup>20</sup> In our system, **2** was not further reduced, since after 15 h, 52 % of acetal and only 11 % of methoxyborane were observed. Unidentified products featuring methylene and methoxy units were observed in the NMR spectra which might account for the decreased amount of **2**.


We then probed the ability of **2** to transfer methylene. As described in Scheme 2, the standard CO<sub>2</sub> reduction was carried

out and followed by the addition of 0.4 equiv. of methanol or diisopropylaniline compare to 9-BBN. This one-pot two-steps procedure led to the corresponding hemiacetal **3** and methyleneaniline **4** in 89 % (72 %) and 83 % (66 %) NMR yields, based on the trapping agent (and 9-BBN), respectively.



**Scheme 2:** Iron-catalyzed selective reduction of CO<sub>2</sub> into bis(boryl)acetal **2** and subsequent functionalization into compounds **3** and **4**.

We then studied the reactivity of compound **2** as a surrogate of formaldehyde in a few representative reactions which were first conducted in NMR tubes (Scheme 3). Unless otherwise stated, yields are based on the trapping agent to account for the efficiency of the trapping reaction. In-situ generated compound **2** was reacted at room temperature with 2-methylaminophenol or N,N-dimethyl-1,2-benzenediamine to afford the cyclic hemiaminal and amination compounds **5** (1 h, 67 % yield) and **6** (3 h, 77 % yield), respectively. Cyclic amination compounds are notably used as organic hydride donors or as N-heterocyclic precursors.<sup>21</sup> The acyclic amination **7** was readily obtained in 92 % yield after 1 h at r.t. by the reaction of 2 equiv. of the corresponding secondary amine. Ammonia is an important building block, but its reactivity is difficult to control. However, the reaction of ammonia with formaldehyde is a known reversible condensation which affords the cage compound hexamethylenetetramine **8**, a versatile organic reagent.<sup>22</sup> In-situ generated compound **2** exposed to an ammonia solution led after 16 h, at r.t. to compound **8** in a 84 % yield based on 9-BBN. This compound features six methylene moieties resulting from the reduction of CO<sub>2</sub>. The characterization of **3-8** completes the list of compounds accessible from the reductive functionalization of CO<sub>2</sub> with amines with the generation of C-N and C-O bonds. With the objective of forming C-C bonds, we turned our attention to phenol reagents. Formaldehyde is indeed used at the industrial scale in phenol resins, via condensation reaction generating C-C bonds. For a proof of concept, we choose to use a t-Bu protected phenol in positions 2 and 4. For comparison and NMR signature in THF, **9** was first generated with an in-situ measured yield of 48 %, from the reaction of para-formaldehyde with the substituted phenol compound in the presence of KOH in 16 h at 120 °C. When in-situ generated compound **2** was used instead of para-formaldehyde, the reaction occurred with the production of **9** in 46 % after 36 h at 120 °C without the need for KOH addition. Encouraged by the generation of such C-C bonds from CO<sub>2</sub>, we then exposed compound **2** to an ylide and observed the expected outcome of a Wittig reaction with the formation of styrene **10** in very good yield (72 %).



**Scheme 3:** Reactions of in-situ generated **2** affording compounds **5-10**.

We used both  $^{13}\text{CO}_2$  and  $^{12}\text{CO}_2$  to confirm by NMR and HRMS that the methylene carbon atom within **3-10** resulted from the reduction of  $\text{CO}_2$ . NMR data for the bis(boryl)acetal **2** and for the resulting compounds **3-10** are provided in Table S4. The important variations of  $^{13}\text{C}$  and  $^1\text{H}$  NMR chemical shifts (from  $\delta(^{13}\text{C}) = 32.6$  and  $\delta(^1\text{H}) = 3.88$  for compound **9** to  $\delta(^{13}\text{C}) = 156.5$  and  $\delta(^1\text{H}) = 7.73$  for compound **4**), indicating different types of methylene, highlight the versatile reactivity of compound **2**. The experiments performed in NMR tubes allowed detailed and convenient monitoring. In a next stage, we sought for isolation of the functionalized products and compounds **8** and **9** were selected. On a 10-fold scale up synthesis, compound **2** was selectively generated at  $60^\circ\text{C}$  in 10 min, and subsequently trapped to afford compounds **8** and **9** in 98 and 37 % isolated yields based on the trapping agent, respectively. A 100-fold scale up synthesis led to compound **8** in 2 days at room temperature with a 70 % (106.2 mg) isolated yield based on 9-BBN.

To conclude, we report herein the use of a (dihydrido)iron complex in the reduction of  $\text{CO}_2$  into either bis(boryl)acetal or methoxyborane depending on the hydroborane used as reductant. Selective reduction of  $\text{CO}_2$  to the acetal level and subsequent functionalization in a one-pot two-steps procedure, allowed to transform  $\text{CO}_2$  into methylene and to considerably enlarge the scope of accessible functions by generating new C-N but also C-O and C-C bonds. Our results further highlight the importance of metal hydride complexes in the controlled transformation of  $\text{CO}_2$ . We are currently conducting mechanistic investigation to understand and improve the catalytic system and pursuing research on the specific reactivity of bis(boryl) and / or bis(silyl)acetal compounds.<sup>23</sup>

## ASSOCIATED CONTENT

Experimental details and NMR data. "This material is available free of charge via the Internet at <http://pubs.acs.org>."

## AUTHOR INFORMATION

### Corresponding Author

[sylviane.sabo@lcc-toulouse.fr](mailto:sylviane.sabo@lcc-toulouse.fr), [sebastien.bontemps@lcc-toulouse.fr](mailto:sebastien.bontemps@lcc-toulouse.fr)

### Author Contributions

All authors have given approval to the final version of the manuscript.

## ACKNOWLEDGMENT

CSC is gratefully acknowledged for the PhD fellowship of G.J. Support from CNRS and the ANR (Programme blanc "IRONHYC" ANR-12) is also acknowledged. We thank Dr. Christian Bijani for NMR analyses.

## REFERENCES

- Some selected reviews: (a) Goepfert, A.; Czaun, M.; Jones, J.-P.; Surya Prakash, G. K.; Olah, G. A., *Chem. Soc. Rev.* **2014**, *43*, 7995-8048; (b) Leitner, W., *Angew. Chem. Int. Ed.* **1995**, *34*, 2207-2221; (c) Aresta, M., Carbon Dioxide Reduction and Uses as a Chemical Feedstock. *Activation of small molecules*, Tolman, W. B., Ed. Wiley: Weinheim, Germany, 2006; pp 1-42; (d) Sakakura, T.; Choi, J.-C.; Yasuda, H., *Chem. Rev.* **2007**, *107*, 2365-2387; (e) Benson, E. E.; Kubiak, C. P.; Sathrum, A. J.; Smieja, J. M., *Chem. Soc. Rev.* **2009**, *38*, 89-99; (f) Fan, T.; Chen, X.; Lin, Z., *Chem. Comm.* **2012**, *48*, 10808-10828; (g) Appel, A. M.; Bercaw, J. E.; Bocarsly, A. B.; Dobbek, H.; DuBois, D. L.; Dupuis, M.; Ferry, J. G.; Fujita, E.; Hille, R.; Kenis, P. J. A.; Kerfeld, C. A.; Morris, R. H.; Peden, C. H. F.; Portis, A. R.; Ragsdale, S. W.; Rauchfuss, T. B.; Reek, J. N. H.; Seefeldt, L. C.; Thauer, R. K.; Waldrop, G. L., *Chem. Rev.* **2013**, *113*, 6621-6658; (h) Costentin, C.; Robert, M.; Saveant, J.-M., *Chem. Soc. Rev.* **2013**, *42*, 2423-2436; (i) Fontaine, F.-G.; Courtemanche, M.-A.; L egar e, M.-A., *Chem. Eur. J.* **2014**, *20*, 2990-2996.
- (a) Centi, G.; Perathoner, S., *Catal. Today* **2009**, *148*, 191-205; (b) Peters, M.; K ohler, B.; Kuckshinrichs, W.; Leitner, W.; Markewitz, P.; M uller, T. E., *ChemSusChem* **2011**, *4*, 1216-1240; (c) Quadrelli, E. A.; Centi, G.; Duplan, J.-L.; Perathoner, S., *ChemSusChem* **2011**, *4*, 1194-1215; (d) Aresta, M.; Dibenedetto, A.; Angelini, A., *Chem. Rev.* **2014**, *114*, 1709-1742.
- (a) Jessop, P. G.; Joo, F.; Tai, C.-C., *Coord. Chem. Rev.* **2004**, *248*, 2425-2442; (b) Wang, W.; Wang, S.; Ma, X.; Gong, J., *Chem. Soc. Rev.* **2011**, *40*, 3703-3727.
- Fernandez-Alvarez, F. J.; Aitani, A. M.; Oro, L. A., *Catal. Sci. Technol.* **2014**, *4*, 611-624.
- (a) Bontemps, S., *Coord. Chem. Rev.* **2015**, DOI: 10.1016/j.ccr.2015.06.003; (b) Chong, C. C.; Kinjo, R., *ACS Catalysis* **2015**, 3238-3259.
- Liu, Q.; Wu, L.; Jackstell, R.; Beller, M., *Nat. Commun.* **2015**, *6*, DOI: 10.1038/ncomms6933.
- (a) Huff, C. A.; Sanford, M. S., *J. Am. Chem. Soc.* **2011**, *133*, 18122-18125; (b) Wesselbaum, S.; vom Stein, T.; Klankermayer, J.; Leitner, W., *Angew. Chem. Int. Ed.* **2012**, *51*, 7499-7502; (c) Wesselbaum, S.; Moha, V.; Meuresch, M.; Brosinski, S.; Thenert, K. M.; Kothe, J.; vom Stein, T.; Englert, U.; Holscher, M.; Klankermayer, J.; Leitner, W., *Chem. Sci.* **2015**, *6*, 693-704; (d) Rezayee, N. M.; Huff, C. A.; Sanford, M. S., *J. Am. Chem. Soc.* **2015**, *137*, 1028-1031.
- Zhang, L.; Han, Z.; Zhao, X.; Wang, Z.; Ding, K., *Angew. Chem. Int. Ed.* **2015**, *54*, 6186-6189.
- (a) Beydoun, K.; vom Stein, T.; Klankermayer, J.; Leitner, W., *Angew. Chem. Int. Ed.* **2013**, *52*, 9554-9557; (b) Li, Y.; Sorribes, I.; Yan, T.; Junge, K.; Beller, M., *Angew. Chem. Int. Ed.* **2013**, *52*, 12156-12160; (c) Beydoun, K.; Ghattas, G.; Thenert, K.; Klankermayer, J.; Leitner, W., *Angew. Chem. Int. Ed.* **2014**, *53*, 11010-11014.
- (a) Tominaga, K.-i.; Sasaki, Y., *Catal. Commun.* **2000**, *1*, 1-3; (b) Liu, Q.; Wu, L.; Fleischer, I.; Selent, D.; Franke, R.; Jackstell, R.; Beller, M., *Chem. Eur. J.* **2014**, *20*, 6888-6894.
- (a) Bontemps, S.; Vendier, L.; Sabo-Etienne, S., *Angew. Chem. Int. Ed.* **2012**, *51*, 1671-1674; (b) Bontemps, S.; Sabo-Etienne, S., *Angew. Chem. Int. Ed.* **2013**, *52*, 10253-10255.
- Very recently, Fontaine, Stephan et al. reported the first X-Ray characterization of a bis(boryl)acetal compound resulting from the reduction of  $\text{CO}_2$ : see Courtemanche, M.-A.; Pulis, A. P.; Rochette, E.;

Legare, M.-A.; Stephan, D. W.; Fontaine, F.-G., *Chem. Comm.* **2015**, 51, 9797.

13. Bontemps, S.; Vendier, L.; Sabo-Etienne, S., *J. Am. Chem. Soc.* **2014**, *136*, 4419-4425.

14. (a) Jacquet, O.; Das Neves Gomes, C.; Ephritikhine, M.; Cantat, T., *J. Am. Chem. Soc.* **2012**, *134*, 2934-2937; (b) Das Neves Gomes, C.; Jacquet, O.; Villiers, C.; Thuéry, P.; Ephritikhine, M.; Cantat, T., *Angew. Chem. Int. Ed.* **2012**, *51*, 187-190; (c) Li, Y.; Fang, X.; Junge, K.; Beller, M., *Angew. Chem. Int. Ed.* **2013**, *52*, 9568-9571; (d) Jacquet, O.; Frogneux, X.; Das Neves Gomes, C.; Cantat, T., *Chem. Sci.* **2013**, *4*, 2127-2131; (e) Das, S.; Bobbink, F. D.; Laurency, G.; Dyson, P. J., *Angew. Chem. Int. Ed.* **2014**, *53*, 12876-12879; (f) Tlili, A.; Blondiaux, E.; Frogneux, X.; Cantat, T., *Green Chem.* **2015**, *17*, 157-168.

15. (a) Tai, C.-C.; Chang, T.; Roller, B.; Jessop, P. G., *Inorg. Chem.* **2003**, *42*, 7340-7341; (b) Federsel, C.; Boddien, A.; Jackstell, R.; Jennerjahn, R.; Dyson, P. J.; Scopelliti, R.; Laurency, G.; Beller, M., *Angew. Chem. Int. Ed.* **2010**, *49*, 9777-9780; (c) Langer, R.; Diskin-Posner, Y.; Leitius, G.; Shimon, L. J. W.; Ben-David, Y.; Milstein, D., *Angew. Chem. Int. Ed.* **2011**, *50*, 9948-9952; (d) Ziebart, C.; Federsel, C.; Anbarasan, P.; Jackstell, R.; Baumann, W.; Spannenberg, A.; Beller, M., *J. Am. Chem. Soc.* **2012**, *134*, 20701-20704; (e) Inoue, Y.; Sasaki, Y.; Hashimoto, H., *J. Chem. Soc., Chem. Commun.* **1975**, 718-719; (f) Evans, G. O.; Newell, C. J., *Inorg. Chim. Acta* **1978**, *31*, L387-L389; (g) Fong, H.; Peters, J. C., *Inorg. Chem.* **2015**, *54*, 5124-5135; (h) Zhang, Y.; MacIntosh, A. D.; Wong, J. L.; Bielinski, E. A.; Williard, P. G.; Mercado, B. Q.; Hazari, N.; Bernskoetter, W. H., *Chem. Sci.* **2015**, *6*, 4291-4299; (i) Pinaka, A.; Vougioukalakis, G. C., *Coord. Chem. Rev.* **2015**, *288*, 69-97.

16. Frogneux, X.; Jacquet, O.; Cantat, T., *Catal. Sci. Technol.* **2014**, *4*, 1529-1533.

17. (a) Whittlesey, M. K.; Mawby, R. J.; Osman, R.; Perutz, R. N.; Field, L. D.; Wilkinson, M. P.; George, M. W., *J. Am. Chem. Soc.* **1993**, *115*, 8627-8637; (b) Field, L. D.; Lawrenz, E. T.; Shaw, W. J.; Turner, P., *Inorg. Chem.* **2000**, *39*, 5632-5638; (c) Allen, O. R.; Dalgarno, S. J.; Field, L. D., *Organometallics* **2008**, *27*, 3328-3330.

18. Dombrey, T.; Werncke, C. G.; Jiang, S.; Grellier, M.; Vendier, L.; Bontemps, S.; Sortais, J.-B.; Sabo-Etienne, S.; Darcel, C., *J. Am. Chem. Soc.* **2015**, *137*, 4062-4065.

19. Except for R<sub>2</sub>BOBR<sub>2</sub>, no other compound in significant amount was observed by NMR during the hydroboration processes.

20. Das Neves Gomes, C.; Blondiaux, E.; Thuéry, P.; Cantat, T., *Chem. Eur. J.* **2014**, *20*, 7098-7106.

21. (a) Zhu, X.-Q.; Zhang, M.-T.; Yu, A.; Wang, C.-H.; Cheng, J.-P., *J. Am. Chem. Soc.* **2008**, *130*, 2501-2516; (b) Benhamou, L.; Chardon, E.; Lavigne, G.; Bellemin-Lapponnaz, S.; César, V., *Chem. Rev.* **2011**, *111*, 2705-2733; (c) Poyatos, M.; Prades, A.; Gonell, S.; Gusev, D. G.; Peris, E., *Chem. Sci.* **2012**, *3*, 1300-1303.

22. Blažević, N.; Kolbah, D.; Belin, B.; Šunjić, V.; Kajfež, F., *Synthesis* **1979**, *1979*, 161-176.

23. During the course of the reviewing process of the present paper, Cantat et al. reported the synthesis of amination from the silane-reduction of CO<sub>2</sub> in the presence of amine. Using a malonate compound instead of an amine, C-C bonds were also generated, see: Frogneux, X.; Blondiaux, E.; Thuéry, P.; Cantat, T., *ACS Catalysis* **2015**, *5*, 3983-3987.

Insert Table of Contents artwork here

