

A New Method for Conversion of Aldoximes into Nitriles: Use of Chlorothionoformates

By D. L. J. CLIVE

(Chemistry Department, Imperial College, London, S.W.7)

Summary *p*-Chlorophenyl chlorothionoformate in the presence of pyridine dehydrates aldoximes under extremely mild conditions.

VERY few mild procedures have been established for dehydration of aldoximes,¹ but in those cases where the vigorous conditions normally prescribed² are acceptable, the sequence constitutes an important preparative route to nitriles. We have examined the response of several aldoximes towards *p*-chlorophenyl chlorothionoformate³ and find that nitriles are produced at room temperature in a process that is both rapid and exceptionally mild.

The method involves addition of a solution of the oxime and pyridine (1 and 2 equiv., respectively) to a solution of the reagent (2 equiv.) followed by isolation of the nitrile, usually by chromatography.[†]

Results with the new process are shown in the Table, in which the yields refer to analytically pure[‡] product obtained, with the exception of heptanonitrile which was prepared on a larger scale, from 1 mmole of aldoxime. In these experiments the reaction time was 1.5 h, but production of the nitrile is probably complete much sooner.

For comparative purposes (*E*)-*p*-chlorobenzaldoxime was treated with *p*-chlorophenyl chloroformate under the above conditions and, though the oxime was rapidly acylated, no

nitrile[§] was detectable (i.r. control) even after a 16 h reaction period. Mechanistically, the new process probably involves formation of an *O*-acyl derivative (1) and the facility with which it decomposes⁴ may reflect a rapid (*E*) → (*Z*) isomerization and subsequent collapse, and/or

TABLE

Oxime	Yield of nitrile
(<i>E</i>)-Veratraldoxime ^{a,b}	61.0%
(<i>E</i>)- <i>p</i> -Nitrobenzaldoxime ^c	70.3
(<i>E</i>)- <i>p</i> -Chlorobenzaldoxime ^b	60.8
(<i>Z</i>)- <i>p</i> -Chlorobenzaldoxime ^b	42.4
Heptanaldoxime ^{b,d}	42.7

^a Nomenclature of J. E. Blackwood, C. L. Gladys, K. L. Loening, A. E. Petrarca, and J. E. Rush, *J. Amer. Chem. Soc.*, 1968, **90**, 509. ^b Diethyl ether used as reaction solvent. ^c Dichloromethane used as reaction solvent. ^d Prepared according to the procedure of E. W. Bousquet, *Org. Synth.*, Coll. Vol. II, p. 313, and crystallised from petroleum, the material was a mixture of the (*Z*)- and (*E*)- isomers in the molar ratio of ca. 6:1. Cf. G. G. Kleinspehn, J. A. Jung, and S. A. Studniarz, *J. Org. Chem.*, 1967, **32**, 460.

the thermodynamic instability of the system (2) with respect to its isomer (3).§

In principle, the substantial driving force available⁵ from



[†] Reaction mixtures were protected from moisture and oxygen as a matter of routine. It was convenient, though not essential, to remove residual *p*-chlorophenol by extraction into dilute aqueous potassium hydroxide.

[‡] Analytical values accurate to ± 0.3%. The compounds were identical with authentic specimens.

[§] There is ample precedent for (*E*)-(Z) interconversion and derivatives of (*Z*)-aldoximes are well known to be much less stable with respect to nitrile production than their (*E*)-isomers.^{2b}

the conversion of (2) into (3) could involve [for an (*E*)-derivative] a cyclic process (4). In so far as the essential features of such an array are comparable with (5) (a retro-

ene reaction⁶), the thermal concerted process represented by (4) is one in which orbital symmetry is conserved. Similar possibilities, generalized in (6), may also apply to the pyrolysis of various imino-ethers⁷ of structure (7) and to the observation, made many years ago but undeveloped in terms of its possibilities as a *mild* synthetic method, that certain combinations of aldoxime and aryl isothiocyanate afford nitriles.⁸



This work was carried out during tenure of an I.C.I. Research Fellowship.

(Received, June 17th, 1970; Com. 943.)

¹ P. J. Foley, jun., *J. Org. Chem.*, 1969, **34**, 2805; T. J. Bentley, J. F. McGhie, and D. H. R. Barton, *Tetrahedron Letters*, 1965, 2497; cf. J. H. Pomeroy and C. A. Craig, *J. Amer. Chem. Soc.*, 1959, **81**, 6340.

² (a) D. T. Mowry, *Chem. Rev.*, 1948, **42**, 189; (b) H. Metzger in "Methoden der Organischen Chemie," Houben-Weyl, Stuttgart, 1968, 4th edn., vol. X, part 4, p. 1.

³ Readily available by the method of D. L. Garmaise, A. Uchiyama, and A. F. McKay, *J. Org. Chem.*, 1962, **27**, 4509.

⁴ Cf. J. H. M. Hill and L. D. Schmookler, *J. Org. Chem.*, 1967, **32**, 4025.

⁵ C. H. DePuy and R. W. King, *Chem. Rev.*, 1960, **60**, 431.

⁶ For a review of the ene reaction see H. M. R. Hoffmann, *Angew. Chem. Internat. Edn.*, 1969, **8**, 556.

⁷ H. Hettler and H. Neygenfind, *Tetrahedron Letters*, 1966, 6031; E. Grigat and R. Pütter, *Chem. Ber.*, 1966, **99**, 2361.

⁸ An. Obregia and C. V. Gheorghiu, *J. prakt. Chem.*, 1930, **128**, 239; C. V. Gheorghiu, *ibid.*, 1931, **130**, 49.

⁹ D. Martin and W. Mucke, *Chem. Ber.*, 1965, **98**, 2059.