

Rhodium(I) complexes with *N*-heterocyclic carbenes bearing a 2,3,4,5-tetraphenylphenyl and its higher dendritic frameworks†

Hiromichi Sato, Tetsuaki Fujihara, Yasushi Obora, Makoto Tokunaga, Junya Kiyosu and Yasushi Tsuji*

Received (in Cambridge, UK) 30th August 2006, Accepted 29th September 2006

First published as an Advance Article on the web 25th October 2006


DOI: 10.1039/b612385f

Novel rhodium(I) complexes with *N*-heterocyclic carbenes bearing a 2,3,4,5-tetraphenylphenyl and its higher dendritic frameworks were synthesized and the complexes were used as catalysts in rhodium-catalyzed hydrosilylation of α,β -unsaturated ketones to show high 1,4-adduct selectivity.

Dendritic frameworks have received considerable attention to develop new materials¹ including catalysts.² Among them, a 2,3,4,5-tetraphenylphenyl (TPPh) moiety developed by Müllen and co-workers have been utilized as building blocks for polyphenylene nanomaterials.³ We are interested in their spatially spread and rigid structures. Palladium acetates with pyridine ligands having a TPPh moiety (**1a**) and its higher dendritic framework (**1b**) successfully suppressed Pd black formation in the air-oxidation of alcohols (Scheme 1).⁴ The diamine (**2**) with TPPh groups was effectively employed as a ligand in a palladium-catalyzed kinetic resolution of various biaryl compounds.⁵ Recently, we found that the phosphines (**3**) having a TPPh moiety are effective ligands to utilize unactivated aryl chlorides in palladium-catalyzed Suzuki–Miyaura and Mizoroki–Heck reactions.⁶

In this communication, we have selected an *N*-heterocyclic carbene (NHC) as a ligand. A use of NHC would be beneficial because stronger bonds to metals as compared with conventional ligands such as phosphines diminish their dissociation from metal centers.⁷ Here, we designed and synthesized novel rhodium(I) complexes with NHC ligands bearing a TPPh and its higher dendritic moieties. These complexes are active catalysts and provide 1,4-adducts predominantly in the hydrosilylation of α,β -unsaturated ketones with Ph_2SiH_2 , which has opposite regioselectivity to the one obtained with conventional Rh-phosphine and Rh-NHC catalyst systems.

Imidazolium salts (**4a** and **4b**) having the TPPh and its higher dendritic frameworks were synthesized in three steps *via* the amination with benzophenone imine^{5,8,9} followed by the diimine formation¹⁰ (Scheme 2). The dendritic moieties of **4b** and **5b** have




Scheme 1 Ligands bearing a 2,3,4,5-tetraphenylphenyl (TPPh) moiety.

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, and CREST, Japan Science and Technology Agency (JST), Kyoto 615-8510, Japan and Catalysis Research Center and Division of Chemistry, Graduate School of Science, Hokkaido University, Sapporo, 001-0021, Japan.

E-mail: ytsuji@scl.kyoto-u.ac.jp; Fax: +81-75-383-2514

† Electronic supplementary information (ESI) available: Experimental procedures and data for characterization of **4** and **5**. See DOI: 10.1039/b612385f









Scheme 2 Synthesis of NHC rhodium(I) complexes.

the methyl substituents to ensure good solubility of these materials.⁴ The MALDI-TOF-MS spectra of **4a** and **4b** showed peaks at $m/z = 981$ ($[M-Cl]^+$) and 2782 ($[M-Cl]^+$), respectively. Rhodium(I) complexes with the corresponding NHC ligands (**5a** and **5b**) were synthesized in good yields (69 and 70%, respectively) by the carbene-transfer method¹¹ using NHC–Ag species derived from **4a** and **4b** (Scheme 2). The new complexes (**5a** and **5b**) were fully characterized, and their mass spectra showed peaks at $m/z = 1227$ (FD-MS: $[M]^+$) and 2992 (MALDI-TOF-MS: $[M-Cl]^+$), respectively. In $^{13}C\{^1H\}$ NMR spectra of **5a** and **5b**, the coordinated carbene carbons appeared with distinct ^{13}C – ^{103}Rh couplings at 184.4 ppm ($^1J_{Rh-C} = 48.7$ Hz) and 184.3 ppm ($^1J_{Rh-C} = 47.7$ Hz), respectively, indicating their tight coordination to the rhodium atoms in solution.

The molecular structure of **5a** has been successfully determined by X-ray crystal structural analysis (Fig. 1a).[‡] It is evident that the TPPh moiety of the NHC ligand spatially spreads out. The structure of **5a** shows that the longest distance between the edge of the two TPPh groups is 2.9 nm. The Rh–C(carbene) bond length of **5a** (2.010(5) Å) is similar to those of other Rh(I)–NHC complexes.^{12,13b} As for **5b**, all the trials to obtain single crystals suitable for crystallographic analysis were unsuccessful. Therefore, the optimized structure of **5b** (Fig. 1b) were obtained by ONIOM calculation¹⁴ (B3LYP/LANL2DZ:UFF) using an initial structure based on the X-ray structure of **5a**. As a result, **5b** is a bigger nano-sized complex with the longest distance between the edge of the two dendritic frameworks being 4.9 nm.

As shown in Fig. 1, the complexes **5a** and **5b** have the very unique rigid and spatially spread structures. Thus, catalytic behavior of **5a** and **5b** was intriguing and compared with the representative Rh(I) NHC complex, $RhCl(COD)[(Mes)_2(C_3H_2N_2)]$ (**5c**) (Scheme 2).¹³ The hydrosilylation¹⁵ of α,β -unsaturated ketones with Ph_2SiH_2 was carried out in the presence of a catalytic amount (1 mol%) of **5** in CH_2Cl_2 at room temperature (Table 1).[§] It is well-established that regioselectivity in the Rh-catalyzed hydrosilylation of α,β -unsaturated ketones was preferentially determined by the nature of silanes employed:¹⁶ use of di- or tri-hydrosilanes such as Ph_2SiH_2 or $PhSiH_3$ lead to 1,2-adducts giving allylic alcohols after the desilylation, whereas mono-hydrosilanes such as Et_3SiH or Me_2PhSiH afford 1,4-adducts giving ketones. The hydrosilylation of 2-cyclohexen-1-one with Ph_2SiH_2 in the presence of **5a** as a

Table 1 Hydrosilylation of α,β -unsaturated ketones catalyzed by **5**^a

		$R^1-CH=CH-C(=O)-R^3 \xrightarrow[CH_2Cl_2, rt, 24 h]{Ph_2SiH_2 (1 \text{ mol\%})} \xrightarrow[Et_2O]{K_2CO_3, MeOH} R^1-CH_2-CH_2-C(=O)-R^3 + R^1-CH_2-CH(OH)-R^3$			
		1,4-Adduct		1,2-Adduct	
Entry	Substrate	Catalyst	Yield ^b /%	Selectivity ^b (1,4-/1,2-)	
1		5a	77	91	/ 9
2		5b	91	88	/ 12
3		5c	92	8	/ 92
4		5a	67	82	/ 18
5		5b	62	81	/ 19
6		5c	88	9	/ 91
7		5a	69	93	/ 7
8		5b	76	100	/ 0
9		5c	53	60	/ 40
10		5a	96	94	/ 6
11		5b	46	50	/ 50
12		5c	43	12	/ 88
13		5a	38	16	/ 84
14		5b	51	4	/ 96
15		5c	72	0	/ 100

^a α,β -Unsaturated ketone (1 mmol), Ph_2SiH_2 (1.2 mmol), **5** (0.01 mmol), CH_2Cl_2 (1 cm³), rt, 24 h. ^b By GC analysis after the desilylation with K_2CO_3 –MeOH.

catalyst afforded the 1,4-adduct (cyclohexanone) in 91% selectivity with a smaller amount of the 1,2-adduct (2-cyclohexen-1-ol) in 77% total yield after the desilylation (entry 1). The complex **5b** as a catalyst also afforded the 1,4-adduct in 88% selectivity (entry 2). Noteworthy is that these 1,4-selectivities obtained with Ph_2SiH_2 in entries 1 and 2 were totally opposite to one obtained with $RhH(PPh_3)_4$ as a catalyst,^{16b} in which only the 1,2-adduct was afforded exclusively. Furthermore, in contrast to **5a** and **5b**, **5c** as a catalyst afforded the 1,2-adduct as a major product in 92% selectivity (entry 3). Thus, the TPPh moieties of **5a** and **5b** play a critical role to reverse the typical regioselectivity associated with the dihydrosilane. Similar effect of the TPPh to reverse the regioselectivity was confirmed in the hydrosilylation of 3-octen-2-one: **5a** and **5b** afforded the 1,4-adduct (2-octanone) in 82 and 81% selectivities, respectively (entries 4 and 5), while **5c** provided the 1,2-adduct (3-octen-2-ol) in 91% selectivity (entry 6). In the hydrosilylation of benzylideneacetone, **5a** and **5b** provided the 1,4-adduct (4-phenyl-2-butanone) in 93 and 100% selectivities, respectively (entries 7 and 8), whereas selectivity of the 1,2-adduct (4-phenyl-3-buten-2-ol) increased considerably by using **5c** as a catalyst (entry 9). This selectivity with **5c** was comparable to that obtained with $RhCl(PPh_3)_3$ as the catalyst.^{16a} With 4-methyl-3-penten-2-one having the two substituents at the β -position, **5a** realized the good 1,4-selectivity (entry 10), while **5b** lowered both the regioselectivity and yield of the products (entry 11). The steric bulk associated with the higher dendritic framework of **5b** might affect the selectivity and catalytic activity. In the hydrosilylation of 3-methyl-2-cyclohexen-1-one, even **5a** and **5b** as well as **5c** all provided the high 1,2-selectivities (84–100%), possibly due to significant steric congestion at the β -position (entries 13–15).

Recently, we have synthesized analogous Rh(I) NHC complexes (**6a–d** in Scheme 3)¹⁷ with very flexible dendritic frameworks consisting of the Fréchet-type benzyl ether moieties.¹⁸ We are interested how rigid^{4–6} and flexible^{17–19} dendritic frameworks





Fig. 1 Molecular structures of **5**. (a) Crystal structure of **5a**. (b) The optimized structure of **5b** calculated by ONIOM method.



Scheme 3 Hydrosilylation of 2-cyclohexen-1-one catalyzed by flexible dendrimer complexes **6**.

operate in the same hydrosilylation of α,β -unsaturated ketones. In the reaction of 2-cyclohexen-1-one, **6a–d** with a series of dendritic moieties of different generations all afforded the 1,2-adduct mainly in similar selectivities (61–69%, Scheme 3). In contrast to **5a** and **5b** having the rigid moieties, **6a–d** could not change the regioselectivity inherent to Ph_2SiH_2 . These flexible dendrimer moieties tend to fold back around an active site.¹⁷ The rigid and spatially spread structures of **5a** and **5b** might be crucial for causing the 1,4-regioselectivity.

In conclusion, novel rhodium(i) NHC complexes having a TPPH and its higher dendritic frameworks were synthesized and fully characterized. These complexes are efficient catalysts in the hydrosilylation of α,β -unsaturated ketones with Ph_2SiH_2 which has the opposite regioselectivity to the one obtained with conventional Rh-phosphine or Rh–NHC catalysts. Further studies on transition-metal catalysts bearing rigid and flexible dendritic frameworks are under investigation.

Notes and references

† Single crystals of **5a**· C_5H_{12} suitable for X-ray diffraction study were obtained by diffusion of *n*-pentane into **5a** in CH_2Cl_2 . Crystal data for **5a**· C_5H_{12} : $\text{C}_{88}\text{H}_{76}\text{ClN}_2\text{Rh}$, $M = 1299.9$, $T = 113$ K, triclinic, space group $P\bar{1}$ (No. 2), $a = 14.85(2)$, $b = 20.51(2)$, $c = 25.04(2)$ Å, $\alpha = 79.38(7)$, $\beta = 71.05(9)$, $\gamma = 81.10(10)^\circ$, $U = 7054(1)$ Å³, $Z = 4$, $\mu(\text{Mo K}\alpha) = 3.30$ cm^{−1}, Unique reflections 31888, Observed reflections 16076 ($I > 3\sigma(I)$), $R_1, wR_2 = 0.056, 0.164$, GOF = 1.01, CCDC 617650. For crystallographic data in CIF or other electronic format see DOI: 10.1039/b612385f

§ General procedure for the hydrosilylation (entry 1 in Table 1): A mixture of **5a** (0.01 mmol, 12 mg), CH_2Cl_2 (1 cm³), 2-cyclohexen-1-one (1 mmol,

9.7×10^{-2} dm³), and bibenzyl (0.25 mmol, 45.6 mg) as an internal standard was placed in a 10-cm³ Schlenk tube under an Ar atmosphere and the resulting solution was stirred for 5 min. Then, Ph_2SiH_2 (1.2 mmol, 0.22 dm³) was added *via* a syringe and the reaction mixture was stirred at room temperature for 24 h. After removal of volatiles under vacuum at -30°C , a residue was dissolved in Et_2O (1 cm³). The solution was further stirred for 1 h with K_2CO_3 (1 mg) and MeOH (1 cm³). Total yield (77%) and a ratio (91/9) of cyclohexanone and 2-cyclohexen-1-ol were determined by a GC (Shimadzu CPB10 column, 25 m length, 0.25 mm id).

- (a) G. R. Newkome, C. N. Moorefield and F. Vögtle, *Dendrimers and Dendrons: Concept, Synthesis, Application*, Wiley-VHC, Weinheim, 2001; (b) *Dendrimer and Other Dendritic Polymers*, ed. J. M. J. Fréchet and D. A. Tomalia, J. Wiley & Sons, New York, 2001; (c) S. Hecht and J. M. J. Fréchet, *Angew. Chem., Int. Ed.*, 2001, **40**, 74.
- (a) B. Helmsa and J. M. J. Fréchet, *Adv. Synth. Catal.*, 2006, **348**, 1125; (b) L. J. Twyman, A. S. H. King and I. K. Martin, *Chem. Soc. Rev.*, 2002, **31**, 69; (c) G. E. Oosterom, J. N. H. Reek, P. C. J. Kamer and P. W. N. M. van Leeuwen, *Angew. Chem., Int. Ed.*, 2001, **40**, 1828; (d) D. Astruc and F. Chardac, *Chem. Rev.*, 2001, **101**, 2991; (e) A. M. Caminade, V. Maraval, R. Laurent and J. P. Majoral, *Curr. Org. Chem.*, 2002, **6**, 739; (f) D. Méry and D. Astruc, *Coord. Chem. Rev.*, 2006, **250**, 1965 and references cited therein.
- (a) F. C. de Schryver, T. Vosch, M. Cotlet, M. van der Auwerre, K. Müllen and J. Hofkens, *Acc. Chem. Res.*, 2005, **38**, 514; (b) M. D. Watson, A. Fechtenkotter and K. Müllen, *Chem. Rev.*, 2001, **101**, 1267; (c) A. J. Berresheim, M. Muller and K. Müllen, *Chem. Rev.*, 1999, **99**, 1747 and references cited therein.
- T. Iwasawa, M. Tokunaga, Y. Obora and Y. Tsuji, *J. Am. Chem. Soc.*, 2004, **126**, 6554.
- H. Aoyama, M. Tokunaga, J. Kiyosu, T. Iwasawa, Y. Obora and Y. Tsuji, *J. Am. Chem. Soc.*, 2005, **126**, 10474.
- T. Iwasawa, T. Komano, A. Tajima, M. Tokunaga, Y. Obora, T. Fujihara and Y. Tsuji, *Organometallics*, 2006, **25**, 4665.
- (a) W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290; (b) V. César, S. Bellemin-Lapponnaz and L. H. Gade, *Chem. Soc. Rev.*, 2004, **33**, 619; (c) D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39 and references cited therein.
- Y. J. Boxhall, P. C. Bulman Page, Y. Chan, C. M. Hayman and H. Heaney, *Synlett*, 2003, 997.
- J. P. Wolfe, J. Ahman, J. P. Sadighi, R. A. Singer and S. L. Buchwald, *Tetrahedron Lett.*, 1997, **38**, 6367.
- A. J. Arduengo, III, R. Krafczyk and R. Schmutzler, *Tetrahedron*, 1999, **55**, 14523.
- H. M. J. Wang and I. J. B. Lin, *Organometallics*, 1998, **17**, 972.
- (a) M. V. Baker, S. K. Brayshaw, B. W. Skelton and A. H. White, *Inorg. Chim. Acta*, 2004, **357**, 2841; (b) M. T. Zarka, M. Bortenschlager, K. Wurst, O. Nuyken and R. Weberskirch, *Organometallics*, 2004, **23**, 4817; (c) R. S. Simons, P. Custer, C. A. Tessier and W. J. Youngs, *Organometallics*, 2003, **22**, 1979; (d) A. R. Chianese, X. Li, M. C. Janzen, J. W. Faller and R. H. Crabtree, *Organometallics*, 2003, **22**, 1663.
- 5c** was prepared by Beller *et al.*^{13a} and X-ray structure of **5c** was recently determined.^{13b} (a) A. M. Seayad, K. Selvakumar, M. Ahmed and M. Bellar, *Tetrahedron Lett.*, 2003, **44**, 1679; (b) P. A. Evans, E. W. Baum, A. N. Fazal and M. Pink, *Chem. Commun.*, 2005, 63.
- (a) F. Maseras and K. Morokuma, *J. Comput. Chem.*, 1995, **16**, 1170; (b) S. Humbel, S. Sieber and K. Morokuma, *J. Chem. Phys.*, 1996, **105**, 1959; (c) M. Svensson, S. Humbel, R. D. J. Froese, T. Matsubara, S. Sieber and K. Morokuma, *J. Phys. Chem.*, 1996, **100**, 19357.
- I. Ojima, in *The Chemistry of Organic Silicon Compounds*, ed. S. Patai and Z. Rapport, Wiley, Chichester, 1989.
- (a) I. Ojima and T. Kogure, *Organometallics*, 1982, **1**, 1390; (b) G. Z. Zheng and T. H. Chan, *Organometallics*, 1995, **14**, 70; (c) Y. Kanazawa, Y. Tsuchiya, K. Kobayashi, T. Shiomi, J.-i. Itoh, M. Kikuchi, Y. Yamamoto and H. Nishiyama, *Chem.–Eur. J.*, 2006, **12**, 63.
- T. Fujihara, Y. Obora, M. Tokunaga, H. Sato and Y. Tsuji, *Chem. Commun.*, 2005, **112**, 4526.
- C. J. Hawker and J. M. Fréchet, *J. Am. Chem. Soc.*, 1990, **112**, 7638.
- B. S. Balaji, Y. Obora, D. Ohara, S. Koide and Y. Tsuji, *Organometallics*, 2001, **20**, 5342.