JOC The Journal of Organic Chemistry

Note

A Suzuki Approach to Quinone Based Diarylethene Photochromes

Dinesh G. Patel, Travis B. Mitchell, Shea Douglas Myers, Dorothy A. Carter, and Frank A. Novak *J. Org. Chem.*, Just Accepted Manuscript • DOI: 10.1021/acs.joc.9b02632 • Publication Date (Web): 03 Jan 2020 Downloaded from pubs.acs.org on January 3, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

A Suzuki Approach to Quinone Based Diarylethene Photochromes

Dinesh G. Patel,^{1,*} Travis B. Mitchell,² Shea D. Myers,¹ Dorothy A. Carter,¹ and Frank A. Novak¹ ¹Department of Chemistry, the Pennsylvania State University at Hazleton, Hazleton, Pennsylvania 18202, United States

²Department of Chemistry, the State University of New York at Buffalo, Buffalo, New York 14260-3000, United States

Supporting Information Placeholder

ABSTRACT: Diarylethene photochromes show promise for use in advanced organic electronic and photonic materials with burgeoning considerations for biological applications; however, these compounds typically require UV-light for photoswitching in at least one direction thus limiting their appeal. We here introduce a naphthoquinone-based diarylethene that switches between open and closed forms with visible light. The synthesis of this quinone diarylethene relies on Suzuki methodology allowing for the inclusion of functional groups not otherwise accessible with current synthetic routes.

Diarylethenes (DAEs, Scheme 1) have enjoyed sustained investigation as p-type¹⁻² photochromes with applications in optical data storage devices,³ photoresponsive quantum dots,⁴⁻⁵ photomechanical constructs,⁶⁻⁷ charge transport systems,⁸⁻¹⁰ vehicles for drug delivery,¹¹ photodynamic therapy agents,¹²⁻¹³ and as linkers in metal organic frameworks.¹⁴⁻¹⁷ Many of these systems are the result of work focusing on hexafluorocyclopentene scaffolds¹⁸⁻²¹ and ultraviolet

light (UV) to achieve ring closure (Scheme 1). Given the limitations inherent to the use of UV for photoswitching, research has moved toward developing DAEs that photoisomerize exclusively with visible light.²²⁻²⁴

Quinone based diarylethenes (qDAEs, compound **1a** as an example), in contrast to their hexafluorocyclopentene analogues, have received miniscule attention. The limited contributions to the literature, however, have suggested that quinone based DAEs could exhibit near ideal visible light facilitated switching.²⁵⁻²⁶ When one considers that quinones have great biological importance²⁷⁻²⁸ and rich redox chemistry, appropriately functionalized qDAEs could open the door to photochromes that are biologically²⁹⁻³⁰ and electrochemically relavent.³¹⁻³²



Scheme 1. (Left) Photoisomerization of a quintessential diarylethene. (Right) The structure of various qDAEs shown in the ring open form.

Katsumura³³ synthesized a series of qDAEs but did not examine their optical properties. Subsequently, Deng and Liebeskind²⁵ demonstrated that complete photoconversion from ringopen to ring-closed isomers was not possible with a similar set of qDAEs (for example **1a** – **1c**). Near complete ring closure was, however, realized in the presence of strong Lewis acids. Most recently, both the groups of Tsuda²⁶ and König²⁹ showed that the ring-open and ring-closed forms of qDAEs exhibited stark differences in redox behavior while the work of König²⁹ further showed that qDAEs are capable of influencing biological systems.

Thus far, all reported synthetic approaches toward qDAEs involve Stille coupling conditions. These Stille routes require the use of lithiation and stannylation steps thus limiting

the scope of functional groups that can survive to the final photochrome product. As examples, these functional groups include esters (**1d**), carboxylic acids, phosphonates, and phosphonic acids,¹⁰ which are easily accessible in non-qDAE systems. The limitations on functional groups in turn limit the possible applications of qDAEs. We seek to enter into the literature qDAEs that can switch in both directions with visible light and that contain otherwise synthetically inaccessible functional groups using current methods. These functional groups (our targets were thienyl esters, carboxylic acids, and halides), additionally, are capable of further modification to allow for their potential use in biological systems³⁴⁻³⁵ and main chain photochromic polymers,³⁶⁻³⁸ both of which have received much attention.

Given that borylation, unlike stannylation, can be accomplished without the need for a lithiation step, we supposed that Suzuki methodology would allow access to qDAEs with increasingly diverse functional groups.³⁹ To date, Suzuki reactions have been employed for the synthesis of DAEs containing hexafluoro- and perhydro-cyclopentene components in addition to some benzothiophene examples.⁴⁰⁻⁴⁴ Suzuki methodology, curiously, has seen negligible use in the synthesis of qDAEs. Previous reports have suggested that decomposition of the halogenated quinone coupling partner prevents the reaction.^{39, 45}



Scheme 2. Selected arylated quinones obtained by Suzuki, direct arylation, and Meerwein coupling reactions.

According to the literature, several non-photochromic monoaryl³⁹ and tetraaryl quinone compounds have been synthesized by a Suzuki approach, but 2,3-dichloro-5,6-dicyano-1,4-

benzoquinone (DDQ) was needed as a final step before the desired products were obtained.⁴⁶⁻⁴⁷ When not employed, it is assumed that oxidation by atmospheric oxygen during the work up and purification helped achieve the same results as DDQ.⁴⁸ Direct arylation using boronic acids has also been explored, but the resulting quinones were only monosubstituted, and FeCl₃ was required to re-oxidize the product mixture, which contained a reduced side product⁴⁹ just as in the reports by Langer and coworkers.⁴⁶⁻⁴⁷ The Meerwein arylation⁵⁰ has also been explored but afforded products in low yield. In all of these reports (Scheme 2), only benzene derived boronic acids or azides were used and thus gave non-photochromic qDAEs. We have found no instances of thiophene boronic acids or thiophene boronic acid esters⁵¹ participating in coupling reactions thus affording photochromic qDAEs.

We sought conditions that would obviate the need for DDQ or other oxidants. Anhydrous Suzuki conditions, we hypothesized, would give products without any needed post-synthetic oxidation and would avoid decomposition of the halogenated quinone precursors. Experiments using thallium carbonate⁵² in the absence of water have been examined previously; however, we turned to fluoride as the activating base.⁵³

Ester containing compound **1d** served as our initial target owing to the ease with which the necessary 5-methyl-2-thiophene carboxylic acid (**2**) can be protected as the methyl ester (**3a**, Scheme 3).⁵⁴ We proceeded with the assumption that protection would prevent side reactions and that deprotection would give the desired carboxylic acid functional group. Unfortunately, attempts at deprotection of **1d** using standard methods⁵⁵ either led to decomposition or returned only starting material.



Scheme 3. Synthesis of required boronic acid 5.

Given the difficulty encountered in hydrolyzing the methyl ester, we turned our attention to synthesis of the more labile *t*-butyl analogue (Scheme 3). Starting with compound **2**, we employed the method of Wright⁵⁶ to protect the acid functionality to give *t*-butyl ester **3b**. Using the methods of Hartwig⁵⁷ and Tatsuo,⁵⁸ the 4-position was borylated to give compound **4b**. While Hartwig has shown that boronic acids can be obtained by treating pinacol esters with sodium periodate and aqueous hydrochloric acid, we were concerned that these conditions would lead to unwanted deprotection of the *t*-butyl ester. To our delight, the use of a weaker acid, oxalic acid, in place of hydrochloric acid gave desired compound **5b** with the *t*-butyl protecting group intact. The corresponding methyl ester compounds **4a** and **5a** are obtained using the same procedures as their *t*-butyl analogues.

To access our desired qDAEs (compounds **7** – **9**; Scheme 4), we started by treating 2,3dibromo-1,4-naphthoquinone (**6**)⁵⁹ with compound **5b**, cesium fluoride, and catalytic PEPPSI- IPr^{60} in anhydrous dioxane. Compound **1d** is obtained in similar fashion. While several traditional methods for deprotection of the *t*-butyl group exist, we found that heating **7** in diphenyl ether (bp = 259 °C) for 2h afforded dicarboxylic acid **8**. Treatment of **8** with tetrabutylamine tribromide⁶¹ affords dibromide **9**. Both **8** and **9** can undergo further synthetic manipulation at either the carboxylate or aryl-bromide positions. It is worth noting that compound **9** has previously been obtained by brominating **1a**, which is available through a Stille coupling.²⁵



Scheme 4. Synthetic route to the ester, carboxylic acid, and halogen functionalized photochromic qDAEs in this study.

We firstly examined **1d**, **7**, **8**, and **9** by ¹H NMR spectroscopy. All show the expected number of and multiplicity for peaks in the open forms. Using **1d** as a model compound, irradiation of an NMR sample with short wavelength visible light (405 – 410 nm) gave ~22% of the ring closed form (Supporting Information). This result is similar to the work of Deng and Liebeskind,²⁵ and no attempt was made to force ring closure using strong Lewis acids. Interestingly, the thienyl proton on **1d**, which appears as a broad singlet centered at 7.42 ppm in the open form shifts to 8.29 ppm, now as a sharp singlet, in the closed form. The downfield shift is expected as the conjugation imparted by cyclization allows for effective electron withdrawal by the quinone ring. Similarly, both the ester methyl group and thienyl methyl group experience downfield shifts, though much less pronounced. Compounds **1d** and **7** have nearly identical spectra; only the ester alkyl groups have differing chemical shifts.



Figure 1. The absorption spectrum of **7** in acetone (5.7 x 10^{-5} M) before and after irradiation with 405 - 410 nm light.

UV-Vis spectroscopy of compounds **7** and **8** showed distinct spectra between ring open and ring closed forms. The open form of **7** (Figure 1) has peak absorption at 412 nm and an onset of absorption at ~500 nm, longer than most previously reported qDAEs. Upon irradiation, a new absorption band appears. The closed form of compound **7** exhibits a long wavelength absorption band centered at 612 nm; a second absorption band at 441 nm is due to the presence of both the ring open and ring closed forms given that complete ring closure was not observed by ¹H NMR. Onset of absorption for the closed form of **7** occurs at ~762 nm. Compound **8**, upon ring closure, exhibits significant broadening of the long wavelength absorption band (Supporting Information), presumably due to hydrogen bonding in solution as this broadening is not observed in the spectrum of **7** in the closed form.

The absorption spectrum for brominated compound **9**, in contrast to that of **7** and **8**, shows only minor changes upon irradiation (see Supporting Information); the reason for this diminished response is not immediately apparent though it may be due to some deactivation pathway available to the excited state as a consequence of the heavy atom effect. The long wavelength absorption due to the closed form has a maximum near 560 nm, much shorter than that for compounds **7** and **8**. This hypsochromic shift informs that the electron withdrawing nature of the thienyl substituents greatly impacts the absorption spectrum and electronics of the closed form qDAE. Bromine is not as strong an electron-withdrawing group when compared to ester or carboxylic acid functionalities. Given this structure-property relationship, we theorize that even stronger electron withdrawing groups would lead to more prominent changes in absorption spectra between open and closed forms while also shifting the long-wavelength closed form absorption band further into the near IR.

Single crystals of compound **8** were obtained by slow evaporation of chromatographic fractions leading to the formation of two different morphologies: the photoinactive form as orange prisms (**8a**, Figure 2) and the photoactive form as yellow plates (**8b**, Figure 3). A suitable crystal from each morphology was analyzed by X-ray diffraction.



Figure 2. (a) Molecular structure of qDAE **8a**. Aromatic and alkyl hydrogen atoms are omitted for clarity. (b) Face-to-face $\pi \cdots \pi$ stacking viewed down [1 1 0]. (c) Hydrogen bonding between $\pi \cdots \pi$ stacked dimers viewed down [1 $\overline{1}$ 0].

The orange prisms of **8a** crystallize in the centrosymmetric space group $P\overline{1}$ with one qDAE molecule in the asymmetric unit. The crystal structure of **8a** assumes the parallel conformation leading to photo-inactivity in the solid state.⁶²⁻⁶³ Dimers of this qDAE form parallel to the $[1 \overline{1} 0]$ through face-to-face $\pi \cdots \pi$ stacking between the quinoid and benzenoid rings separated by a distance of 3.5336(5) Å (Figure 2b). Each dimer hydrogen bonds to four neighboring dimers through the carboxylic acid groups forming 2-D sheets (Figure 2c, Table S1) that possess solvent accessible pores parallel to $[1 \overline{1} 0]$. Attempts to model the electron density within the pores, presumably arising from highly disordered solvent molecules, were unsuccessful. The electron density was subsequently removed using the SQUEEZE routine in PLATON.⁶⁴

The yellow plates of **8b** crystallize in the non-centrosymmetric space group $Pna2_1$ with one qDAE molecule in the asymmetric unit. The crystal structure of **8b** assumes the anti-parallel

conformation with C–C separation distance of 3.284(6) Å for the carbon atoms involved in photocyclization; this value is characteristic of photoactive DAEs in the solid state.⁶⁵ The solid-state photochromism of **8b** will be examined in future work. Each qDAE molecule parallel to the *n*-glide plane (or molecule parallel to $[0 \frac{1}{2} \frac{1}{2}]$) undergoes an electrostatic interaction between the carbonyl carbon adjacent to the thiophene ring of one molecule and the quinoidal oxygen on the backbone of a neighboring molecule (Figure 3b, C14…O1ⁱ = 3.048(6) Å, (i) = $\frac{1}{2}$ -x, $-\frac{1}{2}$ -y, $\frac{1}{2}$ +z).⁶⁶ In addition, each **8b** qDAE molecule hydrogen bonds with adjacent qDAE molecules through the carboxylic acid groups forming infinite chains with two-fold helical symmetry along the crystallographic *a* axis (Figure 3c, Table S2).⁶⁷



Figure 3. (a) Molecular structure of qDAE **8b**. Aromatic and alkyl hydrogen atoms are omitted for clarity. (b) Electrostatic interaction between quinoidal oxygen and carboxyl carbon. (c) Infinite chain imposed by hydrogen bonds between carboxylic acids on adjacent qDAE molecules.

In conclusion, we have demonstrated the use of Suzuki coupling methodology to obtain photochromic quinone-based diarylethenes with previously inaccessible functional groups. These compounds show ideal visible light absorption characteristics and photoswitchability and are

The Journal of Organic Chemistry

amenable to further synthetic modification. Crystal structure data on a carboxylic acid terminated qDAE show extensive hydrogen bonding and that potentially photoactive crystals can be obtained. Future work will focus on computations, photocrystallography, and the realization of qDAE based polymers and metal-organic frameworks.

Experimental Section

All reagents were obtained from commercial sources and used as received unless otherwise noted in the text. Flash column chromatography was accomplished using silica gel purchased from Fisher Scientific (230 – 400 mesh, grade 60, catalog number S825-212). Polyester backed TLC silica plates (grade 60 with UV indicator) were obtained from EMD Millipore (catalog number 1.05735.0001). NMR spectra were obtained on a Bruker AVIII-500 spectrometer operating at 500 MHz for ¹H NMR spectra and 125 MHz for ¹³C NMR spectra or on a Bruker DRX-400 spectrometer operating at 400 MHz for ¹H NMR spectra and 100 MHz for ¹³C NMR spectra; the ¹H NMR spectrum for **1d** after irradiation (Supporting Information) was acquired on a Varian Inova spectrometer operating at 500 MHz. NMR peaks are referenced to residual solvent signals⁶⁸ in either CDCl3 (7.26 ppm for ¹H, 77.0 ppm for ¹³C) or acetone-d₆ (2.05 ppm for ¹H, 29.84 for ¹³C). Solution UV-Vis experiments were preformed in methylene chloride or acetone using a Shimadzu UV-1800 spectrometer. Solvents for these experiments were used as received and not degassed prior to use. Mass spectrometric analyses were performed at the Pennsylvania State University Proteomics and Mass Spectrometry Core Facility, University Park, PA with high-resolution mass spectra (HRMS) obtained on a Waters O-TOF quadrupole time-offlight mass spectrometer. Data were obtained in either positive or negative mode as denoted in each compound's characterization. X-ray diffraction data was collected on either a home source or a synchrotron source as described in the supporting information.

5-methyl-2-thiophene carboxylic acid methyl ester (**3a**). The procedure by Rückle⁵⁴ was followed; in a 500 mL round bottom flask was placed methanol (250 mL). Concentrated sulfuric acid (11 mL, 18 M) was added slowly and with vigorous stirring. Heat was evolved during the addition; without cooling, 5-methyl-2-thiophene carboxylic acid (9.300 g, 65.42 mmol) was added in a single portion followed by reflux for 24 hours. Upon cooling to room temperature, neutralization was achieved by addition of small portions of sodium carbonate. Some solids were present after complete neutralization and were removed by suction filtration. The filtrate was concentrated to approximately 30 mL, then diluted with water (200 mL) and the product extracted with hexanes (3 x 100 mL). The combined hexane phases were dried over magnesium sulfate, filtered, and solvent removed under reduced pressure to afford a faint yellow oil (8.910 g, 87%). ¹H NMR (400 MHz, chloroform-*d*), δ 7.59 (d, 1H, *J* = 3.8 Hz), 6.74 (d, 1H, *J* = 3.3 Hz), 3.83 (s, 3H), 2.49 (s, 3H); ¹³C{¹H} NMR (100 MHz, chloroform-*d*), δ 162.6, 147.8, 133.8, 130.8, 126.3, 51.8, 15.6. HRMS (ESI-TOF) *m/z*: [M + H]⁺ calcd for C₇H₉O₂S 157.0323; found 157.0318.

5-methyl-2-thiophene carboxylic acid *t*-butyl ester (**3b**). This compound has been previously reported,⁶⁹ obtained by a different route than presented here, and has not been fully characterized; based on the procedure by Wright and coworkers,⁵⁶ a 500 mL round bottom flask was charged with magnesium sulfate (42.33 g, 351.7 mmol), methylene chloride (200 mL), and concentrated sulfuric acid (6.90 g, 3.75 mL, 70.3 mmol). After stirring the suspension for 5 minutes, 5-methyl-2-thiophene carboxylic acid (2, 10.000 g, 70.33 mmol) was added in a single portion and immediately followed by the addition of *t*-butanol (20.85 g, 28 mL, 281.3 mmol). The reaction was allowed to stir, tightly capped (a rubber septum was used), for 48 hours. At this point, the magnesium sulfate was removed by suction filtration and the filter cake was washed with hexanes (150 mL). Owing to the small particulate size of the magnesium sulfate resulting from stirring for 48 h, the filtration process took several hours. The washings and filtrate were

combined, and then washed with aqueous sodium hydroxide (1 M, 2 x 40 mL). Upon drying the remaining organic phase over magnesium sulfate and filtering, solvent was removed under reduced pressure. The resulting oil was chromatographed on silica eluting with 100% hexanes to 10% ethyl acetate (EtOAc) in hexanes. Removal of solvent afforded a clear, faint yellow oil (9.348 g, 67%). ¹H NMR (500 MHz, CDCl₃), δ 7.52 (d, 1H, *J* = 3.7 Hz), 6.72 (d, 1H, *J* = 3.7 Hz), 2.49 (s, 3H), 1.55 (s, 9H); ¹³C{¹H} NMR (125 MHz, CDCl₃), δ 161.5, 147.0, 133.1, 133.0, 126.0, 81.2, 28.1, 15.6. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ calcd for C₁₀H₁₄O₂SNa 221.0612; found 221.0602.

Methyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2-carboxylate (4a). As in the procedure by Sawamura and coworkers,⁷⁰ a 250 mL Schlenk tube was charged with compound **3a** (7.000 g, 44.81 mmol), bis(pinacolato)diboron (B₂pin₂, 6.828 g, 26.89 mmol), and 4,4'-di-*tert*-butylbipyridine (dtbpy, 0.024 g, 0.09 mmol). Heptane (40 mL) was added and the solution subjected to three freeze-pump-thaw cycles followed by a nitrogen backfill. Upon (1,5-cvclooctadiene)(methoxy)iridium(I) warming room temperature, dimer to ([Ir(COD)(OMe)]₂, 0.030 g, 0.04 mmol) was added under a stream of nitrogen. After approximately 1 – 2 minutes, the initially yellow solution turned dark brown; heating at 100 °C was carried out for 24 h. Upon cooling, a white precipitate was visible. The reaction was transferred to a separatory funnel and a 1:1 solution of hexanes/methylene chloride was added. This organic phase was washed with potassium acetate (0.10 M aqueous, 2 x 50 mL) and brine (50 mL). The remaining organic phase was passed through a short plug of silica eluting with 1:1 hexanes/methylene chloride. A white solid was obtained upon removal of solvent (12.356 g. 98%); spectral data matched that in the literature.⁷⁰ ¹H NMR (400 MHz, CDCl₃), δ 7.95 (s, 1H), 3.86 (s, 3H), 2.72 (s, 3H), 1.34 (s, 12H); ¹³C{¹H} NMR (100 MHz, CDCl₃), 162.7, 159.7, 140.4, 130.1, 129.1, 83.6, 51.9, 24.9, 16.3. HRMS (ESI-TOF) m/z: [M + H]⁺ calcd for C₁₃H₂₀BO₄S 283.1178; found 283.1167.

tert-butyl 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)thiophene-2carboxylate (**4b**). Following a previously reported procedure,⁵⁷⁻⁵⁸ in a 250 mL Schlenk flask was combined compound **3b** (5.000 g, 25.21 mmol), B₂pin₂ (3.665 g, 0.57 mmol), dtbpy (0.010 g, 0.04 mmol), and heptane (50 mL). The solution was degassed by three freeze-pump-thaw cycles and then backfilled with nitrogen. Under a stream of nitrogen, [Ir(COD)OMe]₂ (0.012 g, 0.02 mmol) was added quickly and the reaction heated to 100 °C and followed until TLC (silica, 5% EtOAc in hexanes) showed that no starting material remained. The reaction solution was passed through a short silica column followed by elution with 100% hexanes to 5% EtOAc in hexanes. The fractions containing product were collected affording a white solid upon removal of solvent (6.743 g, 82%). ¹H NMR (500 MHz, CDCl₃), δ 7.80 (s, 1H), 2.67 (s, 3H), 1.55 (s, 9H), 1.32 (s, 12 H); ¹³C{¹H} NMR (125 MHz, CDCl₃), δ 161.6, 159.1, 139.1, 132.7, 128.7, 83.4, 81.2, 28.2, 24.8, 16.3. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ calcd for C₁₆H₂₅BO₄SNa 347.1464; found 347.1437.

(5-(methoxycarbonyl)-2-methylthiophen-3-yl)boronic acid (**5a**). A modification of the procedure by Hartwig and coworkers was followed.⁵⁷ A 250 mL flask was charged with compound **4b** (7.000 g, 24.81 mmol), oxalic acid dihydrate (6.253 g, 49.62 mmol), water (30 mL), and acetonitrile (150 mL). After stirring for 1 min, NaIO₄ (7.960, 37.21 mmol) was added in a single portion and the resulting mixture stirred for 2 hours. During this time, the reaction increased in cloudiness; a solid mass was apparent when pouring the reaction into a separatory funnel. The material in the separatory funnel was diluted with water (150 mL) and extracted with diethyl ether (5 x 100 mL). A white precipitate remained at the aqueous/organic interface during the extraction process. The combined organic phases were washed with brine (100 mL), dried over magnesium sulfate, filtered, and solvent removed under reduced pressure. The obtained solid was washed with hexanes and the resulting suspension collected by suction filtration to afford a white solid (4.280 g, 86%). ¹H NMR (500 MHz, acetone-*d*₆), δ 7.97 (s, 1H), 7.22 (br s, 2H),

The Journal of Organic Chemistry

3.80 (s, 3H), 2.71 (s, 3H); ¹³C{¹H} NMR (125 MHz, acetone-*d*₆), 163.0, 159.9, 158.6, 141.1, 130.0, 52.1, 16.4. HRMS (ESI-TOF) *m*/*z*: [M − H]⁻ calcd for C₇H₈BO₄S 199.0238; found 199.0229.

(5-(tert-butoxycarbonyl)-2-methylthiophen-3-yl)boronic acid (**5b**). A modification of the procedure by Hartwig and coworkers was followed.⁵⁷ In a 100 mL round bottom flask was combined compound **4b** (3.00 g, 9.25 mmol), oxalic acid dihydrate (2.332 g, 18.50 mmol), acetonitrile (50 mL), and water (10 mL). After stirring for 1 minute NaIO₄ (2.968 g, 13.88 mmol) was added in a single portion. After stirring at room temperature for 7 h, TLC (silica, 1:1 hexanes/EtOAc) showed a new spot of low R_f (~ 0.1 – 0.2) and some starting material; stirring was continued for an additional 9 h at which point TLC showed no starting material. Upon dilution with water (50 mL), the product was extracted with diethyl ether (3 x 50 mL). The ether phases were washed with brine (50 mL), dried over MgSO₄; upon removal of solvent a white solid was obtained. This solid was washed with hexanes, water, and dried under vacuum (1.880 g, 84%). ¹H NMR (500 MHz, acetone-*d*₆), δ 7.90 (s, 1H), 7.19 (s, 2H), 2.69 (s, 3H), 1.53 (s, 9H); ¹³C{¹H} NMR (125 MHz, CDCl₃), δ 162.0, 158.0, 140.4, 132.3, 81.4, 28.4, 16.4. HRMS (ESI-TOF) *m/z*: [M – H]⁻ calcd for C₁₀H₁₄BO₄S [M-H]⁻ 241.0708; found 241.0698.

2,3-dibromonaphthalene-1,4-dione (6). According to the slightly modified procedure by Anuratha and coworkers,⁷¹ a 250 mL round bottom flask was charged with 1,4-naphthoquinone (5.000 g, 31.62 mmol) and sodium acetate (26.934 g, 316.15 mmol). After holding under high vacuum for 30 minutes and then backfilling with nitrogen, acetic acid (AcOH, 125 mL) was added. With vigorous stirring, bromine (15.157 g, 4.88 mL, 94.85 mmol) was introduced in portions over 2 – 3 minutes. After complete addition of bromine, the reaction was refluxed for 2 hours followed by cooling to room temperature overnight. The resulting orange suspension was poured into water (300 mL) and the solids collected by suction filtration. The resulting filter cake was collected and suspended in a 5 wt. % sodium thiosulfate solution with gentle swirling; the solids were again collected and suspended, with stirring, in hot ethanol. Upon cooling to room temperature, a bright yellow solid was collected by suction filtration and dried under vacuum (6.656 g, 67%). ¹H NMR (500 MHz, CDCl₃), δ 8.19 (dd, 2H, *J* = 5.7 Hz, 3.3 Hz), 7.79 (dd, 2H, 5.8 Hz, 3.3 Hz). ¹³C{¹H} NMR (125 MHz, CDCl₃), δ 175.8, 142.6, 134.5, 130.8, 128.2. HRMS (ESI-TOF) *m/z*: [M]⁻ calcd for C₁₀H₄Br₂O₂ 313.8578; found 313.8615.

4,4'-(1,4-dioxo-naphthalene-2,3-diyl)bis(5-methylthiophene-2-carboxylic acid tert-butyl ester) (7). In a 50 mL Schlenk tube was combined Compound **6** (0.746 g, 2.36 mmol), compound **5b** (1.200 g, 4.96 mmol), cesium fluoride (1.510 g, 9.91 mmol), and 1,4-dioxane (10 mL). The yellow mixture was subjected to three freeze-pump-thaw cycles, then backfilled with nitrogen. PEPPSI-IPr (0.064 g, 0.09 mmol) was quickly added under a stream of nitrogen and the reaction refluxed for 15 h, then cooled to room temperature. After dilution with EtOAc (200 mL), the organic phase was washed with water (3 x 50 mL) followed by brine (50 mL). The remaining organic phase was dried over MgSO₄, then adsorbed onto silica followed by chromatography on silica eluting with 100% hexanes to 25% EtOAc in hexanes. The pure fractions were collected; upon removal of solvent under reduced pressure a dark green solid was obtained (0.954 g, 72%). ¹H NMR (500 MHz, CDCl₃), δ 8.18 (dd, 2H, *J* = 5.7 Hz, 3.3 Hz), 7.81 (dd, 2H, *J* = 5.7 Hz, 3.3 Hz), 7.30 (br s, 2H), 2.08 (s, 6H), 1.53 (s, 18H); ¹³C{¹H} NMR (125 MHz, CDCl₃), 183.6, 161.0, 145.9, 141.8, 134.9, 134.1, 132.1, 132.0, 130.6, 126.8, 81.7, 28.2, 15.2. HRMS (ESI-TOF) *m/z*: [M + Na]⁺ calcd for C₃₀H₃₀O₆S₂Na 573.1376; found 573.1355.

4,4'-(1,4-dioxonaphthalene-2,3-diyl)bis(5-methylthiophene-2-carboxylic acid) (8). In a 50 mL flask with reflux condenser was combined diphenyl ether (15 mL) and compound 7 (0.800 g, 1.45 mmol). The dark green solution was refluxed (259 °C) for 2h. After cooling, the product was precipitated by slow addition of the reaction solution to stirring hexanes. The resulting solid was collected by suction filtration, dissolved in acetone, then adsorbed onto silica followed by

 chromatography on silica eluting with 1:1 hexanes/EtOAc to 100% EtOAc to 5% AcOH in EtOAc. The photochromic fractions were collected; upon removal of solvent under reduced pressure, a dark green solid was obtained (0.362 g, 57%). ¹H NMR (400 MHz, acetone- d_6), δ 8.15 (dd, 2H, *J* = 5.7 Hz, 3.3 Hz), 7.92 (dd, 2H, *J* = 5.7 Hz, 3.3 Hz), 7.54 (s, 2H), 2.24 (s, 6H); ¹³C{¹H} NMR (100 MHz, acetone- d_6), 184.3, 162.8, 147.3, 142.8, 137.1, 134.8, 133.2, 132.7, 130.8, 127.1, 15.1. HRMS (ESI-TOF) *m/z*: [M + H]⁺ calcd for C₂₂H₁₅O₆S₂ 439.0310; found 439.0281.

2,3-bis(5-bromo-2-methylthiophen-3-yl)naphthalene-1,4-dione (9). In a 35 mL pressure tube was combined **8** (0.100 g, 0.23 mmol), anhydrous K_3PO_4 (0.102 g, 0.48 mmol), and tetrabutylammonium tribromide (0.231 g, 0.48 mmol). Acetonitrile (5 mL) was added and the capped tube was heated to 100 °C. During the first few minutes, the reaction clarified slightly and went from a white-green mixture to orange. After stirring for 24 h, the reaction was cooled to room temperature and diluted with methylene chloride (50 mL). The cloudy mixture was then washed with water (2 x 30 mL), saturated aqueous NaHCO₃ (30 mL), sodium thiosulfate (10%, 30 mL), and finally with brine (30 mL). The remaining organic phase was dried over magnesium sulfate, filtered, and adsorbed onto silica followed by chromatography on silica eluting with 100% hexanes to 20% EtOAc in hexanes. The vellow band was collected and those fractions deemed pure were combined and solvent removed to give a brown solid with a metallic luster (0.043 g)37%). ¹H NMR (400 MHz, CDCl₃), δ 8.17 (dd, 2H, J = 5.8 Hz, 3.3 Hz), 7.80 (dd, 2H, J = 5.8 Hz, 3.3 Hz,), 6.64 (s, 2H), 2.02 (s, 6H); ¹³C{¹H} NMR (100 MHz, CDCl₃), 183.6, 141.5, 140.9, 134.1, 132.0, 131.5, 130.4, 126.8, 108.0, 14.8. HRMS (ESI-TOF) m/z: [M + Na]⁺ calcd for C₂₀H₁₂Br₂O₂S₂Na 528.8543; found 528.8516.

4,4'-(1,4-dioxo-naphthalene-2,3-diyl)bis(5-methylthiophene-2-carboxylate methyl ester) (1d). Compounds 6 (0.376 g, 1.19 mmol) and 5a (0.500 g, 2.50 mmol) were combined in a 100 mL Schlenk tube followed by the addition of cesium fluoride (0.759 g, 5.00 mmol) and 1,2dimethoxyethane (10 mL). The mixture was degassed by three freeze-pump-thaw cycles, then backfilled with nitrogen. Upon warming to room temperature, and under a stream of nitrogen, PEPPSI-IPr (0.016 g, 0.02 mmol) was quickly added. After refluxing for 18 h, TLC (silica, 10% EtOAc in hexanes) showed a photochromic spot of low R_f. Removal of solvent led to a brown solid that was adsorbed onto silica and chromatographed on silica eluting with 100% hexanes to 30% EtOAc in hexanes. The photochromic spots were collected and solvent removed to afford a yellow powdery solid that was pure by ¹H NMR (0.514 g, 57%). Alternatively, purification can be achieved by passing the crude product (after aqueous workup extracting with chloroform) through a short plug of silica eluting with chloroform. The solid obtained after removal of solvent is heated in methanol until boiling. After storing cold overnight, the remaining solids can be collected by suction filtration, washed with cold methanol, and dried under high vacuum. In this case, a greenish solid is obtained in lower yield (36% yield) than achieved by chromatography but at the same level of purity. ¹H NMR (500 MHz, CDCl₂), δ 8.17 (dd, 2H, *J* = 5.7, 3.3 Hz), 7.81 (dd, 2H, I = 5.8, 3.3 Hz), 7.42 (br s, 2H), 3.84 (s, 6H), 2.09 (s, 6H); ¹³C{¹H} NMR (125 MHz, CDCl₃) 183.5. 162.1, 146.6, 141.6, 135.6, 134.2, 131.9, 130.8, 129.8, 126.8, 52.1, 15.3. HRMS (ESI-TOF) m/z: [M + Na]⁺ calcd for C₂₄H₁₈O₆S₂Na 489.0443; found 489.0436.

Supporting Information.

The Supporting Information is available free of charge on the ACS Publications website at DOI: Copies of ¹H and ¹³C NMR data for all compounds, UV-Vis data for compounds **8** and **9**, hydrogen bonding tables for **8a** and **8b**, and additional experimental information (PDF). Crystal structure of **8a** (cif)

Crystal structure of **8b** (cif)

AUTHOR INFORMATION

Corresponding Author

*E-mail: dgp15@psu.edu

ORCID

Dinesh G. Patel: 0000-0003-0733-5383

Shea D. Myers: 0000-0002-9104-5305

ACKNOWLEDGMENTS

D.G.P. acknowledges support from Penn State Hazleton in the form of a research development grant. Work by T.B.M. was supported by the National Science Foundation under Grant No. DMR-1455039. Access to NSF's ChemMatCARS Sector 15 is supported by the Divisions of Chemistry (CHE) and Materials Research (DMR), National Science Foundation, under grant number NSF/CHE- 1834750. Use of the Advanced Photon Source, an Office of Science User Facility operated for the U.S. Department of Energy (DOE) Office of Science by Argonne National Laboratory, was supported by the U.S. DOE under Contract No. DE-AC02-06CH11357. The authors thank Dr. Tatiana Laremore for help with sample preparation, data interpretation, and providing access to the Pennsylvania State University mass spectrometry facility. Dr. Carlos Pacheco is acknowledged for assistance in the acquisition of NMR data, and Professor Jason Benedict is acknowledged for providing advice on the crystal structure analyses. Dr. Yu-Sheng Chen and Dr. SuYin Grass Wang of NSF's ChemMatCARS are also thanked for their time, expertise, and discussions with regards to the acquired beamline data.

REFERENCES

 1. Dürr, H.; Bouas-Laurent, H., *Photochromism : molecules and systems*. Rev. ed.; Elsevier: Amsterdam ; Boston, 2003.

2. Kobatake, S.; Terakawa, Y. Acid-induced photochromic system switching of diarylethene derivatives between P- and T-types. *Chem. Commun.* **2007**, (17), 1698-1700.

3. Corredor, C. C.; Huang, Z. L.; Belfield, K. D. Two-Photon 3D Optical Data Storage via Fluorescence Modulation of an Efficient Fluorene Dye by a Photochromic Diarylethene. *Adv. Mater.* **2006**, *18* (21), 2910-2914.

4. Díaz, S. A.; Gillanders, F.; Jares-Erijman, E. A.; Jovin, T. M. Photoswitchable semiconductor nanocrystals with self-regulating photochromic Förster resonance energy transfer acceptors. *Nat. Commun.* **2015**, *6*, 6036.

Díaz, S. A.; Menéndez, G. O.; Etchehon, M. H.; Giordano, L.; Jovin, T. M.; Jares-Erijman,
 E. A. Photoswitchable Water-Soluble Quantum Dots: pcFRET Based on Amphiphilic
 Photochromic Polymer Coating. *ACS Nano* 2011, 5 (4), 2795-2805.

6. Morimoto, M.; Irie, M. A Diarylethene Cocrystal that Converts Light into Mechanical Work. J. Am. Chem. Soc. 2010, 132 (40), 14172-14178.

7. Toccafondi, C.; Occhi, L.; Cavalleri, O.; Penco, A.; Castagna, R.; Bianco, A.; Bertarelli, C.; Comoretto, D.; Canepa, M. Photochromic and photomechanical responses of an amorphous

diarylethene-based polymer: a spectroscopic ellipsometry investigation of ultrathin films. *Journal of Materials Chemistry C* **2014**, *2* (23), 4692-4698.

8. Kim, Y.; Hellmuth, T. J.; Sysoiev, D.; Pauly, F.; Pietsch, T.; Wolf, J.; Erbe, A.; Huhn, T.; Groth, U.; Steiner, U. E.; Scheer, E. Charge Transport Characteristics of Diarylethene Photoswitching Single-Molecule Junctions. *Nano Lett.* **2012**, *12* (7), 3736-3742.

9. Wang, Q.; Diez-Cabanes, V.; Dell'Elce, S.; Liscio, A.; Kobin, B.; Li, H.; Brédas, J.-L.; Hecht, S.; Palermo, V.; List-Kratochvil, E. J. W.; Cornil, J.; Koch, N.; Ligorio, G. Dynamically Switching the Electronic and Electrostatic Properties of Indium–Tin Oxide Electrodes with Photochromic Monolayers: Toward Photoswitchable Optoelectronic Devices. *ACS Appl. Nano Mater.* **2019**, *2* (2), 1102-1110.

Wang, Q.; Ligorio, G.; Diez-Cabanes, V.; Cornil, D.; Kobin, B.; Hildebrandt, J.; Nardi, M.
 V.; Timpel, M.; Hecht, S.; Cornil, J.; List-Kratochvil, E. J. W.; Koch, N. Dynamic Photoswitching of Electron Energy Levels at Hybrid ZnO/Organic Photochromic Molecule Junctions. *Adv. Funct. Mater.* 2018, 28 (28), 1800716.

11. Al-Atar, U.; Fernandes, R.; Johnsen, B.; Baillie, D.; Branda, N. R. A Photocontrolled Molecular Switch Regulates Paralysis in a Living Organism. *J. Am. Chem. Soc.* **2009**, *131* (44), 15966-15967.

 Hou, L.; Zhang, X.; Pijper, T. C.; Browne, W. R.; Feringa, B. L. Reversible Photochemical Control of Singlet Oxygen Generation Using Diarylethene Photochromic Switches. *J. Am. Chem. Soc.* 2014, 136 (3), 910-913. 13. Okuda, J.-y.; Tanaka, Y.; Kodama, R.; Sumaru, K.; Morishita, K.; Kanamori, T.; Yamazoe, S.; Hyodo, K.; Yamazaki, S.; Miyatake, T.; Yokojima, S.; Nakamura, S.; Uchida, K. Photoinduced cytotoxicity of a photochromic diarylethene via caspase cascade activation. *Chem. Commun.* **2015**, *51* (54), 10957-10960.

Luo, F.; Fan, C. B.; Luo, M. B.; Wu, X. L.; Zhu, Y.; Pu, S. Z.; Xu, W.-Y.; Guo, G.-C.
 Photoswitching CO2 Capture and Release in a Photochromic Diarylethene Metal–Organic
 Framework. *Angew. Chem. Int. Ed.* 2014, 53 (35), 9298-9301.

Furlong, B. J.; Katz, M. J. Bistable Dithienylethene-Based Metal–Organic Framework
 Illustrating Optically Induced Changes in Chemical Separations. J. Am. Chem. Soc. 2017, 139
 (38), 13280-13283.

16. Patel, D. G.; Walton, I. M.; Cox, J. M.; Gleason, C. J.; Butzer, D. R.; Benedict, J. B. Photoresponsive porous materials: the design and synthesis of photochromic diarylethene-based linkers and a metal–organic framework. *Chem. Commun.* **2014**, *50* (20), 2653-2656.

17. Walton, I. M.; Cox, J. M.; Coppin, J. A.; Linderman, C. M.; Patel, D. G.; Benedict, J. B. Photo-responsive MOFs: light-induced switching of porous single crystals containing a photochromic diarylethene. *Chem. Commun.* **2013**, *49* (73), 8012-8014.

18. Gilat, S. L.; Kawai, S. H.; Lehn, J.-M. Light-triggered electrical and optical switching devices. J. Chem. Soc., Chem. Commun. 1993, (18), 1439-1442.

19. Hanazawa, M.; Sumiya, R.; Horikawa, Y.; Irie, M. Thermally irreversible photochromic systems. Reversible photocyclization of 1,2-bis (2-methylbenzo[b]thiophen-3-yl)perfluorocyclocoalkene derivatives. *J. Chem. Soc., Chem. Commun.* **1992,** (3), 206-207.

20. Irie, M.; Sakemura, K.; Okinaka, M.; Uchida, K. Photochromism of Dithienylethenes with Electron-Donating Substituents. *J. Org. Chem.* **1995**, *60* (25), 8305-8309.

21. Kawai, S. H.; Gilat, S. L.; Ponsinet, R.; Lehn, J.-M. A Dual-Mode Molecular Switching Device: Bisphenolic Diarylethenes with Integrated Photochromic and Electrochromic Properties. *Chem. Eur. J.* **1995**, *1* (5), 285-293.

22. Fukaminato, T.; Hirose, T.; Doi, T.; Hazama, M.; Matsuda, K.; Irie, M. Molecular Design Strategy toward Diarylethenes That Photoswitch with Visible Light. *J. Am. Chem. Soc.* **2014**, *136* (49), 17145-17154.

Sumi, T.; Kaburagi, T.; Morimoto, M.; Une, K.; Sotome, H.; Ito, S.; Miyasaka, H.; Irie, M.
Fluorescent Photochromic Diarylethene That Turns on with Visible Light. *Org. Lett.* 2015, 17 (19), 4802-4805.

24. Xu, J.; Volfova, H.; Mulder, R. J.; Goerigk, L.; Bryant, G.; Riedle, E.; Ritchie, C. Visible-Light-Driven "On"/"Off" Photochromism of a Polyoxometalate Diarylethene Coordination Complex. J. Am. Chem. Soc. **2018**, 140 (33), 10482-10487.

25. Deng, X.; Liebeskind, L. S. A Contribution to the Design of Molecular Switches: Novel Acid-Mediated Ring-Closing–Photochemical Ring-Opening of 2,3-Bis(heteroaryl)quinones (Heteroaryl = Thienyl, Furanyl, Pyrrolyl). *J. Am. Chem. Soc.* **2001**, *123* (31), 7703-7704.

26. Saito, E.; Ako, T.; Kobori, Y.; Tsuda, A. Switching of the π -electronic conjugations in the

reduction of a dithienylethene-fused p-benzoquinone. RSC Advances 2017, 7 (5), 2403-2406.

27. Kumagai, Y.; Shinkai, Y.; Miura, T.; Cho, A. K. The Chemical Biology of Naphthoquinones and Its Environmental Implications. *Annu. Rev. Pharmacool. Toxicol.* **2012**, *52* (1), 221-247.

28. Decosterd, L. A.; Parsons, I. C.; Gustafson, K. R.; Cardellina, J. H.; McMahon, J. B.; Cragg, G. M.; Murata, Y.; Pannell, L. K.; Steiner, J. R. HIV inhibitory natural products. 11. Structure, absolute stereochemistry, and synthesis of conocurvone, a potent, novel HIV-inhibitory naphthoquinone trimer from a Conospermum sp. *J. Am. Chem. Soc.* **1993**, *115* (15), 6673-6679.

29. Simeth, N. A.; Kneuttinger, A. C.; Sterner, R.; König, B. Photochromic coenzyme Q derivatives: switching redox potentials with light. *Chem. Sci.* **2017**, *8* (9), 6474-6483.

30. Wilson, D.; Branda, N. R. Turning "On" and "Off" a Pyridoxal 5'-Phosphate Mimic Using Light. *Angew. Chem. Int. Ed.* **2012**, *51* (22), 5431-5434.

31. Liu, Z.; Wang, H. I.; Narita, A.; Chen, Q.; Mics, Z.; Turchinovich, D.; Kläui, M.; Bonn,
M.; Müllen, K. Photoswitchable Micro-Supercapacitor Based on a Diarylethene-Graphene
Composite Film. J. Am. Chem. Soc. 2017, 139 (28), 9443-9446.

32. Xie, N.; Chen, Y. Combination of fluorescent switch and electrochemical switch based on a photochromic diarylethene. *New J. Chem.* **2006**, *30* (11), 1595-1598.

33. Yoshida, S.; Kubo, H.; Saika, T.; Katsumura, S. Synthesis of 2,3-Diarylquinone by Palladium Catalyzed Cross-Coupling of Dibromoquinones with Heteroarylstannanes. *Chem. Lett.* **1996**, *25* (2), 139-140.

The Journal of Organic Chemistry

34. Roubinet, B.; Weber, M.; Shojaei, H.; Bates, M.; Bossi, M. L.; Belov, V. N.; Irie, M.; Hell, S. W. Fluorescent Photoswitchable Diarylethenes for Biolabeling and Single-Molecule Localization Microscopies with Optical Superresolution. *J. Am. Chem. Soc.* **2017**, *139* (19), 6611-6620.

35. Roubinet, B.; Bossi, M. L.; Alt, P.; Leutenegger, M.; Shojaei, H.; Schnorrenberg, S.; Nizamov, S.; Irie, M.; Belov, V. N.; Hell, S. W. Carboxylated Photoswitchable Diarylethenes for Biolabeling and Super-Resolution RESOLFT Microscopy. *Angew. Chem. Int. Ed.* **2016**, *55* (49), 15429-15433.

36. Harvey, C. P.; Tovar, J. D. Main-chain photochromic conducting polymers. *Polymer Chemistry* **2011**, *2* (12), 2699-2706.

37. Peters, G. M.; Tovar, J. D. Pendant Photochromic Conjugated Polymers Incorporating a
Highly Functionalizable Thieno[3,4-b]thiophene Switching Motif. *J. Am. Chem. Soc.* 2019, 141
(7), 3146-3152.

38. Choi, H.; Lee, H.; Kang, Y.; Kim, E.; Kang, S. O.; Ko, J. Photochromism and Electrical Transport Characteristics of a Dyad and a Polymer with Diarylethene and Quinoline Units. *J. Org. Chem.* **2005**, *70* (21), 8291-8297.

39. Best, W. M.; Sims, C. G.; Winslade, M. Palladium-Catalysed Cross Coupling of Arylboronic Acids with 2-Chloro-1,4-naphthoquinones: the Synthesis of 2-Aryl- and 2,3-Bisaryl-1,4-naphthoquinones. *Aust. J. Chem.* **2001**, *54* (6), 401-404.

40. Hiroto, S.; Suzuki, K.; Kamiya, H.; Shinokubo, H. Synthetic protocol for diarylethenes through Suzuki-Miyaura coupling. *Chem. Commun.* **2011**, *47* (25), 7149-7151.

41. Herder, M.; Eisenreich, F.; Bonasera, A.; Grafl, A.; Grubert, L.; Pätzel, M.; Schwarz, J.; Hecht, S. Light-Controlled Reversible Modulation of Frontier Molecular Orbital Energy Levels in Trifluoromethylated Diarylethenes. *Chem. Eur. J.* **2017**, *23* (15), 3743-3754.

42. Wang, S.; Zhou, Y.; Zhu, L.; Zhang, J.; Zou, Q.; Zeng, T.; Chen, W. A photochromic prototype based on difurylperhydrocyclopentene with remarkable photoswitching behavior and in vivo application. *Chem. Commun.* **2017**, *53* (69), 9570-9573.

43. Fukumoto, S.; Nakashima, T.; Kawai, T. Photon-Quantitative Reaction of a Dithiazolylarylene in Solution. *Angew. Chem. Int. Ed.* **2011**, *50* (7), 1565-1568.

44. Jeong, Y.-C.; Gao, C.; Lee, I. S.; Yang, S. I.; Ahn, K.-H. The considerable photostability improvement of photochromic terarylene by sulfone group. *Tetrahedron Lett.* **2009**, *50* (37), 5288-5290.

45. Tamayo, N.; Echavarren, A. M.; Paredes, M. C. Palladium-catalyzed coupling of 2bromonaphthoquinones with stannanes: a concise synthesis of antibiotics WS 5995 A and C and related compounds. *J. Org. Chem.* **1991**, *56* (23), 6488-6491.

46. Hassan, Z.; Ullah, I.; Ali, I.; Khera, R. A.; Knepper, I.; Ali, A.; Patonay, T.; Villinger, A.; Langer, P. Synthesis of tetraaryl-p-benzoquinones and 2,3-diaryl-1,4-naphthoquinones via Suzuki–Miyaura cross-coupling reactions. *Tetrahedron* **2013**, *69* (2), 460-469.

47. Ullah, I.; Khera, R. A.; Hussain, M.; Villinger, A.; Langer, P. Synthesis of tetraaryl-pbenzoquinones by Suzuki–Miyaura cross-coupling reactions of tetrabromo-p-benzoquinone. *Tetrahedron Lett.* **2009**, *50* (32), 4651-4653.

48. Ye, Q.; Zhang, Z.; Png, Z. M.; Neo, W. T.; Lin, T.; Zeng, H.; Xu, H.; Xu, J. Cyclization of Tetraaryl-Substituted Benzoquinones and Hydroquinones through the Scholl Reaction. *J. Org. Chem.* **2016**, *81* (19), 9219-9226.

49. Molina, M. T.; Navarro, C.; Moreno, A.; Csákÿ, A. G. Arylation of Benzo-Fused 1,4-Quinones by the Addition of Boronic Acids under Dicationic Pd(II)-Catalysis. *Org. Lett.* **2009**, *11* (21), 4938-4941.

50. Takahashi, I.; Takeyama, N.; Mori, H.; Tamamoto, M.; Nishimura, H.; Kitajima, H. Studies on the Selective Preparation of 2-Aryl and 2,3-Diaryl Naphthalene-1,4-diones by Means of Meerwein Arylation. *Chem. Express* **1992**, *7* (9), 709-712.

51. No reaction was observed when pinacol boronic acid esters were used regardless of the reaction conditions employed.

52. Hoshino, Y.; Miyaura, N.; Suzuki, A. Novel Synthesis of Isoflavones by the Palladium-Catalyzed Cross-Coupling Reaction of 3-Bromochromones with Arylboronic Acids or Its Esters. *Bull. Chem. Soc. Jpn.* **1988**, *61* (8), 3008-3010.

53. Brookins, R. N.; Schanze, K. S.; Reynolds, J. R. Base-Free Suzuki Polymerization for the Synthesis of Polyfluorenes Functionalized with Carboxylic Acids. *Macromolecules* **2007**, *40* (10), 3524-3526.

54. Rückle, T.; Biamonte, M.; Grippi-Vallotton, T.; Arkinstall, S.; Cambet, Y.; Camps, M.; Chabert, C.; Church, D. J.; Halazy, S.; Jiang, X.; Martinou, I.; Nichols, A.; Sauer, W.; Gotteland, J.-P. Design, Synthesis, and Biological Activity of Novel, Potent, and Selective (Benzoylaminomethyl)thiophene Sulfonamide Inhibitors of c-Jun-N-Terminal Kinase. *J. Med. Chem.* **2004**, 47 (27), 6921-6934.

55. Wuts, P. G. M.; Greene, T. W., *Greene's protective groups in organic synthesis*. Fifth edition ed.; John Wiley & Sons, Inc.: Hoboken, New Jersey, 2014.

56. Wright, S. W.; Hageman, D. L.; Wright, A. S.; McClure, L. D. Convenient preparations of t-butyl esters and ethers from t-butanol. *Tetrahedron Lett.* **1997**, *38* (42), 7345-7348.

57. Murphy, J. M.; Tzschucke, C. C.; Hartwig, J. F. One-Pot Synthesis of Arylboronic Acids and Aryl Trifluoroborates by Ir-Catalyzed Borylation of Arenes. *Org. Lett.* **2007**, *9* (5), 757-760.

58. Sasaki, I.; Taguchi, J.; Hiraki, S.; Ito, H.; Ishiyama, T. Catalyst-Controlled Regiodivergent C-H Borylation of Multifunctionalized Heteroarenes by Using Iridium Complexes. *Chem. Eur. J.* **2015**, *21* (25), 9236-9241.

59. Jia, Y.; Xu, Y.; Nie, R.; Chen, F.; Zhu, Z.; Wang, J.; Jing, H. Artificial photosynthesis of methanol from carbon dioxide and water via a Nile red-embedded TiO2 photocathode. *J. Mater. Chem. A* **2017**, *5* (11), 5495-5501.

60. O'Brien, C. J.; Kantchev, E. A. B.; Valente, C.; Hadei, N.; Chass, G. A.; Lough, A.; Hopkinson, A. C.; Organ, M. G. Easily Prepared Air- and Moisture-Stable Pd–NHC (NHC=N-Heterocyclic Carbene) Complexes: A Reliable, User-Friendly, Highly Active Palladium Precatalyst for the Suzuki–Miyaura Reaction. *Chem. Eur. J.* **2006**, *12* (18), 4743-4748.

61. Quibell, Jacob M.; Perry, G. J. P.; Cannas, D. M.; Larrosa, I. Transition-metal-free decarboxylative bromination of aromatic carboxylic acids. *Chem. Sci.* **2018**, *9* (15), 3860-3865.

62. Walton, I. M.; Cox, J. M.; Benson, C. A.; Patel, D. G.; Chen, Y.-S.; Benedict, J. B. The role of atropisomers on the photo-reactivity and fatigue of diarylethene-based metal-organic frameworks. *New J. Chem.* 2016, 40 (1), 101-106.
63. Walko, M.; Feringa, B. L. The isolation and photochemistry of individual atropisomers of photochromic diarylethenes. *Chem. Commun.* 2007, (17), 1745-1747.
64. Spek, A. Structure validation in chemical crystallography. *Acta Crystallogr. Sect. D. Biol. Crystallogr.* 2009, 65 (2), 148-155.

65. Kobatake, S.; Uchida, K.; Tsuchida, E.; Irie, M. Single-crystalline photochromism of diarylethenes: reactivity–structure relationship. *Chem. Commun.* **2002**, (23), 2804-2805.

66. Sahariah, B.; Sarma, B. K. Relative orientation of the carbonyl groups determines the nature of orbital interactions in carbonyl–carbonyl short contacts. *Chem. Sci.* **2019**, *10* (3), 909-917.

67. Yamamoto, S.; Matsuda, K.; Irie, M. Photochromism of Diarylethenes Linked by Hydrogen Bonds in the Single-Crystalline Phase. *Chem. Eur. J.* **2003**, *9* (20), 4878-4886.

68. Gottlieb, H. E.; Kotlyar, V.; Nudelman, A. NMR Chemical Shifts of Common Laboratory Solvents as Trace Impurities. *J. Org. Chem.* **1997**, *62* (21), 7512-7515.

69. Chen, T.; Ning, M.; Ye, Y.; Wang, K.; Leng, Y.; Shen, J. Design, synthesis and structure–activity relationship studies of GPR40 agonists containing amide linker. *Eur. J. Med. Chem.* **2018**, *152*, 175-194.

70. Kawamorita, S.; Ohmiya, H.; Sawamura, M. Ester-Directed Regioselective Borylation of Heteroarenes Catalyzed by a Silica-Supported Iridium Complex. *J. Org. Chem.* **2010**, *75* (11), 3855-3858.

71. Anuratha, M.; Jawahar, A.; Umadevi, M.; Sathe, V. G.; Vanelle, P.; Terme, T.; Meenakumari, V.; Milton Franklin Benial, A. SERS investigations of 2,3-dibromo-1,4-naphthoquinone on silver nanoparticles. *Spectrochim. Acta, Part A* **2013**, *105*, 218-222.

