

# Oxidation of Phosphines by Supercritical Nitrous Oxide


Scott Poh, Raquel Hernandez, Mayuko Inagaki, and Philip G. Jessop\*

Department of Chemistry, University of California, Davis, California 95616-5295

jessop@chem.ucdavis.edu

Received May 27, 1999

## ABSTRACT



Despite its reputation for lack of reactivity at moderate temperatures, nitrous oxide is capable of oxidizing at least one class of organic compounds, the phosphines, at temperatures at or below 100 °C. The use of supercritical N<sub>2</sub>O as both the solvent and the reactant simplifies the isolation of the products and allows one to avoid the use of flammable liquid solvents.

Nitrous oxide (N<sub>2</sub>O) is an under-utilized byproduct of chemical processes involving nitric acid. Scrubbing of N<sub>2</sub>O emissions from nylon manufacture has been adopted in recent years<sup>1</sup> in order to combat its contribution to the greenhouse effect.<sup>1–3</sup> Although much of the waste N<sub>2</sub>O is destroyed, one of the more attractive options for recycling N<sub>2</sub>O is its use as an oxidant for organic chemistry. From the point of view of “green” chemistry, nitrous oxide would be preferable to many other oxidants because the only byproduct is N<sub>2</sub>. The most dramatic result of efforts in academia and industry in this direction is the recently announced Monsanto process for the high-temperature heterogeneously catalyzed oxidation of benzene to phenol.<sup>4,5</sup> However, there is almost no information in the literature on low-temperature oxidation of organic functional groups by N<sub>2</sub>O.

Oxidations of most organic compounds by nitrous oxide are thermodynamically favorable but are reputed to be extremely slow except at high temperatures.<sup>6</sup> Buckley et al.<sup>7–9</sup> found that alkenes and alkynes were oxidized by N<sub>2</sub>O,

but the reactions required temperatures of at least 200 °C. Ohtani et al. found that 2-propanol and, to a lesser extent, other small secondary alcohols and primary amines in water were oxidized by N<sub>2</sub>O at room temperature in the presence of platinum “black”.<sup>10</sup> The low-temperature oxidation of inorganic compounds by N<sub>2</sub>O has been more fully explored. For example, catalyzed oxidations of CO<sup>11–13</sup> and NF<sub>3</sub>,<sup>14</sup> and especially the uncatalyzed oxidations of low-valent organo-metallic complexes,<sup>15–20</sup> have been well investigated.

Supercritical N<sub>2</sub>O (scN<sub>2</sub>O) or scN<sub>2</sub>O/CO<sub>2</sub> mixtures offer several advantages as media for the study of N<sub>2</sub>O chemistry. Nitrous oxide is supercritical at temperatures and pressures

(10) Ohtani, B.; Takamiya, S.; Hirai, Y.; Sudoh, M.; Nishimoto, S.; Kagiya, T. *J. Chem. Soc., Perkin Trans. 2* **1992**, 175–179.

(11) Li, C.-S.; Sun, K.-S.; Chang, C.-H. *J. Chem. Soc., Dalton Trans.* **1992**, 1025–1029.

(12) Lee, J.-D.; Fang, W.-P.; Li, C.-S.; Cheng, C.-H. *J. Chem. Soc., Dalton Trans.* **1991**, 1923–1927.

(13) Kazusaka, A.; Lunsford, J. H. *J. Catal.* **1976**, 45, 25–31.

(14) Christe, K. O. *J. Am. Chem. Soc.* **1995**, 117, 6136–6137.

(15) Banks, R. G. S.; Henderson, R. J.; Pratt, J. M. *J. Chem. Soc. A* **1968**, 2886–2889.

(16) Yamamoto, A.; Kitazume, S.; Pu, L. S.; Ikeda, S. *J. Am. Chem. Soc.* **1971**, 93, 371–380.

(17) Bottomley, F.; Lin, I. J. B.; Mukaida, M. *J. Am. Chem. Soc.* **1980**, 102, 5238–5242.

(18) Cummins, C. C.; Schrock, R. R.; Davis, W. M. *Inorg. Chem.* **1994**, 33, 1448–1457.

(19) Avens, L. R.; Barnhart, D. M.; Burns, C. J.; McKee, S. D.; Smith, W. H. *Inorg. Chem.* **1994**, 33, 4245–4254.

(20) The group of Hillhouse has been particularly active in this field. (a) Vaughan, G. A.; Rupert, P. B.; Hillhouse, G. L. *J. Am. Chem. Soc.* **1987**, 109, 5538–5539. (b) Vaughan, G. A.; Hillhouse, G. L.; Lum, R. T.; Buchwald, S. L.; Rheingold, A. L. *J. Am. Chem. Soc.* **1988**, 110, 7215–7217. (c) Koo, K.; Hillhouse, G. L. *Organometallics* **1998**, 17, 2924–2925 and references therein.

(1) The Draft 1998 Inventory of U.S. Greenhouse Gas Emissions and Sinks (1990–1996), U.S. Environmental Protection Agency, 1998.

(2) Troglor, W. C. *J. Chem. Educ.* **1995**, 72, 973–976.

(3) Dickinson, R. E.; Cicerone, R. J. *Nature* **1986**, 319, 109–115.

(4) “Monsanto announces new high-yield process to manufacture phenol”, Monsanto, press release, 1996.

(5) Panov, G. I.; Sheveleva, G. A.; Kharitonov, A. S.; Romannikov, V. N.; Vostrikova, L. A. *Appl. Catal. A-General* **1992**, 82, 31–36.

(6) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 4th ed.; John Wiley: New York, 1980.


(7) Bridson-Jones, F. S.; Buckley, G. D. *J. Chem. Soc.* **1951**, 3009–3016.

(8) Bridson-Jones, F. S.; Buckley, G. D.; Cross, L. H.; Driver, A. P. *J. Chem. Soc.* **1951**, 2999–3008.

(9) Buckley, G. D.; Levy, W. J. *J. Chem. Soc.* **1951**, 2999–3008.

above its critical point of 36.4 °C and 72.5 bar.<sup>21</sup> Depending on the exact conditions, the density is high enough for the scN<sub>2</sub>O to act as a solvent for many nonpolar organic compounds, in the same manner as supercritical carbon dioxide (scCO<sub>2</sub>). *The reader should peruse the safety warnings in the footnotes.*<sup>22</sup> In the past, the few low-temperature oxidations with N<sub>2</sub>O that have been reported were performed in either organic solvents or water. Using the supercritical fluid as the solvent allows one to avoid the use of a flammable liquid solvent and also to dissolve substrates too hydrophobic to dissolve in water. A second advantage is the facile separation of the product from the solvent.<sup>23</sup> The use of scN<sub>2</sub>O as both solvent and oxidant for organic functional group transformations has not been reported previously.

During tests of the effect of scN<sub>2</sub>O on various compounds, we found that attempting to dissolve PPh<sub>3</sub> in scN<sub>2</sub>O at 100 °C and 100–140 bar resulted in its rapid oxidation to phosphine oxide (Table 1 and eq 1).<sup>24</sup> The N<sub>2</sub> byproduct is assumed but was not experimentally confirmed.



The reaction can also be achieved at 65 °C, but it is somewhat slower at this temperature. The pressure of the N<sub>2</sub>O must be supercritical or it will be unable to dissolve the phosphine. For example, the oxidation of PPh<sub>3</sub> at 65 °C proceeds readily at 100 to 140 bar but the starting material was recovered entirely unreacted after an experiment at 10 bar. Note that PPh<sub>3</sub> at this temperature is a solid. Methyl-diphenylphosphine, tricyclohexylphosphine, and triethylphosphine are oxidized more rapidly by scN<sub>2</sub>O than is PPh<sub>3</sub>. Triethylphosphine (470 μmol) was 43% oxidized after 90 min in a mixture of N<sub>2</sub>O (6 bar, ca. 40 mmol) and CO<sub>2</sub> (total pressure 100 bar) at 50 °C. The same experiment but with only CO<sub>2</sub> and no N<sub>2</sub>O gave 0% oxidation.

This oxidation of PR<sub>3</sub> requires no added catalyst, although catalysis by the vessel walls cannot be ruled out. In contrast to the present results, previously reported examples of the

oxidation of PPh<sub>3</sub> by N<sub>2</sub>O have involved metal complexes as catalysts or promoters; the complexes were CoH(N<sub>2</sub>)-(PPh<sub>3</sub>)<sub>3</sub> (6 mol OPPh<sub>3</sub> per mol catalyst, 62% isolated yield, –5 to 20 °C)<sup>16,25</sup> and [MoO(CN)<sub>5</sub>]<sup>3-</sup> (0.5 mol OPPh<sub>3</sub> per mol catalyst, room temperature, 48 h).<sup>26</sup> Presumably the lack of a requirement for a catalyst in the present system is due to the higher temperature and the higher concentration of N<sub>2</sub>O in the reaction phase.

Tests of the solubility of PPh<sub>3</sub> in scN<sub>2</sub>O were not possible because of the rapid reaction. Instead, observations were made of the solubility of PPh<sub>3</sub> in scCO<sub>2</sub>, a supercritical fluid which is quite similar to scN<sub>2</sub>O in all of its physical properties. PPh<sub>3</sub> (up to 65 mg) was completely dissolved in the scCO<sub>2</sub> at 150 bar and 65 or 100 °C. Literature measurements of the solubility of PPh<sub>3</sub> in scCO<sub>2</sub> have only been reported for 47 °C and below, but the mole fraction solubility at 150 bar was found to be roughly temperature independent.<sup>27</sup> The mole fraction used in our experiments is an order of magnitude lower than the solubility limit at this pressure.

It is unlikely that the oxygen source was anything other than N<sub>2</sub>O. The purity of the N<sub>2</sub>O was 99.998%, with less than 2 ppm of O<sub>2</sub> and less than 20 ppm of other impurities. Two different batches of N<sub>2</sub>O were tested. Inadvertent admission of air to the vessel could not have caused the oxidation, because there is little reaction between PPh<sub>3</sub> and low concentrations of O<sub>2</sub> under these conditions. A reaction of PPh<sub>3</sub> in scCO<sub>2</sub> with no N<sub>2</sub>O present gave 0% oxidation, as did an experiment in scCO<sub>2</sub> with 1 bar of added air. The oxidation of PEt<sub>3</sub> in a N<sub>2</sub>O/CO<sub>2</sub> mixture proceeded at a PEt<sub>3</sub>:N<sub>2</sub>O mole ratio of 1:85, further evidence that the oxidation could not have been caused by the presence of trace

(21) Couch, E. J.; Kobe, K. A.; Hirth, L. J. *J. Chem. Eng. Data* **1961**, *6*, 229–237.

(22) **SAFETY WARNINGS:** (a) Nitrous oxide is a thermodynamically powerful oxidant. Never mix high concentrations of organic compounds with scN<sub>2</sub>O. Explosions have occurred with a 9 vol % solution of ethanol in scN<sub>2</sub>O and with a mixture of 1 g of ground coffee in 2.5 mL of scN<sub>2</sub>O.<sup>31–33</sup> To minimize the risk, we choose to keep the combustible substrate to microscale quantities and *very low concentrations*. For example, the experiments with triphenylphosphine were typically performed at 0.00018 g substrate per mL of scN<sub>2</sub>O. In addition to using microscale quantities of substrate, we employ a burst disk, blast shield, and eye protection in all experiments. Diluting scN<sub>2</sub>O with scCO<sub>2</sub> may further enhance the safety. Combustible cosolvents should not be used with scN<sub>2</sub>O under any circumstances. (b) Triethylphosphine ignites on exposure to air<sup>34</sup> or pure N<sub>2</sub>O.

(23) Jessop, P. G.; Leitner, W., Eds. *Chemical Synthesis using Supercritical Fluids*; VCH/Wiley: Weinheim, 1999.

(24) **General Procedure for the Oxidation of a Phosphine by scN<sub>2</sub>O.** A stir bar and 21 mg of PPh<sub>3</sub> were placed in a 160 mL stainless steel vessel while in a glovebox under argon. The vessel cap contains a burst disk, a pressure transducer, a thermocouple, and a valve for N<sub>2</sub>O. The vessel was closed, removed from the glovebox, heated to 100 °C, and then pressurized with 140 bar of scN<sub>2</sub>O. The temperature was monitored by a thermocouple inside a thermowell. After 3 h, the vessel was cooled to room temperature and then placed in a bath of acetone/dry ice until the internal pressure dropped to zero due to condensation of the N<sub>2</sub>O. At this point, the vent valve was opened and the vessel slowly warmed to room temperature, the N<sub>2</sub>O venting as it boiled. The vessel was then opened, and the product was recovered from the bottom of the vessel by dissolution in CDCl<sub>3</sub> and analyzed by <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy without further workup. The product was pure OPPh<sub>3</sub> (100% conversion, 94% isolated yield).

(25) Pu, L. S.; Yamamoto, A.; Ikeda, S. *Chem. Commun.* **1969**, 189–90.

(26) Arzoumanian, H.; Nuel, D.; Sanchez, J. *J. Mol. Catal.* **1991**, *65*, L9–L11.

(27) Schmitt, W. J.; Reid, R. C. *Chem. Eng. Commun.* **1988**, *64*, 155–176.

**Table 1.** Oxidation of Phosphines to Phosphine Oxides by N<sub>2</sub>O

| substrate          | T, °C | SCF                          | P <sub>total</sub> , bar | time, h | % convn |
|--------------------|-------|------------------------------|--------------------------|---------|---------|
| PPh <sub>3</sub>   | 65    | N <sub>2</sub> O             | 10                       | 2       | 0       |
| PPh <sub>3</sub>   | 65    | N <sub>2</sub> O             | 100                      | 2       | 43      |
| PPh <sub>3</sub>   | 65    | N <sub>2</sub> O             | 140                      | 2       | 34      |
| PPh <sub>3</sub>   | 100   | N <sub>2</sub> O             | 130                      | 0.5     | 66      |
| PPh <sub>3</sub>   | 100   | N <sub>2</sub> O             | 140                      | 3       | 100     |
| PPh <sub>3</sub>   | 100   | CO <sub>2</sub> <sup>a</sup> | 100                      | 0.5     | 0       |
| PMePh <sub>2</sub> | 64    | N <sub>2</sub> O             | 140                      | 2       | 71      |
| PMePh <sub>2</sub> | 65    | N <sub>2</sub> O             | 140                      | 4       | 100     |
| PCy <sub>3</sub>   | 70    | N <sub>2</sub> O             | 140                      | 4       | 100     |
| PEt <sub>3</sub>   | 50    | CO <sub>2</sub> <sup>b</sup> | 100                      | 1.5     | 43      |
| PEt <sub>3</sub>   | 50    | CO <sub>2</sub>              | 100                      | 1.5     | 0       |

<sup>a</sup> 1 bar of air added. <sup>b</sup> 6 bar of N<sub>2</sub>O added.

impurities in the  $N_2O$ . A possibility that cannot be ruled out is the conversion of the  $N_2O$  to other nitrogen oxides, either thermally or as a wall-catalyzed reaction, followed by reaction of those nitrogen oxides with  $PPh_3$ . The oxidation of phosphines by  $NO$  and  $N_2O_4$  has been reported.<sup>28,29</sup>

Many other classes of organic compounds are *not* oxidized by dissolution in or exposure to  $scN_2O$ . For example, the following compounds did not react when kept in  $scN_2O$  solution overnight at 90–130 °C: 2-butanol, cyclohexane, cyclododecene, cyclohexanone, triethylamine, and diethyl sulfide. Even aliphatic and aromatic aldehydes were recovered unreacted if air was rigorously excluded at all stages. Thus, aldehydes and phosphines are opposite in their reactivity toward  $O_2$  and  $N_2O$ . Aldehydes are easily oxidized by air at room temperature but are resistant to oxidation by  $N_2O$  even at 130 °C. Neither triphenylphosphine nor methyldiphenylphosphine in acetonitrile solution is oxidized by exposure to air over 17 h at room temperature, but both are readily oxidized by  $N_2O$  at 65 °C. It is not surprising that ease of oxidation by  $O_2$  is not a reliable indicator of ease of

(28) Longhi, R.; Ragsdale, R. O.; Drago, R. S. *Inorg. Chem.* **1962**, *1*, 768–770.

(29) Addison, C. C.; Sheldon, J. C. *J. Chem. Soc.* **1956**, 2705–2708.

oxidation by  $scN_2O$ ; very different mechanisms must be involved. For oxidation of aldehydes and other substrates by  $scN_2O$ , catalysis will be required.<sup>30</sup>

In conclusion, nitrous oxide is capable of oxidizing at least one class of organic compounds, the phosphines, at temperatures well below 200 °C. The use of supercritical  $N_2O$  as the solvent and the reactant simplifies the isolation of the products and allows one to avoid the use of flammable liquid solvents. A kinetic investigation of the reaction and tests of the oxidation of other functional groups by  $scN_2O$  or  $scN_2O/CO_2$  mixtures, with and without catalysts, are in progress.

**Acknowledgment** is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for support of this research.

OL9906918

(30) We have found that Pt/C is capable of catalyzing the oxidation of secondary alcohols by and in  $scN_2O$ . This work will be published separately.

(31) Raynie, D. E. *Anal. Chem.* **1993**, *65*, 3127–3128.

(32) Hansen, B. N.; Hybertson, B. M.; Barkley, R. M.; Sievers, R. E. *Chem. Mater.* **1992**, *4*, 749–752.

(33) Sievers, R. E.; Hansen, B. *Chem. Eng. News* **1991**, *69*, (29), 2.

(34) Lenga, R. E., Ed. *The Sigma-Aldrich Library of Chemical Safety Data*, 2nd ed.; Sigma-Aldrich Corp.: Milwaukee, WI, 1988.

