


This article was downloaded by: [Moskow State Univ Bibliote]

On: 06 February 2014, At: 04:23

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Synthesis of Mixed Alkylphosphites and Alkylphosphates

Gheorghe Iliu^a, Adriana Popa^a, Smaranda Iliescu^a, Alina Bora^a, Gheorghe Dehelean^a & Aurelia Pascariu^a

^a Institute of Chemistry, Romanian Academy, Timisoara, Romania

Published online: 27 Oct 2010.

To cite this article: Gheorghe Iliu, Adriana Popa, Smaranda Iliescu, Alina Bora, Gheorghe Dehelean & Aurelia Pascariu (2003) Synthesis of Mixed Alkylphosphites and Alkylphosphates, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 178:7, 1513-1519, DOI: [10.1080/10426500307865](https://doi.org/10.1080/10426500307865)

To link to this article: <http://dx.doi.org/10.1080/10426500307865>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS OF MIXED ALKYLPHOSPHITES AND ALKYLPHOSPHATES

Gheorghe Ilia, Adriana Popa, Smaranda Iliescu, Alina Bora,
Gheorghe Dehelean, and Aurelia Pascariu
Romanian Academy, Institute of Chemistry, Timisoara,
Romania

(Received December 17, 2002; accepted January 28, 2003)

Some mixed phosphites having two different alkyl chain were obtained as forerunners for mixed phosphates. Mixed dialkyl phosphites were obtained in good yields (40–80%) by phase transfer catalysis in liquid-liquid system, starting from different dialkyl phosphites and aliphatic alcohols. The reaction conditions were optimized in order to obtain good yields in phosphites and phosphates respectively. Compounds were analyzed by IR, P^{31} -NMR.

Keywords: Mixed phosphates; mixed phosphites; phase transfer catalysis

The reaction of phosphorus chlorine derivatives with alcohols is well known and is used for obtaining a wide range of phosphorus esters,¹ but the yields were low. For the synthesis of mixed phosphates Steinberg,² then Kenner and Williams³ proposed the system dialkylphosphite/carbontetrachloride/triethylamine and alcohol. This system has a few major disadvantages: it needs an equivalent mol of amine; reaction products are impure and contain ammonium chloride; reaction times are very long (more than 12 h); only moderate yields.

Zwierzak⁴ showed that alcohols can react with dialkylphosphites in a biphasic system, carbon tetrachloride-aqueous sodium hydroxide (Scheme 1). The studies were limited to a few aliphatic alcohols. We found that this method can be applied to series of aliphatic alcohols⁵ and aliphatic phosphite. This article presents the obtained results in optimal conditions.

Address correspondence to Gheorghe Ilia, Romanian Academy, Institute of Chemistry, 24 M. Viteazu Bvd. 1900 Timisoara, Romania. E-mail: ilia@acad-tim.utt.ro





FIGURE 1 The influence of stirring speed on the yield of phosphates.

Preliminary studies allowed a few observations: a) In order to obtain good yields it needs 50% aqueous sodium hydroxide; b) it needs an excess of dialkylphosphite due to saponification reactions; c) the order of reagent addition is important; and d) the solvent is important.

Based on these observations it can be concluded that, in the phosphorylation reaction of alcohols by the phase transfer catalysis method, the phosphite is oxidized to dialkylchlorophosphate which reacts with alcohols by nucleophilic substitution, and mixed phosphates are obtained (Scheme 3).



SCHEME 3




FIGURE 2 The influence of the reaction time on the yields of phosphates.


TABLE I The Influence of the Catalyst on the Yield

Synthesis	Catalyst	Yields (%)
0	1	2
TEP	TEAC	50,1
DEBP	TEAC	58,2
DBEP	TEAC	54,1
TBP	TEAC	56
TEP	TEAB	55,2
DEBP	TEAB	63,3
DBEP	TEAB	59,8
TBP	TEAB	60,2
TEP	TBAC	63,3
DEBP	TBAC	71,4
DBEP	TBAC	68,7
TBP	TBAC	69,8
TEP	TBAB	68,9
DEBP	TBAB	75
DBEP	TBAB	72,3
TBP	TBAB	71

TEAC(B) = tetraethylammonium chloride (bromide).

TBAC(B) = tetrabutylammonium chloride (bromide).

Side reactions, with formation of dihalocarbene, appear when 50% aqueous sodium hydroxide is used (Scheme 4). Dihalocarbene can react also with alcohols.



SCHEME 4

In accordance with the general procedure, the reactions were performed with different alcohols and diethylphosphite, dipropylphosphite and dibutylphosphite by the phase transfer catalysis method in liquid-liquid system (Table II, Table III, and Table IV).

In all syntheses, the reaction temperature was kept at 15°C and the reaction time was 3 h. Decreasing the reaction temperature below 15°C did increase the reaction time and side reactions and the yields did decrease. Mixed phosphates obtained from C1–C4 were chromatographically pure. The obtained products were analyzed also by IR. In the IR spectra, the disappearance of the band at 2430–2440 cm⁻¹(P–H) was used for the reaction control.

Also mixed phosphites were obtained, by direct hydrolysis of dialkyl phosphites with 20% aqueous tetraethylammonium hydroxide followed

TABLE II Experimental Results in the Synthesis of Some Phosphates Starting from Diethylphosphite (Catalyst-Tetrabutylammonium Bromide 0.5 mmol %)

Synthesis	R ₂ OH	Yield (%)	Yield from lit (%)	b.p. Lit., °C/mm Hg	b.p. °C/mm Hg	³¹ P NMR (ppm)
0	1	5	6	7	8	9
DEMF	Methanol	72	—	100.2/20*	109/30	1.04
TEP	Ethanol	68.9	—	116/30*	123/40	1.00
DEPP	<i>n</i> -Propanol	70.5	67**	50/0.2**	116/35	0.97
DEBP	<i>n</i> -Butanol	75	71**	123/15**	138/25	1.10
DEPnP	<i>n</i> -Pentanol	69	—	—	150/25	1.25
DEHP	<i>n</i> -Hexanol	41	—	—	162/25	1.03
DEHPP	<i>n</i> -Heptanol	47	—	—	173/26	1.20
DEOP	<i>n</i> -Octanol	45	—	—	179/25	1.13

* = [4]; ** = [8].

TABLE III Experimental Results in the Synthesis of Some Phosphates Starting from Dipropylphosphite (Catalyst-Tetrabutylammonium Bromide 0.5 mmol %)

Synthesis	R ₂ OH	Yield (%)	b.p. °C/mm Hg	³¹ P NMR (ppm)
DPMP	Methanol	63	113/15	0.72
DPEP	Ethanol	71	126/14	1.07
TPP	<i>n</i> -Propanol	70	130/15	0.76
DPBP	<i>n</i> -Butanol	73.5	138/14	1.00
DPPnP	<i>n</i> -Pentanol	73	147/12	0.97
DPHP	<i>n</i> -Hexanol	54.5	153/12	0.98
DPHpP	<i>n</i> -Heptanol	55	169/11	1.02
DPOP	<i>n</i> -Octanol	52	179/10	1.09

TABLE IV Experimental Results in the Synthesis of Some Phosphates Starting from Dibutylphosphite (Catalyst-Tetrabutylammonium Bromide 0.5 mmol %)

Synthesis	R ₂ OH	Yield (%)	Yield lit (%)	b.p. Literat. °C/mm Hg	b.p. °C/mm Hg	³¹ P NMR (ppm)
DBMP	Methanol	65	—	—	95/10	0.65
DBEP	Ethanol	72	71*	60–1/0,1*	103/15	1.00
DBPP	<i>n</i> -Propanol	68	—	—	121/20	1.09
TBP	<i>n</i> -Butanol	71.6	—	180-3/25**	190/30	0.97
DBPnP	<i>n</i> -Pentanol	78	—	—	186/15	1.03
DBHP	<i>n</i> -Hexanol	69	—	—	175/10	1.10
DBHpP	<i>n</i> -Heptanol	62	—	—	182/10	1.09
DBOP	<i>n</i> -Octanol	60	—	—	198/10	1.12

* = [4]; ** = [8].

TABLE V Preparation of Tetraethylammonium Hydrogenphosphites (2), Mixed Phosphites (3), and Phosphates

Product	R	R1	R2	³¹ P NMR (ppm)	Yield (%)
2a	C ₂ H ₅	—	—	—	75
2b	<i>n</i> -C ₄ H ₉	—	—	—	80
3a	C ₂ H ₅	CH ₃	—	-4.3	61
3b	C ₂ H ₅	<i>n</i> -C ₄ H ₉	—	-6.6	73
3b	<i>n</i> -C ₄ H ₉	C ₂ H ₅	—	-6.6	72
EMPP	C ₂ H ₅	CH ₃	<i>n</i> -C ₃ H ₇	0.89	67
BEMP	C ₂ H ₅	<i>n</i> -C ₄ H ₉	<i>n</i> -C ₃ H ₇	1.0	76
BEMP	<i>n</i> -C ₄ H ₉	C ₂ H ₅	<i>n</i> -C ₃ H ₇	1.0	75

by reaction with alkyl iodides in acetonitrile. They are the forerunners for mixed phosphates having three different alkyl chains (Table V).

The yields were good in all syntheses. Mixed phosphites were used for obtaining phosphates with three different alkyl radicals.

EXPERIMENTAL

General Procedure for the Synthesis of Mixed Phosphates

To a stirred solution of 50% aqueous sodium hydroxide (60 ml), alkyl alcohol (0.2 mmol), carbon tetrachloride (50 ml), methylene chloride (50 ml) and a phase transfer catalyst, was added dropwise a solution of a phosphite (0.25 moles) in carbon tetrachloride (60 ml). The temperature was kept at 15°C. Then the mixture was kept under stirring 3 h at 15°C. The mixture is diluted with methylene chloride (50 ml), filtered, separated, and the organic layer washed with 2% aqueous hydrogen chloride (25 ml), and water (2 × 25 ml), and dried with anhydrous sodium sulfate and the solvents distilled under reduced pressure.

General Procedure for the Synthesis of Mixed Phosphites

A 20% aqueous solution of tetrabutylammonium hydroxide (0.2 mmol) is added to stirred phosphite (0.2 mmol) at 5–10°C and the mixture is kept at this temperature for 2 h. The salt is extracted with methylene chloride, the solvent is distilled and the crude intermediate mixed with 50 ml acetonitrile and the appropriate alkyl iodide and heated for 6 h at 60°C. The solvent is distilled and the product is purified by distillation under reduced pressure.

CONCLUSIONS

Mixed phosphites were obtained with good yields. Mixed phosphates starting from diethyl-, dipropyl-, dibutyl phosphite and mixed phosphites, respectively, were synthesized using a liquid-liquid system. The best results were obtained when 50% NaOH_{aq} was used, the reaction temperature was 15°C, and the reaction time 3 h and the molar ratio phosphite : alcohol = 1.25:1. P³¹-NMR spectra, performed with a Bruker Avance DRX 400 spectrometer, showed chemical shift values of $\delta = 0.6\text{--}1.25$ ppm (ref. H₃PO₄85%). Mixed phosphates obtained from C₁–C₄ were chromatographically pure.

REFERENCES

- [1] W. E. Keller, *Phase Transfer Reactions* (Georg Thieme Verlag, Stuttgart, 1990), Fluka Compendium, vol. 3.
- [2] G. M. Steinberg, *J. Org. Chem.*, **15**, 637 (1950).
- [3] G. W. Kenner and N. R. Williams, *J. Chem. Soc.*, 522 (1955).
- [4] A. Zwierzak, *Synthesis*, 305 (1976).
- [5] G. Iliu, S. Iliescu, G. Dehelean, A. Popa, L. Pacureanu, L. Macarie, and A. Pascariu, *Phosphorus, Sulfur, and Silicon*, **177**, 2049 (2002).
- [6] G. Iliu, L. Kurunczi, G. Dehelean, S. Iliescu, and L. Macarie, *Phosphorus, Sulfur, and Silicon*, **147**, 191 (1999).
- [7] C. M. Starks and C. Liotta, *Phase Transfer Catalysis* (Academic Press, New York, 1978).
- [8] G. M. Kosolapoff and L. Maier, eds., *Organic Phosphorus Compounds* (Wiley, New York, 1973), vol 6, pp. 409.