



CO₂ TransformationCO₂ Conversion into Esters by Fluoride-Mediated Carboxylation of Organosilanes and Halide DerivativesXavier Frogneux⁺, Niklas von Wolff⁺, Pierre Thuéry, Guillaume Lefèvre,* and Thibault Cantat^{*[a]}

Abstract: A one-step conversion of CO₂ into heteroaromatic esters is presented under metal-free conditions. Using fluoride anions as promoters for the C–Si bond activation, pyridyl, furanyl, and thienyl organosilanes are successfully carboxylated with CO₂ in the presence of an electrophile. The mechanism of this unprecedented reaction has been elucidated based on experimental and computational results, which show a unique catalytic influence of CO₂ in the C–Si bond activation of pyridylsilanes. The methodology is applied to 18 different esters, and it has enabled the incorporation of CO₂ into a polyester material for the first time.

The development of novel methodologies enabling the use of CO₂ is of high interest to benefit from this renewable and environmentally friendly carbon source in the production of chemicals. Recent efforts have targeted the formation of value-added products from CO₂.^[1] For example, the reduction of CO₂ provides an entry to formate and methoxide derivatives, and the reductive functionalization of CO₂ with amines has led to novel transformations over the last ten years.^[2] Because C–C bonds are dominant in organic chemistry, the formation of C–C bonds with CO₂ will be of prominent utility. Transition-metal chemistry has played a substantial role in the development of such transformations, allowing for the coupling of CO₂ with alkynes, dienes, alkenes, and aryl halides using stoichiometric amounts of Ni or Fe, or on the other hand Pd, Cu, Ni, Fe, Rh, or Au-based catalysts.^[3] The formation of carboxylic acids with CO₂ has also been developed with various nucleophiles, such as organoboranes,^[4] organostannanes,^[5] and olefins^[6] in transition-metal-catalyzed systems. In this context, organosilanes are attractive nucleophiles, because they are less toxic and easier to prepare and handle than tin or boron reagents. Yet, organosilanes display a nucleophilicity three to six orders of magnitude lower than organoboranes and organostannanes,^[7] and



Scheme 1. Fluoride-mediated carboxylation of C–Si bonds with CO₂.

their use in CO₂-based carboxylation reactions remains limited to a handful of examples (Scheme 1).

Because the C–Si bond features a low polarity, its activation with a fluoride source is necessary to generate a carbanion synthon, which can undergo a carboxylation with CO₂. Using CsF, the group of Sato recently reported the synthesis of α -amino acid derivatives (Scheme 1a),^[8] α -hydroxy acid derivatives (Scheme 1b),^[9] and the carboxylation of benzylsilane derivatives featuring a directing group (Scheme 1c).^[10] The group of Kondo developed the carboxylation of alkyne-silanes (Scheme 1d).^[11] Early on, Effenberger and Spiegler reported the carboxylation of electron deficient arylsilanes, bearing *ortho*-NO₂ and *ortho*-Cl groups, using KF or CsF under high pressures of CO₂ (50 atm), presumably to avoid the formation of the free aryl anion (Scheme 1e).^[12] These few examples highlight the small scope of reactive organosilanes that is limited to the generation of stabilized carbanions having a corresponding pK_a lower than 35. Additionally, the esterification step must be carried out in a second step to prevent a direct quenching of the carbon nucleophile by the electrophile (RCHN₂ or RX) in the formation of the ester product. Aromatic and heteroaromatic esters being key intermediates in the synthesis of polymers, pharmaceutical intermediates or final products,^[13] we sought to enlarge the scope of ester derivatives available from CO₂

[a] Dr. X. Frogneux,⁺ N. v. Wolff,⁺ Dr. P. Thuéry, Dr. G. Lefèvre, Dr. T. Cantat
NIMBE, CEA, CNRS, Université Paris-Saclay
Gif-sur-Yvette (France)
Fax: (+33) 1-6908-6640
E-mail: thibault.cantat@cea.fr


[⁺] Both authors contributed equally to this work.

Supporting information and ORCID(s) from the author(s) for this article are available on the WWW under <http://dx.doi.org/10.1002/chem.201505092>.

and organosilanes, under metal-free conditions. We show herein that a wide range of heteroaromatic silanes can be used in the direct synthesis of esters from CO₂, organohalides, and a fluoride source; the methodology has been transposed to the formation of a polyester material from CO₂. Mechanistic studies show that this reactivity can be enabled by a unique catalytic influence of CO₂ in the activation of the C–Si bond.

In the absence of any transition metal, the controlled activation of commercially available phenyltrimethylsilane (**1a**) and 2-(trimethylsilyl)pyridine (**2a**) with fluoride ions is challenging, because the corresponding C₆H₅[−] and C₅H₄N[−] anions are unstabilized carbanions with pK_a of 45 and 44, respectively. Indeed, phenyltrimethylsilane (**1a**) exhibited no reactivity when subjected to an atmosphere of CO₂, in the presence of CH₃I and a variety of fluoride sources, such as CsF, TASF (tris(dimethylamino)sulfonium difluorotrimethylsilicate), and TBAT (tetrabutylammonium difluorotriphenylsilicate). In contrast, traces of the methyl picolinate (**4aa**) were detected when **2a** was reacted with CO₂, CH₃I, and CsF, TBAF·3H₂O, or TMAF as a fluoride source (Table 1, entries 1–3). Whereas Brønsted bases, such as Cs₂CO₃ or *t*BuOK, were unable to promote the carboxylation of the substrate (entries 4 and 5); other anhydrous fluoride salts, such as TASF and TBAT, allowed for the formation of the methyl picolinate (**4aa**) in moderate to excellent yields (46% and 93%, respectively, entries 6 and 7). It is worth noting that a procedure to recycle the Ph₃SiF byproduct is possible by the addition of TBAF to reform TBAT.^[14] Carrying the reaction in THF or CH₂Cl₂ allowed for a nearly quantitative conversion into the ester.

The formation of **4aa** represents the first example of the direct carboxylation of a pyridylsilane reagent. Importantly, the



Scheme 2. Synthesis of picolinic esters by fluoride-mediated carboxylation of 2-(trimethylsilyl)pyridine (**2a**) with various electrophiles. NMR yield given; [a] average isolated yield over two runs on 1 mmol scale; [b] 3 h (0.5 equiv).

organosilane is compatible with both CO₂ and the haloalkane, and the reaction proceeds in one pot. In fact, changing the nature of the electrophile enables the formation of a large diversity of picolinic esters (Scheme 2). Primary and secondary alkyl and allyl iodides provided very good yields to the desired esters (**4ab**, 90%; **4ac**, 85%; **4ae**, 93%; **4af**, 99%) after 40 h at room temperature (RT) or 18 h at 40 °C (Scheme 2). Iodobenzene proved to be unreactive in this reaction, indicating that the formation of the O–C bond occurs through a nucleophilic substitution. Importantly, the alkylation of the pyridine ring with the organic halide was not observed, and the carbon nucleophile, released by C–Si bond activation, selectively reacted with CO₂. The utilization of less electrophilic bromoalkanes and chloroalkanes required longer reaction times at RT (72 h) or mild heating (70 °C), as exemplified by the formation of **4ae**, **4af**, and **4ag** in >93% yield from various organic halides (Scheme 2). Interestingly, using a difunctional electrophilic species, such as 1,4-diiodobutane, diester **4ah** formed in 85% yield and crystals suitable for X-ray diffraction analysis could be isolated (see the Supporting Information and Figure S6).

The substituents borne by the silicon center have a notable influence on the reactivity of the pyridylsilane reagent and different kinetic rates were measured when one methyl group in **2a** was replaced with a vinyl (**2a₂**), allyl (**2a₃**) or phenyl (**2a₄**) group. Although the presence of an allyl group slows down the activation of the C–Si bond, introducing a phenyl or vinyl group at the silicon center enhances the kinetics of the carboxylation reaction (see the Supporting Information).



Taken together, these data suggest that the formation of picolinic esters proceeds by a fluoride-mediated activation of the C–Si bond, which releases a carbanion able to undergo carboxylation in the presence of CO₂ prior to the esterification step with the electrophile. Noticeably, no reaction occurred for arylsilanes that do not feature a nitrogen atom on the *ortho* position of the silyl group, and both phenyldimethylvinylsilane

Table 1. Screening of the reaction conditions for the direct esterification of **2a** using CO₂ and CH₃I.

Entry ^[a]	Additive ^[b]	Solvent	Yield [%]
1	CsF	THF	traces
2	TBAF·3H ₂ O	THF	7
3	TMAF	THF	traces
4	Cs ₂ CO ₃	THF	0
5	<i>t</i> BuOK	THF	0
6	TASF	THF	46
7	TBAT	THF	93 (82) ^[c]
8	TBAT	CH ₃ CN	12
9	TBAT	CH ₂ Cl ₂	95
10	TBAT	toluene	2
11	–	THF	0

[a] Reaction conditions: NMR tube, **2a** (0.1 mmol), additive (0.1 mmol), **3a** (0.1 mmol), solvent (0.3 mL), CO₂ (1 bar). NMR yield determined with mesitylene as internal standard; [b] TBAF = [nBu₄N][F], TMAF = [Me₆N][F]; [c] Isolated yield.


(**1a₂**) and 3-(trimethylsilyl)pyridine (**21**) are unreactive under the applied conditions [Eqs. (1) and (2)]. These results are puzzling given that the three anions $C_6H_5^-$, $o-C_5H_4N^-$, and $m-C_5H_4N^-$ have similar pK_a values and hence similar stability. The mechanism of this novel transformation was thus investigated from stepwise reactions and DFT calculations so as to rationalize these findings.




Experimentally, under an argon atmosphere, silane **2a₂** was stable in the presence of TBAT and only a small amount of pyridine (6%) was detected by ¹H NMR after 22 h at 70 °C. However, in the presence of CO₂, the formation of a picolinate species 2-PyCO₂⁻ was observed (68% yield) at room temperature (see the Supporting Information). After addition of CH₃I, the ester **4aa** was obtained as the sole pyridine-based product, along with NBu₄I. These results suggest that the pyridine anion C₅H₄N⁻ does not form under the applied conditions and that CO₂ is required to facilitate the C–Si bond activation.

The different reactivities of **1a₂**, **2a₂**, and **21** were further investigated using DFT calculations (Scheme 3). For both the phenyl (**1a₂**) and pyridine (**2a₂**) derivatives, the formation of the carboxylate anion Ar–CO₂⁻ is favored thermodynamically ($\Delta G < -19.7$ kcal.mol⁻¹) and the esterification step is also mildly exergonic. Using Me₃SiF₂⁻ as fluoride source model, the

formation of the pentavalent anionic species PhSiMe₂(CHCH₂)F⁻ (**A_{Ph}**) from **1a₂** is endothermic yet accessible at RT ($\Delta G = 21.8$ kcal.mol⁻¹). Nevertheless, the direct carboxylation of **A_{Ph}** to Ph–CO₂⁻ (**TS₂**) or the generation of the free C₆H₅⁻ anion (**TS₃**) is incompatible with the reaction conditions, with high-energy transition states (TSs) lying at 38.1 and 38.9 kcal.mol⁻¹, respectively. Replacing the phenyl group with a pyridine core only stabilizes these two TSs by 2.9 kcal.mol⁻¹, revealing that the C–Si bond activation follows a different pathway for **2a₂**. The formation of anion **A_{2-Py}** is accompanied with an increase of negative charge on the nitrogen atom of the pyridine core, from 0.48 in **2a₂** to 0.52 in **A_{2-Py}**, thereby enabling fixation of CO₂ to the nitrogen atom. The formation of the resulting CO₂ adduct **B** involves an energy barrier of 7.5 kcal.mol⁻¹ (TS₄). We found that introducing an electron-withdrawing group (EWG), namely CO₂, significantly facilitates the C–Si bond cleavage and the release of a stabilized anion (**C**) by an accessible TS₅ at 28.6 kcal.mol⁻¹. **C** readily undergoes carboxylation at the carbanionic position to yield **D**, prior to a decarboxylation of the N–CO₂⁻ moiety to 2-PyCO₂⁻. Overall, this mechanism requires a low energy demand of 28.6 kcal mol⁻¹, and it highlights the unique role of CO₂ in this transformation that acts both as a reactant and catalyst able to facilitate the C–Si bond cleavage. In fact, whereas the carboxylation of electron deficient arylsilanes is possible,^[12] the method reported herein uses reversible CO₂ coordination as a means of traceless heteroarene activation. The driving force for the fluoride-mediated carboxylation of **2a₂** derives from the stabilization of anion **C** that is reflected in the decreased pK_a of the pyridine–CO₂ adduct at the 2-position (30.2 versus 43.4 in pyridine, Scheme 3). In contrast, a carbanion at the 3-position of the pyridine–CO₂ adduct is characterized by pK_a of 32.0 and its formation from **21** is 3.9 kcal.mol⁻¹ higher in energy than for **2a₂**, in line with their different reactivities. Based on these findings, the esterification of a variety of heteroaromatic silanes was carried out with CO₂, CH₃I, and TBAT (Scheme 4). Introducing a trifluoromethyl EWG on the 4- or 5-positions of the pyri-



Scheme 3. Fluoride-mediated computed pathway for the carboxylation of **1a₂** and **2a₂** (PBE0-GD3, PCM THF)




Scheme 4. Fluoride-mediated esterification of aromatic silanes with CO₂ and CH₃I. NMR Yields. [a] TBAT (2.0 equiv), **3a** (2.0 equiv).

dine ring somewhat deactivates the reaction at RT, and heating at 70 °C is desirable to reach good conversions into **4fa** and **4ga** (> 99% and 71%, respectively). This behavior results from a destabilization of the pyridine–CO₂ adduct **B**. In contrast, the presence of a methyl group on positions 4, 5, and 6 is beneficial and excellent conversions (72–99%) into the desired methyl picolinates were obtained. Diester **4ha** could be obtained from 2,6-(di-trimethylsilyl)pyridine with excellent 97% yield. The carboxylation of other heterocycles is also feasible; 2-(thienyl)trimethylsilane affords **4ia** in good 75% yield, whereas the furyl derivative **4ja** is obtained under the same conditions in a low yield (27%). For these substrates, it is likely that the activation of heteroaryl silane with CO₂ is unnecessary, as the aromatic anions are stable enough ($pK_a < 35$).^[16] Nonetheless, the pyrrole derivative *N*-methyl(2-trimethylsilyl)pyrrole remained unreacted, even after 40 h at 70 °C. The corresponding *N*-methylpyrrole anion presents a high pK_a of 39.6.^[16] This lack of reactivity can thus be explained by the steric hindrance on the nitrogen center and its absence of nucleophilic character for the coordination of CO₂.

From a practical perspective, heteroaromatic esters **4** are usually prepared by carboxylation of the corresponding *ortho*-lithiated aromatics (e.g. 2-lithiopyridine) with the subsequent addition of an electrophile or by oxidation of methylated heteroaromatics to the carboxylate derivative, followed by condensation of an alcohol.^[15] Given that novel catalytic methods recently appeared to form heteroaromatic silanes by the C–H bond functionalization without the need for a lithiation step,^[17] the carboxylation strategy presented herein offers an attractive pathway to the synthesis of esters **4**. Furthermore, the carboxylation of pyridylsilane derivatives proceeds using CO₂ as a traceless activator (by the formation of stabilized carbanion **C**). This enables the carboxylation of pyridylsilanes at RT and also the use of an electrophile in a one-step procedure for the first time. To further highlight these findings, the polymerization version of the present transformation was attempted [Eq. (3)].

Under an atmosphere of CO₂, disilane **2h** reacted at RT with **3h** to afford a polyester material **4hh**. The gel permeation chromatography (GPC) and NMR data revealed that **4hh** is characterized by a succession of picolinate and butyl moieties. The polymerization degree exceeds four, resulting from the in-



corporation of approximately eight CO₂ molecules (see the Supporting Information). Although many efforts are currently devoted to promoting the incorporation of CO₂ in polycarbonate materials, the formation of oligomer **4hh** represents the first example of CO₂ conversion to a polyester chain, and it offers a novel platform for the utilization of CO₂ in polymer chemistry.^[18] Our current efforts are devoted to expand this proof of concept.

In conclusion, we reported a one-step conversion of CO₂ into heteroaromatic esters under metal-free conditions. Using fluoride anions as promoters for the C–Si bond activation, pyridyl, furanyl, and thienyl organosilanes were successfully carboxylated with CO₂ in the presence of an electrophile. The methodology was applied to 18 different esters and also enabled the incorporation of CO₂ to a polyester material for the first time.

Acknowledgements

For financial support of this work, we acknowledge CEA, CNRS, the University of Paris-Saclay (Fellowship to X.F.), the CHARMMAT Laboratory of Excellence and the European Research Council (ERC Starting Grant Agreement no. 336467). T.C. thanks the Foundation Louis D. - Institut de France for its formidable support. We thank Dr. G. Carrot (CEA) for collecting GPC data.

Keywords: carbon dioxide · density functional calculations · fluorides · silanes · synthetic methods

- [1] a) S. Perathoner, G. Centi, *ChemSusChem* **2014**, *7*, 1274–1282; b) A. Goepfert, M. Czaun, J.-P. Jones, G. K. S. Prakash, G. A. Olah, *Chem. Soc. Rev.* **2014**, *43*, 7995–8048; c) F. J. Fernández-Alvarez, A. M. Aitani, L. A. Oro, *Catal. Sci. Technol.* **2014**, *4*, 611–624; d) A. Tlili, X. Frogneux, E. Blondiaux, T. Cantat, *Angew. Chem. Int. Ed.* **2014**, *53*, 2543–2545; *Angew. Chem.* **2014**, *126*, 2577–2579; e) G. Fiorani, W. Guo, A. W. Kleij, *Green Chem.* **2015**, *17*, 1375–1389.
- [2] a) A. Tlili, E. Blondiaux, X. Frogneux, T. Cantat, *Green Chem.* **2015**, *17*, 157–168; b) C. Das Neves Gomes, O. Jacquet, C. Villiers, P. Thuéry, M. Ephritikhine, T. Cantat, *Angew. Chem. Int. Ed.* **2012**, *51*, 187–190; *Angew. Chem.* **2012**, *124*, 191–194; c) O. Jacquet, X. Frogneux, C. D. N. Gomes, T. Cantat, *Chem. Sci.* **2013**, *4*, 2127–2131; d) Y. Li, X. Fang, K. Junge, M. Beller, *Angew. Chem. Int. Ed.* **2013**, *52*, 9568–9571; *Angew. Chem.* **2013**, *125*, 9747–9750; e) Y. Li, I. Sorribes, T. Yan, K. Junge, M. Beller, *Angew. Chem. Int. Ed.* **2013**, *52*, 12156–12160; *Angew. Chem.* **2013**, *125*, 12378–12382; f) K. Beydoun, T. vom Stein, J. Klankermayer, W. Leitner, *Angew. Chem. Int. Ed.* **2013**, *52*, 9554–9557; *Angew. Chem.* **2013**, *125*, 9733–9736; g) X. Cui, X. Dai, Y. Zhang, Y. Deng, F. Shi, *Chem. Sci.* **2014**, *5*, 649–655.
- [3] a) Y. Tsuji, T. Fujihara, *Chem. Commun.* **2012**, *48*, 9956–9964; b) K. Huang, C.-L. Sun, Z.-J. Shi, *Chem. Soc. Rev.* **2011**, *40*, 2435–2452; c) M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann, F. E. Kühn, *Angew.*

- Chem. Int. Ed.* **2011**, *50*, 8510–8537; *Angew. Chem.* **2011**, *123*, 8662–8690.
- [4] a) T. Ohishi, M. Nishiura, Z. Hou, *Angew. Chem. Int. Ed.* **2008**, *47*, 5792–5795; *Angew. Chem.* **2008**, *120*, 5876–5879; b) Y. Makida, E. Marelli, A. M. Z. Slawin, S. P. Nolan, *Chem. Commun.* **2014**, *50*, 8010–8013; c) X. Zhang, W.-Z. Zhang, L.-L. Shi, C.-X. Guo, L.-L. Zhang, X.-B. Lu, *Chem. Commun.* **2012**, *48*, 6292–6294; d) K. Ukai, M. Aoki, J. Takaya, N. Iwasawa, *J. Am. Chem. Soc.* **2006**, *128*, 8706–8707.
- [5] a) M. Shi, K. M. Nicholas, *J. Am. Chem. Soc.* **1997**, *119*, 5057–5058; b) R. Johansson, O. F. Wendt, *Dalton Trans.* **2007**, 488–492; c) D. P. Hruszkewycz, J. Wu, N. Hazari, C. D. Incarvito, *J. Am. Chem. Soc.* **2011**, *133*, 3280–3283; d) J. Wu, N. Hazari, *Chem. Commun.* **2011**, *47*, 1069–1071.
- [6] a) H. Ohmiya, M. Tanabe, M. Sawamura, *Org. Lett.* **2011**, *13*, 1086–1088; b) H. A. Duong, P. B. Huleatt, Q.-W. Tan, E. L. Shuying, *Org. Lett.* **2013**, *15*, 4034–4037.
- [7] a) F. Corral-Bautista, L. Klier, P. Knochel, H. Mayr, *Angew. Chem. Int. Ed.* **2015**, *54*, 12497–12500; b) G. Hagen, H. Mayr, *J. Am. Chem. Soc.* **1991**, *113*, 4954–4961.
- [8] a) T. Mita, M. Sugawara, K. Saito, Y. Sato, *Org. Lett.* **2014**, *16*, 3028–3031; b) T. Mita, Y. Higuchi, Y. Sato, *Chem. Eur. J.* **2013**, *19*, 1123–1128.
- [9] T. Mita, Y. Higuchi, Y. Sato, *Org. Lett.* **2014**, *16*, 14–17.
- [10] T. Mita, K. Michigami, Y. Sato, *Org. Lett.* **2012**, *14*, 3462–3465.
- [11] M. Yonemoto-Kobayashi, K. Inamoto, Y. Tanaka, Y. Kondo, *Org. Biomol. Chem.* **2013**, *11*, 3773–3775.
- [12] F. Effenberger, W. Spiegler, *Chem. Ber.* **1985**, *118*, 3900–3914.
- [13] a) C.-P. Lin, P. Florio, E. M. Campi, C. Zhang, D. P. Fredericks, K. Saito, W. R. Jackson, M. T. W. Hearn, *Tetrahedron* **2014**, *70*, 8520–8531; b) A. C. Carmo Jr., L. K. C. de Souza, C. E. F. da Costa, E. Longo, J. R. Zamian, G. N. da Rocha Filho, *Fuel* **2009**, *88*, 461–468; c) S. Saleh, B. Sweileh, S. O. Taha, R. Mahmoud, M. O. Taha, *Molecules* **2011**, *16*, 933–950; d) A. Riahi, M. Wurster, M. Lalk, U. Lindequist, P. Langer, *Bioorg. Med. Chem.* **2009**, *17*, 4323–4326.
- [14] C. Eaborn, *J. Chem. Soc.* **1952**, 2846–2849.
- [15] a) G. Black, E. Depp, B. B. Corson, *J. Org. Chem.* **1949**, *14*, 14–21; b) D. G. Anderson, D. E. Webster, *J. Organomet. Chem.* **1968**, *13*, 113–116; c) T. Kamei, K. Fujita, K. Itami, J. Yoshida, *Org. Lett.* **2005**, *7*, 4725–4728.
- [16] K. Shen, Y. Fu, J.-N. Li, L. Liu, Q.-X. Guo, *Tetrahedron* **2007**, *63*, 1568.
- [17] A. A. Toutov, W.-B. Liu, K. N. Betz, A. Fedorov, B. M. Stoltz, R. H. Grubbs, *Nature* **2015**, *518*, 80–84.
- [18] R. Nakano, S. Ito, K. Nozaki, *Nat. Chem.* **2014**, *6*, 325–331.

Received: December 18, 2015

Published online on January 25, 2016