

phthalic anhydrides has been known for over half a century.^{2,3}

(2) Tiemann and Krüger, *Ber.*, **29**, 901 (1896).

(3) Fuller and Kenyon, *J. Chem. Soc.*, **126**, 2309 (1924).


DEPARTMENT OF CHEMISTRY
BATTERSEA POLYTECHNIC

LONDON S. W. 11, ENGLAND RECEIVED JANUARY 27, 1948

Rearrangement of Phenylethynylcarbinol

BY WARREN S. MACGREGOR

Rupe¹ reported that several tertiary acetylenic alcohols containing a free acetylenic hydrogen were rearranged upon boiling with 85% formic acid to the corresponding α,β -unsaturated aldehydes. Thus, the ethynylcarbinols from tetrahydrocarvone and fenchone were reported to yield 5 - isopropyl - 2 - methylcyclohexylidene - 1 - acetaldehyde and 1,3,3-trimethylbicyclo[2,2,1]heptylidene-2-acetaldehyde, respectively.² The reaction represents a special case of a Meyer-Schuster rearrangement³ and was considered by Rupe¹ to proceed in analogous manner



Reinvestigation of several of the reported rearrangements by Fischer and Lowenberg,⁴ Hurd and Christ⁵ and others demonstrated that the principal products were unsaturated ketones presumably formed by dehydration of the carbinols followed by hydration of the triple bond.⁵ Thus, 1-ethynylcyclohexanol yielded 1-acetylcyclohexene^{4,5} rather than cyclohexylidene acetaldehyde. Chanley⁶ showed that both the ketone and aldehyde products resulted from the rearrangement of 1-ethynyl-1-cyclohexanols although the aldehydes were formed in very low yield (0.8 to 6%).

The reaction probably involves initial elimination of the hydroxyl group to form a carbonium ion. The loss of a proton from an adjacent carbon followed by hydration of the triple bond and ketonization would yield the observed unsaturated ketone. Shifting of the carbonium ion bonds to the allenic structure followed by hydroxylation at the positive terminal carbon would yield the enolic form of the unsaturated aldehyde which is in equilibrium with the predominantly favored aldehyde.

A compound such as phenylethynylcarbinol would yield a carbonium ion that could not form an unsaturated ketone by the above mechanism and hence, barring alternative modes of reaction, should yield cinnamaldehyde as the principal rearranged product. When the carbinol was heated with 30% sulfuric acid, 85% phosphoric acid or phthalic anhydride the odor of cinnamaldehyde

was evident but the material was largely polymerized to a tar. The rearrangement was achieved with less polymerization by steam distilling the carbinol through 28% sulfuric acid. The resultant crude cinnamaldehyde in two experiments was converted directly to the phenylhydrazone (35% over-all yield). In a third experiment the aldehyde was converted to α -bromocinnamaldehyde (21% over-all yield).

Experimental

Rearrangement of Phenylethynylcarbinol.—Phenylethynylcarbinol (5.4 g.) and 5 cc. of water were placed in a 25-cc. distilling flask fitted for steam distillation with the side arm extending nearly to the bottom of the 50-cc. flask of a Clark acetyl apparatus⁷ containing 25 cc. of 28% sulfuric acid. The flasks were heated in oil-baths to 115°, a steam generator connected to the small flask, and steam distillation started adjusting the bath temperatures to maintain approximately constant volume in the flasks. A water-insoluble oil (3.1 g.) separated in the distillate and dissolved in 5 cc. of ethanol. A solution of 17.5 g. phenylhydrazine hydrochloride and 10.0 g. sodium acetate in 150 cc. of water then was added to the oil and the mixture warmed for fifteen minutes on a water-bath. After filtering, washing with water, ethanol and ether, and drying, 3.0 g. (33%) of crude cinnamalphenylhydrazone (m. p. 154–157°) was obtained. Two recrystallizations from ethanol raised the m. p. to 168–169° and a mixed m. p. with an authentic sample showed no depression. A second experiment yielded 1.2 g. (36%) of the crude phenylhydrazone (m. p., 161–165°) from 2.0 g. of the carbinol.

The crude cinnamaldehyde from a similar rearrangement of 6.2 g. of phenylethynylcarbinol was extracted from the distillate with ether, and upon removal of solvent the oil was taken up in 10 cc. of glacial acetic acid. A solution of 10.7 g. of bromine in 19.9 g. of acetic acid was added dropwise with stirring and cooling in an ice-bath until the bromine color remained for ten minutes (10.3 g. of the solution, corresponding to 3.0 g. cinnamaldehyde, was required.) Potassium carbonate (1.5 g.) was then added and the mixture allowed to stand overnight at room temperature. After refluxing thirty-five minutes, the mixture was cooled and poured into 30 cc. of water, whereupon an oil separated that partially crystallized upon shaking. The oil and crystals were removed and taken up in 10 cc. of warm ethanol. Upon cooling and seeding 2.1 g. (21%) of α -bromocinnamaldehyde, m. p., 71.5–72°, was obtained. Recrystallization from ethanol did not alter the m. p., and the mixed m. p. with a sample prepared following the directions of Allen and Edens⁸ showed no depression.

(7) Clark, *Ind. Eng. Chem., Anal. Ed.*, **8**, 487 (1936).

(8) Allen and Edens, "Organic Syntheses," **25**, 92 (1945).

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF PORTLAND
PORTLAND 3, OREGON

RECEIVED JUNE 1, 1948

Indium Orthovanadate¹

BY W. O. MILLIGAN, HENRY H. RACHFORD, JR., AND L. MERTEN WATT

Forty-seven years ago Renz² prepared a gel which he designated as $\text{In}(\text{VO}_3)_3 \cdot 2\text{H}_2\text{O}$ by the addition of a sodium metavanadate solution to an indium chloride solution. The supposed constitution of the gel was based merely upon a chemical analysis, and there does not exist in the literature.

(1) Presented before the Texas Regional Meeting of the American Chemical Society held in Austin, Texas, December 7–8, 1945.

(2) Renz, *Ber.*, **34**, 2765 (1901).

- (1) Rupe and Kambli, *Helv. Chim. Acta*, **9**, 672 (1926).
- (2) Rupe and Keunzy, *Helv. Chim. Acta*, **14**, 708 (1931).
- (3) Meyer and Schuster, *Ber.*, **55**, 819 (1922).
- (4) Fischer and Lowenberg, *Ann.*, **475**, 183 (1929).
- (5) Hurd and Christ, *THIS JOURNAL*, **59**, 118 (1937).
- (6) Chanley, *THIS JOURNAL*, **70**, 244 (1948).