Nov., 1937

## Summary

From an analysis and reinterpretation of the results of Tschesche and Haupt it was concluded that the formula for convallatoxin tentatively suggested by these investigators is inadmissible and that the (unisolated) aglycone in question probably is identical with strophanthidin. Comparisons made both in this Laboratory and by Dr. Tschesche have now established the correctness of this hypothesis.

Converse Memorial Laboratory Cambridge, Massachusetts

**RECEIVED SEPTEMBER 23, 1937** 

[CONTRIBUTION FROM THE CHEMISTRY LABORATORY OF THE UNIVERSITY OF MICHIGAN] The Reaction of Alkali Metals with Polycyclic Hydrocarbons. II

BY W. E. BACHMANN AND L. H. PENCE<sup>1</sup>

It was shown recently that sodium and lithium add to the meso positions of 1,2-benzanthracene, 1.2.5.6-dibenzanthracene and 3-methylcholanthrene.<sup>2</sup> We have now extended this reaction to include 5-methyl-1,2-benzanthracene, 5-phenyl-1,2-benzanthracene, 1,2,3,4-dibenzanthracene, 6methyl-1,2,3,4-dibenzanthracene, 1,2,6,7-dibenzanthracene and 1,2,3,4,5,6-tribenzanthracene. Intensely colored addition compounds are formed when solutions of these hydrocarbons in etherbenzene are shaken with sodium or with lithium. Thus, 1.2.3.4-dibenzanthracene gives an intensely lavender colored solution of 9,10-disodium-9,10dihydro-1,2,3,4-dibenzanthracene (I), while with lithium the intensely purple 9,10-dilithium addition product is formed.



As a rule the color of the disodium addition product is different from that of the dilithium derivative. This difference is ascribed to a difference in the mode of addition of the two metals, the one adding in *cis* positions, the other in *trans* positions. On treatment with methanol, the alkali addition products of the hydrocarbons yield the 9,10-dihydro derivatives (II, for example). The same dihydro derivative is obtained from both the disodium and the dilithium addition compounds, except in the case of 1,2,6,7-dibenzanthracene (III). In the reaction between the alkali metals and 1,2,6,7-dibenzanthracene it was expected that addition could take place at the 5,8- or the 9,10-positions. Actually, a mixture of the two possible dihydro derivatives was obtained from the sodium addition product and methanol. A single dihydro compound was isolated from the dilithium addition product, although the structure of this compound was not determined.

The structures assigned to the dihydro derivatives and hence to the alkali metal addition products are based on the oxidation of the dihydro derivatives to the 9,10-diones by chromic acid. All of the 9,10-dihydro derivatives are dehydrogenated readily by sulfur to the parent hydrocarbons.

In order to obtain water-soluble derivatives of the polycyclic hydrocarbons, we have investigated the action of carbon dioxide on a number of the



alkali addition products. In this reaction a distinct difference in behavior was encountered

<sup>(1)</sup> From the Ph.D. dissertation of L. H. Pence.

<sup>(2)</sup> Bachmann, J. Org. Chem., 1, 347 (1936).

between the disodium and the dilithium addition compounds. Thus, while the dilithium compound obtained from 1,2,5,6-dibenzanthracene yielded the expected dilithium salt of 9,10-dicarboxy-9,10-dihydro-1,2,5,6-dibenzanthracene (IV) exclusively, the disodium compound behaved in an erratic manner, not always giving the same compound even when the identical conditions appeared to be employed. Usually the monocarboxylic acid (V) was obtained from the disodium compound. The dicarboxylic acid (V1) was obtained from the dilithium addition product of 3-methylcholanthrene, the disodium compound yielding a monocarboxylic acid. A similar result was obtained with 1,2,3,4-dibenzanthracene.

From the reaction between carbon dioxide and the sodium addition compound of anthracene, Schlenk and Bergmann<sup>3</sup> isolated a dicarboxylic acid. When, however, anthracene was added to the disodium addition compound prior to the introduction of carbon dioxide, a monocarboxylic acid was formed; the latter compound was also obtained when the reaction between the anthracene and sodium had been incomplete. The excellent yields of dihydro compounds obtainable from our sodium addition compounds by reaction with methanol precluded an explanation of the formation of a monocarboxylic acid on the basis of an incomplete reaction between the hydrocarbon and sodium. Moreover, none of the parent hydrocarbon was isolated from the reaction mixture after treatment with carbon dioxide. It seems likely that the disodium compound yields an unstable dicarboxylic acid which subsequently loses a molecule of carbon dioxide.

The sodium and potassium salts of the acids are soluble in water and it is in this form that they are being tested for carcinogenic activity. Work is in progress on the reaction between the alkali metal addition compounds and alkyl and acyl halides.

#### Experimental

1,2,3,4-Dibenzanthracene.—A mixture of 14.8 g. of 9-(o-toluoyl)-phenanthrene<sup>4</sup> and 5 g. of zinc dust was heated at 400–420° for three hours. The hydrocarbon, which was obtained by distillation at 2 mm. pressure, was purified by recrystallization from xylene–alcohol; m. p. 200– 201.5°; weight, 7.8 g. (56%), a yield higher than that reported previously.

6-Methyl-1,2,3,4-dibenzanthracene.—To the Grignard reagent prepared from 30 g. of 1-bromo-2,4-dimethyl-

(4) Bachmann and Pence, THIS JOURNAL, 57, 1130 (1935).

benzene<sup>5</sup> in 50 cc. of ether was added 100 cc. of benzene and 30 g. of 9-cyanophenanthrene. After being refluxed for two days, the mixture was hydrolyzed with ice-cold hydrochloric acid, which precipitated a solid ketone imine hydrochloride. The ketone, 9-(2',4'-dimethylbenzoyl)-phenanthrene, resulting from the hydrolysis of the ketone-imine hydrochloride by boiling water, was obtained as a colorless oil by distillation at 200-250° at 3 mm.; weight, 34 g. (75%).

The aforementioned ketone was kept at  $420^{\circ}$  for two and one-half hours and the product was distilled at 3 mm. pressure. Two recrystallizations from xylene, using decolorizing charcoal, gave 4.0 g. (12%) of 6-methyl-1,2,3,4dibenzanthracene as fine, colorless needles; m. p. 157.5-158°. The hydrocarbon is only slightly soluble in cold alcohol and acetic acid.

Anal. Calcd. for C<sub>22</sub>H<sub>16</sub>: C, 94.5; H, 5.5. Found: C, 94.6; H, 5.5.

6-Methyl-1,2,3,4-dibenzanthraquinone was formed by refluxing a mixture of 100 mg. of the hydrocarbon, 200 mg. of sodium dichromate dihydrate and 10 cc. of acetic acid for thirty minutes. After sublimation at 230-260° (0.2 mm.), the quinone crystallized from acetic acid in long yellow needles; yield 65%; m. p. 187.5-188.5°.

Anal. Calcd. for  $C_{23}H_{14}O_2$ : C, 85.7; H, 4.4. Found: C, 85.2; H, 4.6.

1,2,6,7-Dibenzanthracene.—This hydrocarbon has been prepared previously but not from crystalline 3-(o-toluoyl)phenanthrene. By heating a mixture of 6.6 g. of the crystalline ketone<sup>4</sup> and 2.2 g. of zinc dust at 400-420° for two hours, distilling the product at 3 mm. pressure and recrystallizing the distillate twice from xylene, 1.34 g. of 1,2,6,7-dibenzanthracene was obtained as golden-yellow crystals. The highest melting point which we observed was 262°, while Clar<sup>6</sup> reported a value of 263-264°.

1,2,3,4,5,6-Tribenzanthracene.—After heating 15.4 g. of 9-(2'-methyl-1'-naphthoyl)-phenanthrene<sup>4</sup> with 5 g. of zinc dust at 410° for two and three-quarter hours, the product was distilled at 3 mm. pressure. By recrystallization from xylene 6.4 g. (44%) of 1,2,3,4,5,6-tribenzanthracene was obtained; m. p. 225-228°. Fieser and Dietz<sup>7</sup> obtained a 24% yield of the hydrocarbon melting at 224° by pyrolysis of the ketone without the addition of zinc dust.

Preparation of the 9,10-Dihydro Derivatives of the Polycyclic Hydrocarbons.—A mixture of 0.5 g. of the hydrocarbon, 1 g. of powdered sodium, 50 cc. of anhydrous ether, 50 cc. of dry benzene and a few sharp particles of glass<sup>8</sup> was shaken in absence of air for two days. The intense color which formed after a few minutes usually was replaced by still another color, that of the 9,10-disodium addition compound. After the color had been completely discharged by addition of methanol, the solution was separated from the excess of sodium, shaken with dilute hydrochloric acid, filtered and evaporated. The dihydro compound which remained was then recrystallized.

For the reaction with lithium, the same procedure was

<sup>(3)</sup> Schlenk and Bergmann, Ann., 463, 134 (1928).

<sup>(5)</sup> Morgan and Coulson, J. Chem. Soc., 2203 (1929).

<sup>(6)</sup> Clar, Ber., 62, 1574 (1929).

<sup>(7)</sup> Fieser and Dietz, ibid., 62, 1827 (1929).

<sup>(8)</sup> Compare Bachmann and Wiselogle, THIS JOURNAL, 58, 1943 (1936).

used except that 0.8 g. of lithium wire (0.5 mm. diameter) was employed. The reaction was initiated by pressing the lithium wire in several places with a glass rod. The mixture was then shaken for five to seven days.

9,10 - Dihydro - 5 - methyl - 1,2 - benzanthracene obtained from the intensely wine-red disodium addition product of 5-methyl-1,2-benzanthracene crystallized from acetone-ethanol in thin, colorless plates; yield 57%; m. p. 128-129.5°.

Anal. Calcd. for  $C_{10}H_{16}$ : C, 93.4; H, 6.6. Found: C, 93.5; H, 6.5.

9,10 - Dihydro - 5 - phenyl - 1,2 - benzanthracene obtained from the brownish-purple disodium compound of 5-phenyl-1,2-benzanthracene<sup>9</sup> crystallized from acetic acid in glistening, colorless needles; yield, 90%; m. p. 96-96.5°.

Anal. Calcd. for  $C_{24}H_{18}$ : C, 94.1; H, 5.9. Found: C, 94.4; H, 6.0.

9,10-Dihydro-1,2,3,4-dibenzanthracene obtained from the dark lavender disodium addition compound of 1,2,-3,4-dibenzanthracene crystallized from benzene-ethanol in colorless needles; yield 84%; m. p. 202-203°. The identical compound was obtained in 64% yield from the purple dilithium addition compound of the hydrocarbon.

Anal. Calcd. for  $C_{22}H_{16}$ : C, 94.2; H, 5.8. Found: C, 93.8; H, 6.1.

9,10 - Dihydro - 6 - methyl - 1,2,3 4 - dibenzanthracene was obtained in 80% yield from both the reddish-purple disodium compound and the bluish-purple dilithium compound of 6-methyl-1,2,3,4-dibenzanthracene. It crystallized from benzene in fine, colorless needles; m. p. 207-209.5°.

Anal. Caled. for  $C_{23}H_{18}$ : C, 93.8; H, 6.2. Found: C, 93.6; H, 6.3.

9,10 - Dihydro - 1,2,3,4,5,6 - tribenzanthracene, obtained from the dark blue disodium compound of 1,2,3,4,5,6-tribenzanthracene in 73% yield and from the blue dilithium derivative in 40% yield, crystallized from xylene in colorless needles; m. p.  $281-283^{\circ}$ .

Anal. Calcd. for  $C_{26}H_{18}$ : C, 94.5; H, 5.5. Found: C, 94.1; H, 5.6.

The mixture of compounds obtained from the bluishgreen sodium addition compound of 1,2,6,7-dibenzanthracene and methanol melted at  $170-200^{\circ}$ . From the bluishgreen lithium addition product a 73% yield of one of the two possible dihydro-1,2,6,7-dibenzanthracenes was isolated as fine, colorless needles; m. p.  $157-160.5^{\circ}$ . It was not possible to determine whether this compound was the 5,8- or the 9,10-dihydro derivative.

Anal. Calcd. for  $C_{22}H_{16}$ : C, 94.2; H, 5.8. Found: C, 94.2; H, 5.8.

Dehydrogenation of the 9,10-Dihydro Derivatives.—A mixture of 50 mg. of the dihydro derivative and 10 mg. of sulfur was heated at  $200-230^{\circ}$  for one hour. The product was dissolved in toluene and the solution was shaken with mercury for several hours in order to remove the excess of sulfur. From the solution the parent hydrocarbon was isolated in good yield. Thus, a 78% yield of 1,2,3,4-

dibenzanthracene was obtained by dehydrogenation of 9,10-dihydro-1,2,3,4-dibenzanthracene.

Oxidation of the 9,10-Dihydro Derivatives.—A solution of 50 mg. of the 9,10-dihydro derivative and slightly more than the calculated amount of sodium dichromate dihydrate in 10 cc. of acetic acid was heated at the boiling point for forty-five minutes. The 9,10-dione, which precipitated on adding dilute sulfuric acid to the mixture, was purified by sublimation at 0.2 mm., followed by recrystallization from acetic acid. In each case, the 9,10-dione proved to be identical with the quinone obtained by oxidizing the parent hydrocarbon. For example, from 9,10dihydro-1,2,3,4-dibenzanthracene a 77% yield of 1,2,3,4dibenzanthraquinone was obtained, which was identical with the product formed by oxidation of 1,2,3,4-dibenzanthracene.

Our 1,2,3,4,5,6-tribenzanthraquinone melted at 246.5-247.5°, while the reported value<sup>7</sup> is 244°. Oxidation of the dihydro-1,2,6,7-dibenzanthracene (m. p. 157-160.5°) gave a mixture of quinones of the same nature as that obtained by oxidation of 1,2,6,7-dibenzanthracene.

Reaction of the Alkali Addition Compounds with Carbon Dioxide.—Carbon dioxide (dried by concentrated sulfuric acid and phosphorus pentoxide) was passed into a cooled solution of the 9,10-disodium or 9,10-dilithium addition compound of the hydrocarbons until the color of the mixture was discharged completely. The suspension of the alkali salt of the carboxylic acid was decanted from the excess of the metal and was shaken with ice water. Acidification of the separated aqueous solution precipitated the carboxylic acid, which was then recrystallized. The ether-benzene solution from which the sodium salt had been extracted always yielded a small amount of the 9,10dihydro derivative but none of the parent hydrocarbon. The methyl esters of the acids were all prepared by means of diazomethane.

9,10-Dicarboxy-9,10-dihydro-1,2-benzanthracene, obtained from the dilithium addition compound of 1,2benzanthracene, formed thin, colorless sheets from acetone-benzene; yield 58%; m. p. 252° with evolution of gas.

Anal. Calcd. for  $C_{20}H_{14}O_4$ : C, 75.4; H, 4.4. Found: C, 75.4; H, 4.8.

The **dimethyl ester** crystallized from acetone-ethanol in clusters of thick, colorless plates; m. p. 143.5-145°.

Anal. Calcd. for  $C_{22}H_{18}O_4$ : C, 76.3; H, 5.2. Found: C, 76.1; H, 5.4.

In one run the disodium addition compound gave a crystalline acid but in half a dozen other runs the acid and the methyl ester derived from it proved to be uncrystallizable oils.

9,10-Dicarboxy-9,10-dihydro-1,2,3,4-dibenzanthracene, obtained from the dilithium compound of 1,2,3,4dibenzanthracene, crystallized from xylene in clusters of colorless prisms; yield 65%; m. p.  $270-274^{\circ}$  with evolution of gas. The acid was analyzed in the form of its dimethyl ester, which crystallized from acetone-ethanol in colorless prisms; m. p.  $236-238^{\circ}$ .

Anal. Calcd. for  $C_{26}H_{20}O_4$ : C, 78.8; H, 5.1. Found: C, 78.7; H, 5.3.

The 9-carboxy-9,10-dihydro-1,2,3,4-dibenzanthracene,

<sup>(9)</sup> Bachmann and Bradbury, J. Org. Chem., 2, 175 (1937).

which was obtained from the sodium addition compound in 24% yield, crystallized from acetone-xylene in colorless plates; m. p.  $227.5-230^{\circ}$  with evolution of gas. Since the acid was difficult to purify, it was converted to its methyl ester for analysis. The ester crystallized from acetone ethanol in clusters of colorless prisms; m. p.  $158-164^{\circ}$ .

Anal. Calcd. for  $C_{24}H_{18}O_2$ : C, 85.2; H, 5.4. Found: C, 84.3; H, 5.4.

9,10-Dicarboxy-9,10-dihydro-1,2,5,6-dibenzanthracene (IV), obtained from the dilithium addition compound of 1,2,5,6-dibenzanthracene in 59% yield, crystallized from acetone-benzene in clusters of colorless prisms; m. p.  $262-264^\circ$  with gas evolution. The acid is very soluble in acetone but only slightly soluble in benzene. It was analyzed in the form of its dimethyl ester, which crystallized from acetone-ethanol in stout, colorless needles; m. p.  $255-257^\circ$ .

Anal. Calcd. for  $C_{26}H_{20}O_4$ : C, 78.8; H, 5.1. Found: C, 78.1; H, 5.0.

The reaction between the disodium compound of 1,2,5,6dibenzanthracene and carbon dioxide did not always proceed in the same manner. Usually a 60-70% yield of an acid was obtained which crystallized from benzene in colorless needles; m. p.  $242-255^{\circ}$  with decomposition. Since the acid was difficult to purify, it was converted to its methyl ester, which crystallized from benzene-methanol in broad, colorless needles; m. p.  $174-175^{\circ}$ . Although the ester was not obtained quite pure, analysis showed that the acid was 9-carboxy-9,10-dihydro-1,2,5,6-dibenzanthracene (V).

Anal. Calcd. for C<sub>24</sub>H<sub>18</sub>O<sub>2</sub>: C, 85.2; H, 5.4. Found: C, 84.1; H, 5.3.

In some runs the disodium compound yielded the same dicarboxylic acid that was obtained from the dilithium addition compound. In one run there was obtained a 79% yield of an acid, m. p.  $255-258^\circ$ , whose methyl ester

crystallized in broad, colorless needles which melted at 100-101°. Unfortunately, this product was all used up in certain tests and we were unable to secure any more of it in subsequent experiments.

6,12b-Dicarboxy-6,12b-dihydro-3-methylcholanthrene (VI), was obtained in 42% yield from the dilithium addition compound of 3-methylcholanthrene.<sup>2</sup> From hot benzene the dicarboxylic acid separated in brownish-yellow crystals; m. p.  $242-246^{\circ}$ . Considerable losses were experienced in the process of recrystallization, for the acid appeared to be unstable. Its dimethyl ester crystallized from acetone-methanol in clusters of short, brownish prisms; m. p.  $205-208^{\circ}$ .

Anal. Calcd. for  $C_{2b}H_{22}O_4$ : C, 77.7; H, 5.7. Found: C, 77.1; H, 5.9.

The acid isolated in 59% yield from the disodium addition product crystallized in pale yellow needles; its methyl ester formed pale yellow cubes from acetone-methanol; m. p. 178-179°. Analysis of the latter showed that the acid was a monocarboxy-6,12b-dihydro-3-methylcholanthrene, but whether the carboxyl group was in the 6or 12b-position was left undecided.

Anal. Calcd. for  $C_{23}H_{20}O_2$ : C, 84.1; H, 6.1. Found: C, 83.9; H, 5.5.

### Summary

The reaction between a number of polycyclic hydrocarbons with sodium and with lithium has been investigated. From the intensely colored alkali metal addition compounds of the hydrocarbons the dihydro derivatives were obtained by treatment with methanol. With carbon dioxide the addition compounds yielded salts of carboxylic acids.

ANN ARBOR, MICHIGAN RECEIVED SEPTEMBER 18, 1937

# The Hydroxylation of Unsaturated Substances. III. The Use of Vanadium Pentoxide and Chromium Trioxide as Catalysts of Hydroxylation

## By NICHOLAS A. MILAS

The catalytic hydroxylation of unsaturated substances has been under way in this Laboratory for some time. Of particular usefulness has been the method recently described<sup>1</sup> and extended to embrace a great number of substances.<sup>2</sup> In this method hydrogen peroxide in anhydrous tertiary butyl or tertiary amyl alcohols and in the presence of osmium tetroxide adds onto carbon-carbon double bonds to form glycols. Although the mechanism of this reaction has not been entirely elucidated, evidence has been accumulated to indicate that it proceeds through the formation of perosmic acid which probably dissociates into hydroxyl radicals and osmium tetroxide. In this way the catalyst seems to act, like ultraviolet light,<sup>3</sup> as though it dissociates the hydrogen peroxide into hydroxyl radicals which subsequently add onto double bonds or cause other oxidations.

Like osmium there are a number of metals belonging to the IV, V and VI sub-groups of the Periodic System which are known to form very

(3) Milas, Kurz and Anslow, Jr., ibid., 59, 543 (1937).

<sup>[</sup>Contribution from the Research Laboratory of Organic Chemistry, Massachusetts Institute of Technology, No. 158]

<sup>(1)</sup> Milas and Sussman, THIS JOURNAL, 58, 1302 (1936).

<sup>(2)</sup> Milas and Sussman, *ibid.*, **59**, 2545 (1937); also unpublished results.