Contribution from Rocketdyne, a Division of Rockwell International, Canoga Park, California 91304

# **Perfluoroammonium and Cesium Fluorotungstates**

## WILLIAM W. WILSON and KARL O. CHRISTE\*

#### Received May 22, 1981

The syntheses of NF4<sup>+</sup> salts containing fluorotungstate anions were studied. The new NF4WOF5 salt was prepared by the reaction of WOF<sub>4</sub> with a concentrated solution of  $NF_4HF_2$  in anhydrous HF. It is a white solid, stable up to 60 °C At higher temperatures it decomposes to  $NF_4W_2O_2F_9$ ,  $NF_3$ ,  $OF_2$ , and  $WF_6$ . The ionic nature of  $NF_4WOF_5$  in the solid state was established by vibrational spectroscopy, but in dilute HF solutions the  $WOF_5^-$  anion solvolyzes according to  $2WOF_5^-$ + HF  $\rightleftharpoons$  W<sub>2</sub>O<sub>2</sub>F<sub>9</sub><sup>-</sup> + HF<sub>2</sub><sup>-</sup>. In HF solution WF<sub>6</sub><sup>-</sup> is readily oxidized by NF<sub>4</sub><sup>+</sup> to yield WF<sub>6</sub> and NF<sub>3</sub>. The syntheses and some reaction chemistry and properties of CsWOF<sub>5</sub>, CsW<sub>2</sub>O<sub>2</sub>F<sub>9</sub>, Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>, and CsWF<sub>6</sub> are discussed.

# Introduction

The synthesis of a stable NF4<sup>+</sup> salt containing a fluorotungstate anion is of significant interest for solid-propellant  $WF_6$  gas generators. In this paper the synthesis and characterization of the first known example of a stable NF<sub>4</sub><sup>+</sup> fluorotungstate salt are reported, and some reaction chemistry of various cesium fluorotungstates in anhydrous HF is discussed.

#### **Experimental Section**

Apparatus. Volatile materials used in this work were handled in a stainless steel-Teflon FEP vacuum line. The line and other hardware used were well passivated with ClF<sub>3</sub> and, if HF was to be used, with HF. Nonvolatile materials were handled in the dry nitrogen atmosphere of a glovebox. Metathetical reactions were carried out in HF solution with use of an apparatus consisting of two FEP U-traps interconnected through a coupling containing a porous Teflon filter.<sup>1</sup> Thermal decomposition measurements were carried out in a previously described<sup>2</sup> sapphire reactor.

Infrared spectra were recorded in the range 4000-200 cm<sup>-1</sup> on a Perkin-Elmer Model 283 spectrophotometer. Spectra of solids were obtained by using dry powders pressed between AgCl windows in an Econo press (Barnes Engineering Co.). Spectra of gases were obtained by using a Teflon cell of 5-cm path length equipped with AgCl windows.

Raman spectra were recorded on a Cary Model 83 spectrophotometer using the 4880-Å exciting line of an Ar ion laser and a Claassen filter<sup>3</sup> for the elimination of plasma lines. Sealed glass, Teflon FEP, or Kel-F tubes were used as sample containers in the transverse-viewing transverse-excitation mode. Lines due to the Teflon or Kel-F sample tubes were suppressed by the use of a metal mask.

The <sup>19</sup>F NMR spectra were recorded at 84.6 MHz on a Varian Model EM390 spectrometer equipped with a variable-temperature probe. Chemical shifts were determined relative to external CFCl<sub>3</sub> with positive shifts being downfield from the standard.<sup>4</sup>

Elemental analyses were carried out as previously described.<sup>5</sup>

**Materials.** Literature methods were used for the syntheses of  $NF_4SbF_6$ ,  $^6NF_4HF_2$  solutions in HF,  $^7$  and WOF<sub>4</sub>.<sup>8</sup> Hydrogen fluoride (Matheson) was dried by storage over  $BiF_5$  to remove the  $H_2O.9$ Tungsten hexafluoride (Alfa) and CsI (Alfa, 99.9%) were used as received. Cesium fluoride (KBI) was dried by fusion in a platinum crucible and ground in the drybox. Sulfur dioxide (Matheson) was dried over  $P_2O_5$  and purified by fractional condensation prior to use.

Preparation of CsWOF5. Cesium fluoride (9.93 mmol) was loaded inside the drybox into a prepassivated Teflon FEP ampule. Anhydrous

- (6) Christe, K. O.; Schack, C. J.; Wilson, R. D. J. Fluorine Chem. 1976, , 541.
- Christe, K. O.; Wilson, W. W.; Wilson, R. D. Inorg. Chem. 1980, 19, (7) 1494.
- Paine, R. T.; McDowell, R. S. Inorg. Chem. 1974, 13, 2366. Christe, K. O.; Wilson, W. W.; Schack, C. J. J. Fluorine Chem. 1978,
- (9) 11, 71.

HF (15 mL, liquid) was added to the ampule on the vacuum line, and 9.9 mmol of distilled H<sub>2</sub>O was added with a syringe. The mixture was allowed to homogenize, and WF<sub>6</sub> (9.96 mmol) was added in vacuo at -196 °C. After the ampule was kept at ambient temperature for several hours, all material volatile at 55 °C was pumped off. On the basis of its weight and vibrational spectra, the solid residue was mainly CsWOF<sub>5</sub><sup>10,11</sup> containing only small amounts of CsW<sub>2</sub>O<sub>2</sub>F<sub>9</sub><sup>12</sup> and CsHF<sub>2</sub>.

Preparation of CsW2O2F9 by Solvolysis of CsWOF5 in HF. A sample of CsWOF<sub>5</sub> (2.35 mmol) in a Teflon ampule was stirred in anhydrous HF (10 mL, liquid) for 12 h at 24 °C. The undissolved material was separated from the solution by filtration at 24 °C, and all volatile material was pumped off at 55 °C. On the basis of the observed mass balance and vibrational spectra, the filter cake consisted of  $C_8W_2O_2F_9^{12}$  (~1.1 mmol) and the filtrate residue consisted of CsHF<sub>2</sub> ( $\sim$ 1.2 mmol) containing as an impurity Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>,<sup>13</sup> which was formed by hydrolysis of some CsWOF<sub>5</sub>.

Preparation of Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>. Cesium fluoride (37 mmol) and 20 mL of 48% aqueous HF were combined in a Teflon FEP ampule to which  $WF_6$  (18.47 mmol) was added at -196 °C on the vacuum line. When the mixture warmed to room temperature, initially a white precipitate formed, which subsequently dissolved, resulting in a clear solution. The solution was poured into a 250-mL Teflon beaker, and the aqueous HF solvent was evaporated on a hot plate at 60 °C. The resulting white residue (10.24 g, weight calculated for 18.47 mmol of Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub> = 10.40 g) was shown by vibrational spectroscopy to be essentially pure cis-Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>.<sup>13</sup> Treatment of Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub> with anhydrous HF resulted in the partial conversion of Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub> to CsWOF<sub>5</sub>, Cs-W<sub>2</sub>O<sub>2</sub>F<sub>9</sub>, and CsHF<sub>2</sub>

Preparation of CsWF<sub>6</sub>. Cesium iodide (21.2 mmol) was loaded in the drybox into a prepassivated Teflon U-tube, equipped with a Teflon filter, and anhydrous HF (26 mL, liquid) was added at -196 °C on the vacuum line. On warming toward ambient temperature, the HF reacted with the CsI to form HI, which accumulated in the bottom of the U-tube as a separate liquid phase. Tungsten hexafluoride (32.46 mmol) was added to the U-tube at -196 °C, and the mixture was allowed to warm to ambient temperature. A copious precipitate of black iodine crystals formed. The mixture was kept at 25 °C for 12 h, and the clear solution was filtered into a second prepassivated Teflon U-tube attached through Teflon tubing and valves to the first U-tube. The HF solvent and small amounts of  $I_2$  were pumped off from the contents of the second U-tube at 55 °C for 2 h, resulting in a slightly off-white solid (4.856 g, weight calculated for 21.2 mmol of  $C_{s}WF_{6} = 9.144$  g), which was identified by vibrational spectroscopy<sup>14-18</sup> as  $C_{s}WF_{6}$ . A second extraction of the precipitate with 16 mL of liquid HF resulted in an additional 3.011 g of  $CsWF_6$ , indicating that CsWF<sub>6</sub> is only moderately soluble in HF.

- Beuter, A.; Sawodny, W. Z. Anorg. Allg. Chem. 1976, 427, 37. (10)
- Beuter, A.; Sawodny, W. Angew. Chem., Int. Ed. Engl. 1972, 11, 1020. Bougon, R.; Bui Huy, T.; Charpin, P. Inorg. Chem. 1975, 14, 1822. (11)
- 12) Pausewang, G.; Schmitt, R.; Dehnicke, K. Z. Anorg. Allg. Chem. 1974, (13)
- 408, 1 and references cited therein.
- (14)
- Peacock, R. D.; Sharp, D. W. A. J. Chem. Soc. 1959, 2762. Beuter, A. Dissertation, University of Ulm, Ulm, West Germany, 1974. Heath, G. A.; Hefter, G. T.; Boyle, T. W.; Desjardins, C. D.; Sharp, D. W. A. J. Fluorine Chem. 1978, 11, 399. (16)
- Prescott, A.; Sharp, D. W. A.; Winfied, J. M. J. Chem. Soc., Dalton Trans. 1975, 936. (17)
- (18) Shamir, J.; Malm, J. G. Inorg. Nucl. Chem. Lett. 1976, 107.

Christe, K. O.; Schack, C. J.; Wilson, R. D. Inorg. Chem. 1977, 16, 849. (1)(2) Christe, K. O.; Wilson, R. D.; Goldberg, I. B. Inorg. Chem. 1979, 18, 2572.

Claassen, H. H.; Selig, H.; Shamir, J. Appl. Spectrosc. 1969, 23, 8. Pure Appl. Chem. 1972, 29, 627. Rushworth, R.; Schack, C. J.; Wilson, W. W.; Christe, K. O. Anal. (5) Chem. 1981, 53, 845.

Reaction of CsI with SO<sub>2</sub>. Dry CsI (6.30 mmol) was loaded in the drybox into a Teflon ampule, and SO<sub>2</sub> (51.4 mmol), which had been stored over  $P_2O_5$ , was added at -196 °C on the vacuum line. On warming, all the CsI reacted with SO2 to form a red solution. Removal of the excess of unreacted SO<sub>2</sub> at -10 °C resulted in a red-orange solid of the approximate composition CsI-4SO<sub>2</sub>, which at 25.5 °C exhibited a dissociation pressure of 450 torr. Exhaustive dissociation of the adduct at 25.5 °C produced pure SO<sub>2</sub> and CsI (6.3 mmol).

Preparation of NF4WOF5. A mixture of CsF (21.99 mmol) and NF<sub>4</sub>SbF<sub>6</sub> (22.14 mmol) was loaded in the drybox into one half and  $WOF_4$  (14.59 mmol) into the other half of a prepassivated Teflon double-U metathesis apparatus. Dry HF (16 mL, liquid) was added on the vacuum line to the half containing the NF<sub>4</sub>SbF<sub>6</sub>-CsF mixture and was warmed to 25 °C for 30 min with stirring. After this mixture was cooled to -78 °C, the metathesis apparatus was inverted and the  $\mathbf{NF_4HF_2}$  solution was filtered into the other half of the apparatus containing the WOF<sub>4</sub>. The NF<sub>4</sub>HF<sub>2</sub>-WOF<sub>4</sub> mixture in HF was stirred for 30 min at 25 °C, and then the volatile material was pumped off for 12 h at 25 °C and 0.8 h at 50 °C. The volatile material, trapped at -196 °C, was shown by infrared spectroscopy to consist of HF and NF3 and did not contain any tungsten species. The filter cake (8.06 g; weight calculated for 21.99 mmol of  $CsSbF_6 = 8.11$  g) and the filtrate residue (4.99 g; weight calculated for 14.59 mmol of NF<sub>4</sub>WOF<sub>5</sub> = 5.61 g) were shown by vibrational spectroscopy to consist of  $CsSbF_6$ and mainly NF4WOF5, respectively. On the basis of its elemental analysis, the filtrate residue had the composition (wt %): NF<sub>4</sub>WOF<sub>5</sub>, 96.1; CsSbF<sub>6</sub>, 2.0; NF<sub>4</sub>SbF<sub>6</sub>, 1.9. Anal. Calcd: NF<sub>3</sub>, 18.14; Cs, 0.72; W, 45.91; Sb, 1.37. Found: NF<sub>3</sub>, 18.05; Cs, 0.7; W, 46.0; Sb, 1.32.

**Preparation of NF<sub>4</sub>W<sub>2</sub>O<sub>2</sub>F<sub>9</sub>.** A sample of NF<sub>4</sub>WOF<sub>5</sub> was placed into a sapphire tube equipped with a stainless steel valve. The contents of the tube were heated in a dynamic vacuum to 155 °C for 6.5 h and to 180 °C for 4 h. On the basis of the observed weight change and the vibrational spectra, the conversion of NF4WOF5 to NF4- $W_2O_2F_9$  was essentially complete.

In an alternate method,  $NF_4WOF_5$  was dissolved in anhydrous HF and formed a precipitate. This precipitate was separated from the solution by filtration and, on the basis of its vibrational spectra, consisted of NF4W2O2F9.

## **Results and Discussion**

Syntheses and Reaction Chemistry of CsWOF<sub>5</sub>, CsW<sub>2</sub>O<sub>2</sub>F<sub>9</sub>,  $Cs_2WO_2F_4$ ,  $CsWF_6$ ,  $WOF_4$ , and  $H_3OWOF_5$ . The synthesis of  $CsWOF_5$  was first reported<sup>19</sup> in 1958 by Hargreaves and Peacock and involved the reaction of moist CsF with  $WF_6$  in IF<sub>5</sub> solution. In subsequent reports,<sup>10,20</sup> either the original or slightly modified methods were used replacing the IF<sub>5</sub> solvent by CH<sub>3</sub>CN and using WOF<sub>4</sub> as a starting material. Since in Hargreave and Peacock's method an excess of water can result in  $WO_2F_4^{2-}$  formation, a better control of the amount of water used in the reaction was desirable. Furthermore, the intended reactions with NF4<sup>+</sup> salts required the use of anhydrous HF as a solvent. Consequently, our CsWOF<sub>5</sub> preparation was carried out in such a manner that dry CsF, WF<sub>6</sub>, and a stoichiometric amount of water were used in anhydrous HF solution. On the basis of its vibrational spectra, the resulting solid product consisted of CsWOF5<sup>10,11</sup> containing small amounts of  $CsW_2O_2F_9^{12}$  and  $CsHF_2$ . During an attempt to determine the solubility of CsWOF<sub>5</sub> in anhydrous HF, it was found that the salt did not completely dissolve in anhydrous HF. The insoluble material was filtered off and identified by vibrational spectroscopy as  $CsW_2O_2F_9$ .<sup>12</sup> The residue isolated from the filtrate after solvent removal consisted mainly of CsHF<sub>2</sub> contaminated by a small amount of cis-Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>.<sup>13</sup> The latter was most likely produced by hydrolysis of some CsWOF<sub>5</sub> by a trace of water present in the large amount of HF used as a solvent. These results demonstrate the existence of the equilibrium reaction in anhydrous HF shown by eq 1.

$$2CsWOF_5 + HF \rightleftharpoons CsW_2O_2F_9 \downarrow + CsHF_2 \qquad (1)$$

A large excess of HF and the low solubility of  $C_{s}W_{2}O_{2}F_{9}$  can

shift this equilibrium all the way to the right. Separation of the  $CsW_2O_2F_9$  precipitate from the  $CsHF_2$  solution, followed by washing with anhydrous HF, affords essentially pure  $C_{S}W_{2}O_{2}F_{9}$  and thus offers a convenient synthetic method for the preparation of this compound. Equilibrium 1 is in excellent agreement with that observed by Bougon and co-workers<sup>12</sup> for the HF-WOF<sub>4</sub> system (eq 2). If the CsHF<sub>2</sub> and CsW<sub>2</sub>O<sub>2</sub>F<sub>9</sub>

$$2WOF_4 + 2HF \rightleftharpoons W_2O_2F_9^- + H_2F^+$$
(2)

are not separated prior to HF removal, equilibrium 1 is shifted back to the left on HF removal and in our experiments resulted in a mixture containing about 95 mol % CsWOF<sub>5</sub> and 5 mol %  $CsW_2O_2F_9$ .

The above described synthesis of CsWOF<sub>5</sub> by controlled hydrolysis of  $WF_6$  in a CsF-containing HF solution is similar to the previously reported<sup>8</sup> hydrolysis of WF<sub>6</sub> in HF according to eq 3. However, on the basis of the results of this study,

$$SiO_{2} + 4HF \rightarrow SiF_{4} + 2H_{2}O$$

$$H_{2}O + WF_{6} \xrightarrow{HF} WOF_{4} + 2HF$$
(3)

care must be taken to use an excess of  $WF_6$  in this reaction. In the presence of an excess of water,  $H_3O^+WOF_5^-$  and  $H_3O^+W_2O_2F_9^-$  are formed according to eq 4 and 5. Since

$$2H_2O + WF_6 \xrightarrow{HF} H_3O^+WOF_5^- + HF$$
 (4)

$$3H_2O + 2WF_6 \xrightarrow{HF} H_3O^+W_2O_2F_9^- + 3HF$$
 (5)

these oxonium salts are also volatile, they are difficult to separate from  $WOF_4$  by sublimation. It should be noted that in the previous report<sup>21</sup> on the synthesis of  $H_3O^+WOF_5^-$  the observed anion spectra were incorrectly ascribed to a distorted WOF<sub>5</sub><sup>-</sup> anion. The reported spectra<sup>21</sup> are identical with those published<sup>12</sup> for  $W_2O_2F_9^-$ , which were also confirmed by the results of this study (see below).

For the synthesis of  $Cs_2WO_2F_4$ , the use of an HF-soluble tungsten species<sup>22</sup> and of an aqueous HF solution are important. In agreement with a previous report,<sup>22</sup> WO<sub>3</sub> or  $H_2WO_4^{23}$  was found difficult to dissolve completely in aqueous HF, but WF<sub>6</sub> and CsF in a 1:2 mole ratio were found to be a convenient alternate to  $Cs_2WO_4^{22}$  as a starting material for the preparation of pure *cis*-Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub>. If anhydrous HF is used in place of aqueous HF as a solvent, equilibria 6 and 1

$$Cs_2WO_2F_4 + 3HF \rightleftharpoons CsWOF_5 + CsHF_2 + H_2O$$
 (6)

were found to become important. Equilibrium 1 results in a precipitation of  $C_{s}W_{2}O_{2}F_{9}$ , which can be isolated by filtration, while the residue obtained from the filtrate consists of Cs<sub>2</sub>W- $O_2F_4$  and  $CsHF_2$ . The vibrational spectra observed for  $\bar Cs_2\text{-}WO_2F_4$  agree well with those^{13,23,24} previously reported and confirm that  $WO_2F_4^{2-}$  is present as the cis isomer.

For the synthesis of  $CsWF_6$ , the original method<sup>25</sup> based on the reaction given by eq 7 did not lead to a pure product,

$$2C_{sI} + 2WF_{6} \xrightarrow{SO_{2}} 2C_{s}WF_{6} + I_{2}$$
(7)

in agreement with previous reports.<sup>16,26</sup> Even when the reaction was carried out in a well-passivated sapphire reactor with carefully dried reagents, the product always contained

- (23) Mattes, R.; Müller, G.; Becher, H. J. Z. Anorg. Chem. 1972, 389, 177.
  (24) Beuter, A.; Sawodny, W. A. Z. Anorg. Allg. Chem. 1971, 381, 1.
  (25) Hargreaves, G. B.; Peacock, R. D. J. Chem. Soc. 1957, 4212.
  (26) Bond, A. M.; Irvine, I.; O'Donnel, T. A. Inorg. Chem. 1975, 14, 2408.

<sup>(21)</sup> Selig, H.; Sunder, W. A.; Schilling, F. C.; Falconer, W. E. J. Fluorine Chem. 1978, 11, 629.

<sup>(22)</sup> Pausewang, G. Z. Naturforsch. B: Anorg. Chem., Org. Chem. 1974, 29B. 49.



**Figure 1.** Vibrational spectra of solid CsWF<sub>6</sub>: trace A, infrared spectrum of the dry powder pressed between AgCl disks; trace B, Nujol mull between CsI windows; trace C, Nujol alone between CsI windows; trace D, Raman spectrum of the solid. The Raman band at 488 cm<sup>-1</sup>, marked by an asterisk, is probably due to the Pyrex capillary.

significant amounts of CsWOF<sub>5</sub>, and sometimes CsW<sub>2</sub>O<sub>2</sub>F<sub>9</sub>, as byproducts. The replacement of SO<sub>2</sub> as a solvent by HF eliminated this problem and offered the additional advantage of easy separation of the HF-insoluble I<sub>2</sub> from HF-soluble CsWF<sub>6</sub>. This preparation of the CsWF<sub>6</sub> is similar to that<sup>26</sup> previously published for NaWF<sub>6</sub> with HF as a solvent, NaF, and cadmium in place of iodide as a reducing agent. Other solvents that have successfully been used for the syntheses of WF<sub>6</sub><sup>-</sup> salts include CH<sub>3</sub>CN,<sup>16,17</sup> CH<sub>3</sub>NO<sub>2</sub>,<sup>17</sup> and SF<sub>4</sub>.<sup>27</sup>

Since only incomplete spectroscopic data had previously been reported for CsWF<sub>6</sub>, its vibrational and <sup>19</sup>F NMR spectra were recorded. The vibrational spectra and observed frequencies are given in Figure 1. The observed spectra, particularly the infrared spectra, are too complicated for an ideal octahedral anion. It must be kept in mind, however, that CsWF<sub>6</sub> samples usually are poor Raman scatterers and also hydrolyze very easily to CsWOF<sub>5</sub>. The <sup>19</sup>F NMR spectrum of CsWF<sub>6</sub> in HF solution at -80 °C consisted of a singlet at  $\phi$  159 with J<sub>183WF</sub> of about 53 Hz.

The compatibility of SO<sub>2</sub> with CsI and WF<sub>6</sub> was briefly studied in an effort to explain the above described side reaction resulting in CsWOF<sub>5</sub> formation. In agreement with a previous report,<sup>28</sup> CsI was found to form with SO<sub>2</sub> an orange-red 1:4 adduct according to eq 8. This adduct was unstable at room

$$CsI + 4SO_2 \xrightarrow{SO_2} CsI(SO_2)_4$$
 (8)

temperature, possessing a dissociation pressure of about 450 torr at 25.5 °C. Its decomposition resulted in SO<sub>2</sub> and CsI as the only products. For the  $WF_6$ -SO<sub>2</sub> system, little interaction was noted. Even after an equimolar mixture of  $WF_6$  and SO<sub>2</sub> was kept at 25 °C for 5 days, less than 0.2% of the starting material had reacted to form WOF<sub>4</sub> and SOF<sub>2</sub> according to eq 9. It therefore appears that the interaction of

$$WF_6 + SO_2 \rightarrow SOF_2 + WOF_4$$
 (9)

ences cited therein.

 $WF_6$  with SO<sub>2</sub> is either catalyzed by CsI or one of its reaction products or that the desired CsWF<sub>6</sub> product is more reactive than WF<sub>6</sub> toward SO<sub>2</sub>. Such behavior is in accord with a previous report<sup>29</sup> on the enhanced reactivity of SO<sub>2</sub> toward WF<sub>6</sub> in the presence of bases capable of forming adducts with WF<sub>6</sub>.

Syntheses of NF<sub>4</sub><sup>+</sup> Fluorotungstate Salts. The possibility was explored to synthesize stable NF<sub>4</sub><sup>+</sup> salts derived from one of the following complex fluorotungstate anions: WF<sub>6</sub><sup>-</sup>, WO<sub>2</sub>F<sub>4</sub><sup>2-</sup>, WOF<sub>5</sub><sup>-</sup>, and W<sub>2</sub>O<sub>2</sub>F<sub>9</sub><sup>-</sup>.

For the synthesis of  $NF_4WF_6$  the metathetical reaction eq 10 proved unsatisfactory because of product separation

$$NF_{4}SbF_{6} + C_{8}WF_{6} \xrightarrow{HF} C_{8}SbF_{6}\downarrow + NF_{4}WF_{6}$$
(10)

problems due to the limited solubility of the  $WF_6^-$  salts in HF especially at low temperature and because of the competing redox reaction (eq 11) during removal. Although the solubility

$$NF_4^+ + 2WF_6^- + HF \xrightarrow{HF} NF_3 + 2WF_6 + HF_2^-$$
 (11)

problem was successfully circumvented by separately generating HF solutions of  $NF_4HF_2$  and  $WF_5$  (eq 12 and 13) and

$$NF_4BiF_6 + C_8HF_2 \xrightarrow{HF} C_8BiF_6 \downarrow + NF_4HF_2$$
 (12)

$$CsWF_6 + BiF_5 \xrightarrow{HF} CsBiF_6 \downarrow + WF_5$$
(13)

combining them, the redox reaction (12) remained a major obstacle, particularly when an excess of  $NF_4HF_2$  was employed in the reaction.

....

The synthesis of  $(NF_4)_2WO_2F_4$  was attempted with use of the metathetical reaction given by eq 14. Although  $WO_2F_4^{2-1}$ 

$$2NF_4SbF_6 + Cs_2WO_2F_4 \xrightarrow{Hr} (NF_4)_2WO_2F_4 + 2CsSbF_6 \downarrow$$
(14)

appeared to be compatible with NF<sub>4</sub><sup>+</sup>, the desired (NF<sub>4</sub>)<sub>2</sub>W-O<sub>2</sub>F<sub>4</sub> could not be isolated due to WO<sub>2</sub>F<sub>4</sub><sup>2-</sup> interacting with anhydrous HF according to reactions 6 and 1. Consequently, the filter cake contained a significant amount of W<sub>2</sub>O<sub>2</sub>F<sub>9</sub><sup>-</sup> salts, whereas the filtrate residue contained a significant amount of WOF<sub>5</sub><sup>-</sup> salts. The volatile reaction products contained some NF<sub>3</sub> and OF<sub>2</sub>. Nitrogen trifluoride is the product expected for either thermal decomposition of NF<sub>4</sub>HF<sub>2</sub><sup>7</sup> or redox reactions involving NF<sub>4</sub><sup>+</sup>. Oxygen difluoride could have been formed by fluorination of H<sub>2</sub>O generated from WO<sub>2</sub>F<sub>4</sub><sup>2-</sup> and HF according to reaction 6.

Since the products from the NF<sub>4</sub>SbF<sub>6</sub> + Cs<sub>2</sub>WO<sub>2</sub>F<sub>4</sub> metathetical reactions had indicated that NF<sub>4</sub>WOF<sub>5</sub> and NF<sub>4</sub>-W<sub>2</sub>O<sub>2</sub>F<sub>9</sub> might be stable but are difficult to separate from the CsSbF<sub>6</sub> precipitate, this separation problem was overcome by first preparing an NF<sub>4</sub>HF<sub>2</sub> solution according to eq 15, then

$$NF_4SbF_6 + C_8HF_2 \xrightarrow{HF} C_8SbF_6\downarrow + NF_4HF_2$$
 (15)

separating the CsSbF<sub>6</sub> precipitate from the highly soluble  $NF_4HF_2$  by filtration, and reacting a twofold excess of this  $NF_4HF_2$  solution with WOF<sub>4</sub> according to eq 16. The excess

$$NF_4HF_2 + WOF_4 \xrightarrow{Hr} NF_4WOF_5 + HF$$
 (16)

of unreacted  $NF_4HF_2$  was decomposed at room temperature to  $NF_3$ ,  $F_2$ , and HF, leaving behind the thermally more stable  $NF_4WOF_5$  salt.

The NF<sub>4</sub>W<sub>2</sub>O<sub>2</sub>F<sub>9</sub> salt was obtained by controlled vacuum pyrolysis of NF<sub>4</sub>WOF<sub>5</sub> at 180 °C in a sapphire reactor. The vibrational spectra of the solid residue showed only the bands characteristic for NF<sub>4</sub><sup>+ 1,7,9</sup> and W<sub>2</sub>O<sub>2</sub>F<sub>9</sub><sup>-.12</sup> The volatile

<sup>(27)</sup> Kemmit, R. D. W.; Sharp, D. W. A. J. Chem. Soc. 1963, 2496.
(28) Burow, D. F. In "The Chemistry of Nonaqueous Solvents"; Lagowski, J. J., Ed.; Academic Press: New York, 1970; Vol. 3, p 164 and refer-

<sup>(29)</sup> Tebbe, F. N.; Muetterties, E. L. Inorg. Chem. 1968, 7, 172.

Table I. Vibrational Spectra of Solid  $NF_4W_2O_2F_9$ ,  $CsW_2O_2F_9$ , and  $NF_4WOF_5$ 

obsd freq, $cm^{-1}$ , and rel intens <sup>a</sup>						assignment (point group) <sup>b</sup>		
NF <sub>4</sub> W <sub>2</sub> O <sub>2</sub> F <sub>9</sub>		CsW2O2F,		NF <sub>4</sub> WOF <sub>5</sub>		NF +		WOF
IR	Raman	IR	Raman	IR	Raman	$(T_d)$	W <sub>2</sub> O <sub>2</sub> F <sub>9</sub> <sup>-</sup>	$(C_{4v})$
2320 vw				2315 vw		$2 \nu_3 (A_1 + E + F_2)$		
2060 vw		2060 vw					(2 × 1035)	
2015 w				1996 w		$v_1 + v_3 (F_2)$		
1768 vw				1760 wv		$\nu_3 + \nu_4 (A_1 + E + F_2)$		
1461 w				1453 w		$\nu_1 + \nu_4 (F_2)$		
1343 w		1338 w					(709 + 640)	
1240 sh		1 <b>24</b> 0 w					(640 + 610)	
1224 mw				1221 mw		$2 \nu_4 (A_1 + E + F_2)$		
1165 vs	1165 (0.2)			1160 vs	1165 (0.7)	$\nu_3 (F_2)$		
1035 vs	1034 (10)	{1048 vs {1035 vs	1036 (10)	991 vs	996 (10)		$\nu$ W=O out of phase $\nu$ W=O in phase	$\nu_1\left(\mathbf{A}_1\right)$
900 vw	850 (1.5)				852 (8.4)	$\nu_1$ (A <sub>1</sub> )	•	
		822 vw 790 vw						
709 s	698 (5.4)	704 s	700 (6.3)	688 vs	690 (5.4)		$\nu_{s}$ WF <sub>4</sub> in phase	$\nu_{2}(A_{1})$
640 vs, br		628 vs, br	. ,	620 vs, br	. ,		vas WF	ν. (E)
610 s	610 (0. <b>9</b> )		610 (0.6)		613 (4.9)	$\nu_{4}(F_{2})$	$\nu_{s}$ WF <sub>4</sub> out of phase	$\nu_{5}$ (B <sub>1</sub> )
				515 vs			•	$\nu_{3}(A_{1})$
462 s) 440 s)	440 (0.4)	440 vs			446 (1.6)	$\nu_2$ (E)	$\nu_{as}$ WFW	• • •
400 vw		400 vw						
	335 sh		340 (0.6)	١				
	316 (6.2)		320 (5.3)		329 (6.8)			$\nu_{\rm o}$ (E)
	305 sh		309 sh					
	283 (0.5)		284 (0.2)	<b>}</b>	285 (0.5)		def vibrations	$\nu_{4}$ (A <sub>1</sub> )
	211 (0.6)		217 (0.5)					
	194 (0.2)		200 (0.2)	1				
	120 sh		122 (0.3)	/	140 sh			

<sup>a</sup> Uncorrected Raman intensities. <sup>b</sup> The actual site symmetry of NF<sub>4</sub><sup>+</sup> and WOF<sub>5</sub><sup>-</sup> might be lower than  $T_d$  and  $C_{4v}$ , respectively. <sup>c</sup> Assignments from ref 10 and 12.



Figure 2. Infrared and Raman spectra of solid  $NF_4WOF_5$ . The broken line indicates absorption due to the AgCl window material.

products consisted of NF<sub>3</sub>, OF<sub>2</sub>, and WF<sub>6</sub> in a mole ratio of about 2:1:1, indicating the reaction shown in eq 17.

$$3NF_4WOF_5 \rightarrow 2NF_3 + OF_2 + WF_6 + NF_4W_2O_2F_9$$
 (17)

**Properties of NF**<sub>4</sub>WOF<sub>5</sub> and NF<sub>4</sub>W<sub>2</sub>O<sub>2</sub>F<sub>9</sub>. Both salts are white crystallinic solids stable at room temperature. The thermal stability of the salts was studied in a static vacuum with a sapphire reactor equipped with a pressure transducer. The NF<sub>4</sub>WOF<sub>5</sub> salt was found to be stable at 55 °C but at 85 °C started to undergo slow decomposition according to reaction 17 at an approximate rate of 0.03% h<sup>-1</sup>. At 155 °C the pyrolysis proceeded at a much faster rate, and after 6.5



Figure 3. Infrared and Raman spectra of solid  $NF_4W_2O_2F_9$  compared to those of solid  $CsW_2O_2F_9$ .

h about 90% of the starting material was converted to NF<sub>4</sub>- $W_2O_2F_9$  and to the other products shown in reaction 17. Complete conversion to NF4W2O2F9 was achieved by heating to 180 °C for an additional 4 h.

The ionic structure of NF<sub>4</sub>WOF<sub>5</sub> and NF<sub>4</sub>W<sub>2</sub>O<sub>2</sub>F<sub>9</sub> in both the solid state and HF solution was demonstrated by vibrational and <sup>19</sup>F NMR spectroscopy. The infrared and Raman spectra of solid NF<sub>4</sub>WOF<sub>5</sub> and NF<sub>4</sub>W<sub>2</sub>O<sub>2</sub>F<sub>9</sub> are shown in Figures 2 and 3, respectively. For comparison, the vibrational spectra of  $CsW_2O_2F_9$  are also shown in Figure 3. The observed frequencies, together with their assignments to the  $NF_4^{+,1,7,9}$  $WOF_5^{-,10,11}$  and  $W_2O_2F_9^{-12}$  ions, are summarized in Table I and clearly establish the nature of these salts in the solid state. In HF solution, however, a significant change was observed. First of all, both NF4WOF5 and NF4W2O2F9 were only partially soluble in anhydrous HF. In both cases the insoluble phases were shown by Raman spectroscopy to consist of  $NF_4W_2O_2F_9$ , and the HF solution showed only  $NF_4^+$  and  $W_2O_2F_9^-$  ions. In the HF solution obtained from dissolving  $NF_4WOF_5$ , the ratio of  $NF_4^+:W_2O_2F_9^-$  was, as expected from eq 18, considerably higher than that found for the  $NF_4W_2O_2F_9$ 

$$2NF_4WO_5 + HF \stackrel{HF}{\longleftrightarrow} NF_4HF_2 + NF_4W_2O_2F_9$$
 (18)

solution. The presence of these ions and the lack of  $WOF_5^$ in these solutions were confirmed by <sup>19</sup>F NMR spectroscopy. These spectra showed a triplet of equal intensity at  $\phi$  215.5 with  $J_{\rm NF}$  = 239.5 Hz, characteristic for NF<sub>4</sub><sup>+,1,7</sup> and a singlet at  $\phi$  54 with two <sup>183</sup>W satellites with  $J_{183}_{WF}$  = 84 Hz, characteristic for  $W_2O_2F_9^-$  in HF solution.<sup>12</sup> The latter signal has been attributed to the eight equivalent fluorines of  $W_2O_2F_9^$ with the bridging ninth fluorine undergoing rapid exchange with the HF solvent.<sup>12</sup> This exchange was still rapid at -60°C, and no evidence for F-F coupling was observed. These Raman and NMR data suggest that NF4WOF5 undergoes the solvolysis in anhydrous HF given by eq 18. This solvolysis is analogous to that observed for CsWOF<sub>5</sub> in reaction 1 and explains why NF<sub>4</sub>WOF<sub>5</sub> could not be prepared by a direct metathesis. Due to their limited solubility in anhydrous HF, some of the  $W_2O_2F_9^-$  salt is precipitated out. This was confirmed by <sup>19</sup>F NMR peak area measurements on the HF solution of the original NF4WOF5 sample; it was shown that the ratio of the fluorines on nitrogen to the equivalent fluorines on tungsten was 1.5:1, indicating that on the basis of (18) about half of the  $NF_4W_2O_2F_9$  had precipitated out under the given conditions.

Conclusion. The perfluoroammonium cation is capable of forming stable salts derived from the WOF<sub>5</sub><sup>-</sup> and the  $W_2O_2F_9^$ anion but rapidly oxidizes the  $WF_6^-$  anion to  $WF_6$ . The new  $NF_4WOF_5$  and  $NF_4W_2O_2F_9$  salts were characterized, and the syntheses and reaction chemistry of tungsten fluoride and oxyfluoride anions were studied in more detail.

Acknowledgment. We thank Drs. C. J. Schack and L. R. Grant and Mr. R. D. Wilson for helpful support and the Office of Naval Research and the Army Research Office for financial support of this work.

Registry No. NF4WOF5, 79028-46-9; NF4W2O2F9, 79056-35-2;  $CsWOF_5$ , 58341-09-6;  $CsW_2O_2F_9$ , 79056-36-3;  $cis-Cs_2WO_2F_4$ , 79005-35-9; CsWF<sub>6</sub>, 19175-38-3; WF<sub>6</sub>, 7783-82-6; WOF<sub>4</sub>, 52049-91-9.

Contribution from the Department of Chemistry, College of Liberal Arts and Sciences, and the Department of Industrial Chemistry, Faculty of Engineering, Kyoto University, Sakyo-ku, Kyoto 606, Japan

# Formation Constant of the Tetracyanocuprate(II) Ion and the Mechanism of Its Decomposition

AKIRA KATAGIRI,\*1a SEIJI YOSHIMURA,1b and SHIRO YOSHIZAWA1b

Received May 29, 1981

The overall formation constant ( $\beta$ ) of tetracyanocuprate(II) ion (Cu<sup>II</sup>(CN)<sub>4</sub><sup>2-</sup>), which is formed as a reaction intermediate in the anodic oxidation of cyanide ion in the presence of copper ion, has been obtained by measuring the standard electrode potential ( $E^{\circ}$ ) of the redox couple  $Cu^{II}(CN)_4^{2-}/Cu^{I}(CN)_4^{3-}$ . The results are  $E^{\circ}(Cu^{II}(CN)_4^{2-}/Cu^{I}(CN)_4^{3-}) = 0.66$  V vs. NHE and  $\beta(Cu^{II}(CN)_4^{2-}) = 10^{22.0}$  M<sup>-4</sup> at 25 °C. The mechanism of the decomposition of the  $Cu^{II}(CN)_4^{2-}$  is discussed by using the  $\beta(Cu^{II}(CN)_4^{2-})$  value and the previously obtained kinetic data. It is concluded that a binuclear complex,  $Cu^{II}_2(CN)_6^{2-}$ , is formed from  $Cu^{II}(CN)_4^{2-}$ . This complex subsequently decomposes to form  $Cu^{II}(CN)_2^{-}$  and  $(CN)_2$ , this step being rate determining. The kinetic studies of the chemical reaction between Cu<sup>2+</sup> and CN<sup>-</sup>, which have been reported by several groups of workers, are also interpreted by the same mechanism.

#### Introduction

In previous studies of the anodic oxidation of cyanide ion in the presence of copper ion, we have found that copper ion has a marked catalytic effect on this reaction<sup>2a</sup> and that the primary reaction product is cyanogen, which gives various secondary products according to the pH of the solution.<sup>3</sup>

We have proposed the reaction mechanism given by eq 1 and 2 from some electrochemical studies.<sup>2</sup> We have further

$$Cu^{I}(CN)_{4}^{3^{-}} \longrightarrow Cu^{II}(CN)_{4}^{2^{-}} + e^{-} \qquad (1)$$

$$u^{I}(CN)_{4}^{2^{-}} + CN^{-} \longrightarrow Cu^{I}(CN)_{4}^{3^{-}} + \frac{1}{2}(CN)_{2} \qquad (2)$$

studied the kinetics of reaction 2 by means of ESR spectroscopy<sup>4,5</sup> and have obtained the rate equation given by eq 3. On the basis of this result, we have proposed two possible

<sup>(1)</sup> (a) Department of Chemistry. (b) Department of Industrial Chemistry. (2)

 <sup>(</sup>a) S. Yoshimura, A. Katagiri, and S. Yoshizawa, Denki Kagaku oyobi Kogyo Butsuri Kagaku, 47, 360 (1979);
 (b) ibid., 47, 488 (1979). (3)S. Yoshimura, A. Katagiri, and S. Yoshizawa, Nippon Kagaku Kaishi, 1327 (1980).

<sup>(4)</sup> S. Yoshimura, A. Katagiri, Yasuo Deguchi, and S. Yoshizawa, Bull.

Chem. Soc. Jpn., 53, 2434 (1980).

S. Yoshimura, A. Katagiri, Yasuo Deguchi, and S. Yoshizawa, Bull. Chem. Soc. Jpn., 53, 2437 (1980).