

Enantioselective Syntheses and Configuration Assignments of γ -Chiral Butenolides from *Plagiomnium undulatum*: Butenolide Synthesis from Tetronic Acids

Tobias Kapferer,^[a] Reinhard Brückner,^{*[a]} Axel Herzig,^[a] and Wilfried A. König^{†[b]}


Abstract: Both enantiomers of the γ -chiral α,β -dimethylated butyrolactones *nat-1* and *nat-2* from the moss *Plagiomnium undulatum* were synthesized stereoselectively through butenolides and tetronic acids, respectively. The configuration of the natural products was determined by GLC comparisons with mono(3-*O*-acetyl-6-*O*-*tert*-butyldimethylsilyl-2-*O*-methyl)hexakis(6-*O*-*tert*-butyldimethylsilyl-2,3-di-*O*-methyl)- β -cyclodextrin as a stationary phase.

Keywords: Arndt-Eistert homologation • asymmetric synthesis • butyrolactones • configuration determination • dihydroxylation • tetronic acids

Introduction

Bryophytes (mosses) are known to generate a great variety of secondary metabolites. However, this property mainly applies to liverworts (Hepaticae), whereas common mosses (Musci) are considered to be poor in their production of volatile constituents. A recent investigation of the hydrodistillation products of a selection of mosses revealed that many of them also produce complex mixtures of volatiles, although in much smaller amounts (approximately 10%) than liverworts. From the genus *Plagiomnium undulatum*, in addition to (+)-dauca-8,11-diene (a new sesquiterpene hydrocarbon), two Δ^2 -butenolides,^[1] *nat-1* and *nat-2* (Scheme 1), with the latter exhibiting a very intense, pleasant, floral fragrance, were isolated and their structures were derived by NMR spectroscopy and mass spectrometry analysis.^[2] However, the absolute configuration at their stereocenters remained unknown.

In order to establish the latter, we compared extracts from *Plagiomnium undulatum*, which contained both *nat-1*



Scheme 1. Butenolides from *Plagiomnium undulatum*.^[2]

and *nat-2* (plus other compounds), with synthetic specimens of (*S*)-**1**, (*R*)-**1**, (*S*)-**2**, and (*R*)-**2** by enantioselective gas chromatography with appropriately modified cyclodextrins.^[3] This was the only way of determining the absolute configuration of compound *nat-1* because it had been inaccessible in pure form from the natural source. In contrast, GLC comparisons between *nat-2* (which had been isolated in pure form) and stereochemically unambiguously assigned reference compounds (*S*)-**2** and (*R*)-**2** were an obvious means of absolute configuration assignment, since “easier” methods like CD spectroscopy^[4] or ¹H NMR spectroscopy analysis in the presence of an enantiopure lanthanide shift reagent^[5] were not readily applicable.

Reference compounds (*S*)-**1**, (*R*)-**1**, (*S*)-**2**, and (*R*)-**2** qualified, in principle, for being accessible by the asymmetric dihydroxylation^[6] (AD) of sterically homogeneous β,γ -unsaturated carboxylic esters, since this reaction provides β -hydroxy- γ -butyrolactones in a single step and in essentially enantiopure form.^[7] Having accessed a variety of saturated^[8] and unsaturated butyrolactones^[9] in this manner,^[7,10] we found the latter did not include trisubstituted (for example, **1**) or tetrasubstituted (for example, **2**) Δ^2 -butenolides. This gap is closed by the present study. Concomitantly, we disclose the first examples each of a novel access to optically active tetronic acids (**11**→**10**→**18**; **5**→**4**→**21**) and of a novel preparation of β -substituted butenolides (**18**→**19**→**2**).


[a] Dipl.-Chem. T. Kapferer, Prof. Dr. R. Brückner, Dr. A. Herzig
Institut für Organische Chemie und Biochemie
Albert-Ludwigs-Universität
Albertstrasse 21, 79104 Freiburg (Germany)
Fax: (+49) 40-4283-82893
E-mail: reinhard.brueckner@organik.chemie.uni-freiburg.de

[b] Prof. Dr. W. A. König[†]
Institut für Organische Chemie
Universität Hamburg
Martin-Luther-King-Platz 6, 20146 Hamburg (Germany)
Fax: (+49) 761-203-6100
E-mail: wkoenig@chemie.uni-hamburg.de

[†] Deceased, November 19, 2004.

Results and Discussion


Following the β,γ -unsaturated ester $\rightarrow$ β -hydroxy- γ -lactone strategy, the two enantiomers of trisubstituted butenolides (*S*)- and (*R*)-**1** were traced back to ester *trans*-**5**^[11] (Scheme 2). This originated from the deconjugative Knoeven-




Scheme 2. Approaching butenolides (*S*)- and (*R*)-**1**. a) Malonic acid (1.1 equiv), NEt_3 (3.0 equiv), 90°C , 18 h; b) H_2SO_4 (cat.), MeOH, Δ , 2 h; 82%; c) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQD})_2\text{PHAL}$ (0.01 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 12 h; 91%; d) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQD})_2\text{PHAL}$ (0.01 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 12 h; 90%; e) $p\text{BrC}_6\text{H}_4\text{CO}_2\text{H}$ (1.1 equiv), DCC (1.1 equiv), DMAP (cat.), CH_2Cl_2 , RT, 4 h, Δ , 1 h; 81%. A Pluton/Povray plot^[15] of the solid-state structure of (*S,S*)-**6** is also depicted. Pent = pentyl, $(\text{DHQD})_2\text{PHAL}$ = 1,4-bis(dihydroquinidyl)phthalazine, DCC = *N,N'*-dicyclohexylcarbodiimide, DMAP = 4-dimethylaminopyridine.

nagel condensation^[12] of aldehyde **3** with malonic acid followed by methyl esterification. Oxidation of ester *trans*-**5** with AD mixes α and β (containing $(\text{DHQ})_2\text{PHAL}$ and $(\text{DHQD})_2\text{PHAL}$, respectively, as the chiral auxiliary) provided hydroxylactones (*S,S*)-**4** (94% *ee*;^[13] absolute configuration ascertained by X-ray crystal structure analysis^[14] of a crystal of the *para*-bromobenzoate (*S,S*)-**6**) and (*R,R*)-**4** (97% *ee*^[13]), respectively.

Application of our unsaturated ester $\rightarrow$ β -hydroxy- γ -lactone strategy to obtain the optically active tetrasubstituted butenolides (*S*)- and (*R*)-**2** allowed us to start both from ester (*E*)-**11**^[16] (Scheme 3) or from its isomer (*Z*)-**11** (Scheme 4). AD mix α converted the former into a hydroxylactone, (*4S,5S*)-**10**, with 91% *ee*,^[13] and the latter into the hydroxylactone (*4S,5R*)-**10** with 80% *ee*.^[13] Likewise, dihydroxylation of esters (*E*)- and (*Z*)-**11** with AD mix β occurred with 97% *ee*^[13] ($\rightarrow$ (*4R,5R*)-**10**) and 85% *ee*^[13] ($\rightarrow$



Scheme 3. Approaching butenolides (*S*)- and (*R*)-**2**, variant 1. a) SOCl_2 (1.0 equiv), DMF (cat.), CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{RT}$, 3 h; b) CH_2N_2 (2.0 equiv), Hünig's base (1.0 equiv), Et_2O , $0^\circ\text{C} \rightarrow \text{RT}$, 1 h; 55% over 2 steps; c) AgOBz (0.3 equiv), NEt_3 (4.7 equiv), MeOH, RT, 5 h; 88%; d) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQ})_2\text{PHAL}$ (0.016 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 1 d; 83%; e) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQD})_2\text{PHAL}$ (0.016 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 1 d; 82%. DMF = *N,N*-dimethylformamide.




Scheme 4. Approaching butenolides (*S*)- and (*R*)-**2**, variant 2. a) SOCl_2 (1.1 equiv), DMF (cat.), CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{RT}$, 2.5 h; b) CH_2N_2 (3.0 equiv), Hünig's base (1.0 equiv), Et_2O , $0^\circ\text{C} \rightarrow \text{RT}$; 57% over 2 steps; c) AgOBz (0.3 equiv), NEt_3 (4.7 equiv), MeOH, RT, 4 h; 95%; d) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQD})_2\text{PHAL}$ (0.016 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 1 d; 85%; e) $\text{K}_2\text{OsO}_2(\text{OH})_4$ (0.8 mol %), $(\text{DHQ})_2\text{PHAL}$ (0.016 equiv), $\text{K}_3\text{Fe}(\text{CN})_6$ (3.0 equiv), K_2CO_3 (3.0 equiv), MeSO_2NH_2 (1.0 equiv), $t\text{BuOH}/\text{H}_2\text{O}$ (1:1), 0°C , 1 d; 92%; f) $p\text{BrC}_6\text{H}_4\text{COCl}$ (1.2 equiv), NEt_3 (1.3 equiv), DMAP (cat.), CH_2Cl_2 , RT, 6 h, Δ , 2 h; 56%. A Pluton/Povray plot^[15] of the solid-state structure of (*4S,5R*)-**12** is also depicted.

(*4R,5S*)-**10**), respectively. The β,γ -unsaturated C_{10} esters (*E*)- and (*Z*)-**11** had been obtained stereospecifically from the α,β -unsaturated C_9 acids (*E*)-**7**^[17] and (*Z*)-**7**,^[18] respectively by Arndt–Eistert homologations^[19] proceeding through the

corresponding acid chlorides (*E*)- and (*Z*)-**8** and the derived diazoketones (*E*)- and (*Z*)-**9**.^[20]

The anomalous X-ray diffraction^[14] of the *para*-bromobenzoate first obtained from these lactones as nice crystals revealed its stereostructure to be that depicted for (4*S*,5*R*)-**12**; this proves that the configuration of the underlying hydroxylactone, namely the α -osmylation product of ester (*Z*)-**11**, is (4*S*,5*R*)-**10** (Scheme 4). The configurations of the hydroxylactone isomers obtained according to Scheme 3 emerged from correlations with those shown in Scheme 4. According to chiral GLC,^[13] α -methylation/oxidation of the α -osmylation product (4*S*,5*S*)-**10** of ester (*E*)-**11** (Scheme 3) provided the *same* tetronic acid enantiomer ((*S*)-**18**, Scheme 6) as α -methylation/oxidation of compound (4*R*,5*S*)-**10** from Scheme 4. Similarly, α -methylation/oxidation of the β -osmylation product of ester (*E*)-**11** (Scheme 3)—which was hence designated (4*R*,5*R*)-**10**—gave the identical tetronic acid enantiomer (not depicted) to that obtained by α -methylation/oxidation of hydroxylactone (4*S*,5*R*)-**10** (Scheme 4).

Hydroxylactones (*S*,*S*)-**4** and (*R*,*R*)-**4** were transformed




Scheme 5. Completing the synthesis of butenolides (*S*)- and (*R*)-**1**. a) MsCl (1.5 equiv), NEt₃ (2.8 equiv), CH₂Cl₂, 0°C, 30 min; 92%, 94% ee; b) CH₂N₂ (5.0 equiv), Et₂O, 0°C → RT, 15 h; 96%; c) LDA (1.7 equiv), THF, -78°C, 30 min; MeI (1.8 equiv), -78°C, 18 h; 80%; d) 1,4-dioxane, 110°C, 5 d; 75%. Ms = mesyl = methanesulfonyl, LDA = lithium diisopropylamide, THF = tetrahydrofuran.

into the trisubstituted butenolides (*S*)-**1** and (*R*)-**1**, respectively, as detailed in Scheme 5 for the former compound. Dehydration with mesyl chloride/triethylamine produced the expected Δ^1 -butenolide in 94% ee (92% yield). Its C ^{β} H=C ^{α} H moiety was elaborated into the desired C ^{β} Me=C ^{α} Me moiety by Hanessian and Murray's β -methylation/ α -alkylation protocol for butenolides.^[21] Cycloaddition of diazomethane gave the Δ^1 -pyrazoline **13** with the C ^{β} -Me

“bond-to-be” (96% yield). Enolate formation with LDA and treatment with methyl iodide afforded a Δ^2 : Δ^1 mixture of pyrazolines **14** and **15**, thereby establishing the C ^{α} -Me bond (80% yield). Pyrolysis in refluxing dioxane provided the desired butenolide (*S*)-**1** in 75% yield with an undiminished ee value (95%). A specimen of (*R*)-**1** (97% ee) was similarly obtained.

The methodology of Hanessian and Murray^[21] was unsuitable for converting hydroxylactones (4*R*,5*S*)-**10** and (4*S*,5*R*)-**10** into the tetrasubstituted butenolides (*S*)-**2** and (*R*)-**2**, respectively, since butenolide **16**—obtained by dehydrating an isomeric mixture of hydroxylactones **10**—did not react with diazomethane (Scheme 6). We circumvented this inertia by




Scheme 6. Completing the synthesis of butenolides (*S*)- and (*R*)-**2**. a) CH₂N₂ (5.0 equiv), Et₂O, RT, 5 d; b) LDA (2.8 equiv), THF, -78°C, 30 min; c) MeI (1.8 equiv), THF/DMPU, -78°C → -40°C, 14 h; 74%; d) DMSO (3.5 equiv), (F₃CCO)₂O (2.0 equiv), CH₂Cl₂, -78°C, 2 h; e) NEt₃ (4.0 equiv), -78°C, 14 h; 99%; f) MeLi (2.0 equiv), THF, -50°C → 0°C, 2 h; g) HCl, 30 min, RT; 87%, 86% ee; h) Me₃O⁺·BF₄⁻ (3.0 equiv), CH₂Cl₂, RT, 28 h; 85%. DMPU = 1,3-dimethyltetrahydro-2-(1*H*)-pyrimidinone.

adding tetronic acids to the product palette of AD reactions of β , γ -unsaturated esters and by establishing a novel way for converting the C ^{β} -OH motif of tetronic acids into the C ^{β} -hydrocarbon subunit of Δ^2 -butenolides.

The *S* enantiomer of butenolide **2** was targeted from hydroxylactone (4*R*,5*S*)-**10** by lithioalkoxide/enolate formation followed by methylation^[22] (Scheme 6). A single diastereomer of the expected α -methyl- β -hydroxylactone was isolated. Working in THF/DMPU, the α -methyl and β -hydroxy groups were probably oriented *trans*.^[10a,22] Swern oxidation^[23] including enol formation yielded tetronic acid (*S*)-**18** almost quantitatively. There appears to be little prece-

dence^[24] for such an approach to these compounds.^[25] The method could be extended to tetrone acid **21** which, unlike (*S*)-**18**, might—but did not—racemize due to the possibility of C³,H acidity (Scheme 7).



Scheme 7. Synthesis of an optically active tetronic acid, **21**, from a β,γ -unsaturated ester. a) Same as step d of Scheme 2; b) LDA (5.0 equiv), THF, -78°C , 1 h; MeI (10 equiv), THF, -78°C , 2 h; 83%; c) DMSO (3.5 equiv), $(\text{F}_3\text{CCO})_2\text{O}$ (2.0 equiv), CH_2Cl_2 , -78°C , 1 h; NEt_3 (4.0 equiv), -78°C , 45 min; 92%.

Tetronic acid (*S*)-**18** was methylated by Meerwein's salt regioselectively at $\text{C}^2=\text{O}$, rather than at C^4-OH , thereby yielding methoxyfuranone (*S*)-**19** (Scheme 6).^[26] Addition of MeLi followed by acid hydrolysis provided target structure (*S*)-**2** in 87% yield. This reaction, to the best of our knowledge, represents the first example of the overall transformation $\text{O}=\text{C}-\text{C}=\text{C}(\text{OR}^1)_2 + \text{R}^2\text{M} \rightarrow \text{R}^2-\text{C}=\text{C}-\text{C}(=\text{O})(\text{OR}^1)$ in heterocycle synthesis. It consists of the sequential transformations 1) $\text{HO}-\text{C}=\text{CH}-\text{C}(=\text{O})\text{OR}^1 + \text{R}^1\text{X} \rightarrow \text{O}=\text{C}-\text{C}=\text{C}(\text{OR}^1)_2$ and 2) $\text{O}=\text{C}-\text{C}=\text{C}(\text{OR}^1)_2 + \text{R}^2\text{M} \rightarrow \text{R}^2-\text{C}=\text{C}-\text{C}(=\text{O})(\text{OR}^1)$. The same sequence of steps might be applicable to the conversion of *other* β -ketolactones into *other* α,β -unsaturated lactones containing a β -substituent; conceivably, this could entail the two-step conversion of tetrahydropyran-2,4-diones into 4-substituted 5,6-dihydro-(2*H*)-2-pyrans or the two-step conversion of benzo-3,4-dihydro-(2*H*)-2,4-pyranones into 4-substituted coumarins. It should be noted that the *related* transformation $\text{O}=\text{C}-\text{C}=\text{C}-\text{OR}^1 + \text{R}^2\text{M} \rightarrow \text{R}^2-\text{C}=\text{C}-\text{C}=\text{O}$, which proceeds at a lower oxidation state than (*S*)-**19** $\rightarrow$ (*S*)-**2**, has been amply used for the preparation of 3-substituted 2-cycloalken-1-ones.^[27]

Having made enantiomer (*R*)-**2** available by the same sequence of steps (Scheme 6, bottom), we were in possession of all the materials needed for the GLC analysis of compounds *nat-1* and *nat-2*. As demonstrated by enantioselective GLC (Figure 1), the predominant *nat-2* is the enantiopure *S* enantiomer, while the minor component *nat-1* is a mixture of enantiomers with a small excess of the *R* enantiomer (9% *ee*).

Experimental Section

General: All reactions were performed in oven-dried (110°C) glassware under N_2 . THF was freshly distilled from K. CH_2Cl_2 was distilled from CaH_2 . Diazomethane was prepared as an ethanol-free diethyl ether solution according to the literature^[28] by using the ALDRICH Diazald kit; the diazomethane concentrations were determined prior to use.^[28] Products were purified by flash chromatography^[29] on Merck silica gel 60. Yields refer to analytically pure samples. ^1H NMR (CHCl_3 ($\delta=7.26$ ppm)




Figure 1. GLC analyses of natural plant extract and synthetic reference compounds by using a 25 m fused-silica capillary column with monokis(3-*O*-acetyl-6-*O*-*tert*-butyldimethylsilyl-2-*O*-methyl)hexakis(6-*O*-*tert*-butyldimethylsilyl-2,3-di-*O*-methyl)- β -cyclodextrin^[3a] at 150°C (isothermal). H_2 at an inlet pressure of 0.5 bar was used as the carrier gas with split injection (ratio 1:30). Bottom: natural plant extract; center: synthetic reference compounds; top: coinjection of natural plant extract and synthetic reference compounds.

as the internal standard in CDCl_3 ; C_6HD_5 ($\delta=7.15$ ppm) as the internal standard in C_6D_6) and ^{13}C NMR (CDCl_3 (center peak of the triplet $\delta=77.0$ ppm) as the internal standard in CDCl_3 ; C_6D_6 (center peak of the triplet $\delta=128.0$ ppm) as the internal standard in C_6D_6) spectroscopy was performed on Varian Mercury VX 300 and Bruker DRX 500 spectrometers. Integrals are in accordance with the assignments; coupling constants are given in Hz. The assignments of ^1H and ^{13}C NMR signals refer to the IUPAC nomenclature except within substituents where primed numbers are used. Combustion analyses were performed by E. Hickl, Institut für Organische Chemie und Biochemie, Universität Freiburg. MS analyses were obtained by Dr. J. Wörth and C. Warth, Institut für Organische Chemie und Biochemie, Universität Freiburg. IR spectra were obtained on a Perkin-Elmer Paragon 1000 apparatus. Optical rotations measured with a Perkin-Elmer polarimeter 341 at 589 nm and 20°C and were calculated according to the Drude equation ($[\alpha]_D = (\alpha_{\text{exp}} \times 100)/(c \times d)$); rotational values are the average of five measurements of α_{exp} in a given solution of the respective sample. Melting points were measured on a Dr. Tottoli apparatus (Büchi) and are uncorrected. The *ee* values were determined by chiral GC, by using a Carlo Erba Instruments HRC 5160 Mega series apparatus with a Varian CP7502 (β -cyclodextrin/dimethylpolysiloxane) column.

(*S*)-3,4-Dimethyl-5-pentyl-2-(5*H*)-furanone ((*S*)-1**):** A solution of **15** (70 mg, 0.33 mmol) in 1,4-dioxane (3 mL) was heated at 110°C for 5 d. The solvent was evaporated in vacuo. Subsequent purification by flash chromatography (cyclohexane/EtOAc 10:1) afforded the title compound (45 mg, 75%); $[\alpha]_D = -6.5$ ($c=0.6$ in CHCl_3); IR (film): $\tilde{\nu}=2955, 2930, 2860, 1755, 1685, 1465, 1460, 1440, 1385, 1330, 1130, 1115, 1095, 1060,$

presumably interpretable as d, $^4J_{\text{allyl}}=1.2$ Hz, 3-CH₃), 5.69 (incomplete resolved dq, $^4J_{2,3-\text{Me}}=^4J_{2,4}=1.2$ Hz, 2-H), 11.89 (brs, COOH) ppm; elemental analysis calcd (%) for C₉H₁₆O₂ (156.21): C 69.32, H 10.16.

(Z)-3-Methyl-2-octenoic acid ((Z)-7): At -5°C MeLi (1.6 M in diethyl ether, 57.5 mL, 92.0 mmol, 4.6 equiv) was added to a suspension of CuI (8.76 g, 46.0 mmol, 2.3 equiv) in THF (200 mL). The solution was stirred for 20 min at this temperature and then cooled to -78°C . A solution of 2-octynoic acid (2.80 g, 20.0 mmol) in THF (9 mL) was added. After stirring for 1.5 h at this temperature, the mixture was warmed to -15°C and stirred for another 5 h. The mixture was poured into cold (0°C) aq. HCl (1 M, 400 mL). After extraction with EtOAc (4×100 mL), the combined organic extracts were dried over MgSO₄ and evaporated in vacuo. The residue was purified by flash chromatography (cyclohexane/EtOAc 10:1) to afford the title compound (3.04 g, 97%): ^1H NMR (500 MHz, CDCl₃): $\delta=0.89$ (m, 8-H₃), 1.28–1.37 (m, 6-H₂, 7-H₂), 1.47 (m, 5-H₂), 1.92 (d, $^4J_{3-\text{Me},2}=1.4$ Hz, 3-CH₃), 2.63 (m, 4-H₂), 5.67 (m, 2-H), 11.89 (brs, COOH) ppm.

(E)-3-Methyl-2-octenoyl chloride ((E)-8): At 0°C , SOCl₂ (0.33 mL, 0.53 g, 4.5 mmol, 1.0 equiv) and DMF (1 drop) were added to a solution of (E)-3-methyl-2-octenoic acid (701 mg, 4.49 mmol) in CH₂Cl₂ (19 mL). The solution was stirred for 3 h at room temperature. After evaporation in vacuo, the crude residue was used for the preparation of (E)-9 without further purification.

(Z)-3-Methyl-2-octenoyl chloride ((Z)-8): At 0°C , SOCl₂ (0.051 mL, 83 mg, 0.70 mmol, 1.1 equiv) and DMF (1 drop) were added to a solution of (Z)-3-methyl-2-octenoic acid (0.10 g, 0.64 mmol) in CH₂Cl₂ (4 mL). The solution was stirred for 2.5 h at room temperature. After evaporation in vacuo, the crude residue was used for the preparation of (Z)-9 without further purification.

(E)-1-Diazo-4-methyl-3-nonen-2-one ((E)-9): At 0°C , the crude acid chloride (E)-8 was added to a solution of CH₂N₂ (0.42 M in diethyl ether, 21.4 mL, 8.98 mmol, 2.0 equiv) and Hünig's base (0.78 mL, 0.58 g, 4.5 mmol, 1.0 equiv). After the mixture had been stirred for 1 h at room temperature, the solvent was evaporated in vacuo. The residue was purified by flash chromatography (cyclohexane/EtOAc 30:1) to afford the diazoketone (E)-9 (447 mg, 55% over 2 steps) as a yellow oil: ^1H NMR (500 MHz, CDCl₃; * = distinguishable by a C,H correlation spectrum): $\delta=0.89$ (t, $J_{9,8}=7.2$ Hz, 9-H₃), 1.23–1.36 (m, 7-H₂, 8-H₂), 1.43–1.50 (m, 6-H₂), 2.10 (m, 5-H₂), 2.19 (d, $^4J_{4-\text{Me},3}=0.8$ Hz, 4-CH₃), 5.18 (brs, 1-H)*, 5.74 (brs, 3-H)* ppm; IR (film): $\tilde{\nu}=2960, 2930, 2870, 2860, 2095, 1690, 1645, 1615, 1445, 1385, 1335, 1145, 1115, 1085, 785$ cm⁻¹; HRMS: m/z calcd for C₁₀H₁₆N₂O: 180.126211; found 180.126263.

(Z)-1-Diazo-4-methyl-3-nonen-2-one ((Z)-9): At 0°C , a solution of the crude acid chloride (Z)-8 in Et₂O (3 mL) was added to a solution of CH₂N₂ (0.43 M in diethyl ether, 4.5 mL, 1.92 mmol, 3.0 equiv) and Hünig's base (0.11 mL, 84 mg, 0.65 mmol, 1.0 equiv). After the mixture had been stirred for 15 min at 0°C and for 1 h at room temperature, the solvent was evaporated in vacuo. The residue was purified by flash chromatography (petroleum ether/diethyl ether 10:1) to afford the diazoketone (Z)-9 (66 mg, 57% over 2 steps) as a yellow oil: ^1H NMR (500 MHz, CDCl₃; * = distinguishable by a C,H correlation spectrum): $\delta=0.89$ (m, 9-H₃), 1.29–1.37 (m, 7-H₂, 8-H₂), 1.43–1.51 (m, 6-H₂), 1.87 (d, $^4J_{4-\text{Me},3}=1.4$ Hz, 4-CH₃), 2.65 (brt, $J_{5,6}=7.9$ Hz, 5-H₂), 5.16 (brs, 1-H)*, 5.73 (brs, 3-H)* ppm; IR (film): $\tilde{\nu}=3080, 2960, 2930, 2870, 2860, 2095, 1645, 1615, 1445, 1385, 1335, 1145, 1115, 1085, 1005, 960, 845, 785, 725$ cm⁻¹; elemental analysis calcd (%) for C₁₀H₁₆N₂O (180.1): C 66.69, H 8.88, N 15.55; found: C 66.57, H 8.97, N 15.62.

(4R,5R)-4,5-Dihydro-4-hydroxy-5-methyl-5-pentyl-2-(3H)-furanone ((4R,5R)-10): (DHQD)₂PHAL (6 mg, 0.008 mmol, 1.6 mol %), K₃Fe(CN)₆ (494 mg, 1.50 mmol, 3.0 equiv), K₂CO₃ (207 mg, 1.50 mmol, 3.0 equiv), MeSO₂NH₂ (48 mg, 0.50 mmol, 1.0 equiv), and K₂OsO₂(OH)₄ (1.5 mg, 0.004 mmol, 0.8 mol %) were dissolved in *t*BuOH (2.5 mL) and H₂O (2.5 mL). After the solution had been cooled to 0°C , (E)-11 (92 mg, 0.50 mmol) was added. After 24 h at 0°C , aq. Na₂SO₃ (2.5 mL) was added and the mixture was stirred for 30 min at room temperature. The solution was extracted with EtOAc (3×40 mL). The combined organic extracts were dried over MgSO₄. After evaporation in vacuo, the residue

was purified by flash chromatography (cyclohexane/EtOAc 2:1) to afford the title compound (76 mg, 82%): $[\alpha]_D^{25}=+15.7$ ($c=1.2$ in CHCl₃); ^1H NMR (500 MHz, CDCl₃; * = interchangeable): $\delta=0.90$ (m, 5'-H₃), 1.29–1.47 (m, 2'-H₂, 3'-H₂, 4'-H₂), superimposed with 1.33 (s, 5-CH₃), AB signal ($\delta_A=1.73$, $\delta_B=1.81$, $J_{AB}=13.8$ Hz, A part in addition split by $J_{A,2'-H(1)}=10.1$, $J_{A,2'-H(2)}=6.1$ Hz, B part in addition split by $J_{B,2'-H(1)}=10.5$ *, $J_{B,2'-H(2)}=5.8$ * Hz, 1'-H₂), 2.46 (brs, 4-OH), 2.52 (dd, $J_{\text{gem}}=18.0$, $J_{3-H(1),4}=2.5$ Hz, 3-H¹), 2.95 (dd, $J_{\text{gem}}=18.0$, $J_{3-H(2),4}=6.2$ Hz, 3-H²), 4.21 (m, 4-H) ppm; IR (film): $\tilde{\nu}=3445, 2955, 2935, 2870, 1755, 1460, 1385, 1320, 1295, 1265, 1235, 1200, 1170, 1135, 1120, 1100, 1075, 955, 930$ cm⁻¹; $t_r(4S,5R)=39.82$ min, $t_r(4S,5S)=45.61$ min (140°C , 100 kPa); 97% *ee*; elemental analysis calcd (%) for C₁₀H₁₈O₃ (186.2): C 64.50, H 9.74; found: C 64.46, H 9.72.

(4S,5S)-4,5-Dihydro-4-hydroxy-5-methyl-5-pentyl-2-(3H)-furanone ((4S,5S)-10): The title compound (167 mg, 83%) was prepared from (E)-11 (200 mg, 1.09 mmol) by an analogous procedure to that described for (4R,5R)-10 but with (DHQ)₂PHAL as the ligand: $[\alpha]_D^{25}=-16.1$ ($c=1.1$ in CHCl₃); $t_r(4S,5S)=42.19$ min, $t_r(4R,5R)=37.79$ min (140°C , 100 kPa); 91% *ee*.

(4R,5S)-4,5-Dihydro-4-hydroxy-5-methyl-5-pentyl-2-(3H)-furanone ((4R,5S)-10): The title compound (909 mg, 85%) was prepared from (Z)-11 (1.06 g, 5.75 mmol) by an analogous procedure to that described for (4R,5R)-10 but with (DHQD)₂PHAL as the ligand: $[\alpha]_D^{25}=-4.0$ ($c=1.2$ in CHCl₃); $t_r(4R,5S)=39.50$ min, $t_r(4S,5R)=37.33$ min (140°C , 100 kPa); 85% *ee*.

(4S,5R)-4,5-Dihydro-4-hydroxy-5-methyl-5-pentyl-2-(3H)-furanone ((4S,5R)-10): The title compound (618 mg, 92%) was prepared from (Z)-11 (665 mg, 3.61 mmol) by an analogous procedure to that described for (4R,5R)-10: $[\alpha]_D^{25}=+3.8$ ($c=1.1$ in CHCl₃); ^1H NMR (500 MHz, CDCl₃): $\delta=0.89$ (t, $J_{5',4'}=7.0$ Hz, 5'-H₃), 1.25–1.45 (m, 2'-H₂, 3'-H₂, 4'-H₂), superimposed with 1.40 (s, 5-CH₃), 1.59 (m, 1'-H₂), 2.55 (dd, $J_{\text{gem}}=18.1$, $J_{3-H(1),4}=4.1$ Hz, 3-H¹), superimposes, in line with the integral, on the brs of 4-OH), 2.91 (dd, $J_{\text{gem}}=18.1$, $J_{3-H(2),4}=6.8$ Hz, 3-H²), 4.26 (m, 4-H) ppm; IR (film): $\tilde{\nu}=3445, 2955, 2935, 2870, 1755, 1385, 1320, 1295, 1265, 1195, 1175, 1135, 1120, 1105, 1065, 1040, 950$ cm⁻¹; $t_r(4S,5R)=36.79$ min, $t_r(4S,5S)=40.54$ min (140°C , 100 kPa); 80% *ee*; elemental analysis calcd (%) for C₁₀H₁₈O₃ (186.2): C 64.49, H 9.74; found: C 64.27, H 9.87.

Methyl (E)-4-methyl-3-nonen-2-one ((E)-11): Under exclusion of light, a solution of silver benzoate (130 mg, 0.568 mmol, 0.29 equiv) in triethylamine (1.28 mL, 929 mg, 9.18 mmol, 4.70 equiv) was added dropwise to a solution of (E)-9 (352 mg, 1.95 mmol) in methanol (8 mL). After the mixture had been stirred for 5 h at room temperature, the solvent was evaporated in vacuo and the residue was purified by flash chromatography (petroleum ether/diethyl ether 50:1) to afford the title compound (318 mg, 88%) as a colorless liquid: ^1H NMR (500 MHz, CDCl₃): $\delta=0.88$ (t, $J_{9,8}=7.2$ Hz, 9-H₃), 1.21–1.34 (m, 7-H₂, 8-H₂), 1.37–1.43 (m, 6-H₂), 1.62 (brs, 4-CH₃), 2.01 (brt, $J_{5,6}=7.6$ Hz, 5-H₂), 3.05 (brd, $J_{2,3}=7.1$ Hz, 2-H₂), 3.68 (s, COOCH₃), 5.31 (tm, $J_{3,2}=6.7$ Hz, 3-H) ppm; proof of the *E* configuration was obtained from the NOE observed at 3-H ($\delta=5.31$ ppm) while irradiating 5-H₂ ($\delta=2.01$ ppm).

Methyl (Z)-4-methyl-3-nonen-2-one ((Z)-11): The title compound (799 mg, 95%) was prepared from (Z)-9 (824 mg, 4.58 mmol) by an analogous procedure to that described for (E)-11: ^1H NMR (500 MHz, CDCl₃): $\delta=0.89$ (t, $J_{9,8}=7.2$ Hz, 9-H₃), 1.19–1.41 (m, 6-H₂, 7-H₂, 8-H₂), 1.73 (dt, $^4J_{4-\text{Me},3}=^5J_{4-\text{Me},2}=1.3$ Hz, 4-CH₃), 2.01 (brt, $J_{5,6}=7.7$ Hz, 5-H₂), 3.04 (m, presumably interpretable as an incomplete resolved dq, $J_{2,3}=7.2$, $^5J_{2,4-\text{Me}}=1.2$ Hz, 2-H₂), 3.68 (s, COOCH₃), 5.31 (tm, $J_{3,2}=7.2$ Hz, 3-H) ppm.

(4S,5R)-4-(4-Bromobenzyloxy)-4,5-dihydro-5-methyl-5-pentyl-2-(3H)-furanone ((4S,5R)-12): Triethylamine (0.08 mL, 0.05 g, 0.5 mmol, 1.3 equiv), *p*-bromobenzoyl chloride (0.11 g, 0.49 mmol, 1.2 equiv), and DMAP (cat.) were added to a solution of (4S,5R)-10 (70 mg, 0.41 mmol) in CH₂Cl₂ (1 mL). After the mixture had been stirred for 6 h at room temperature, the solution was heated under reflux for 2 h. It was then allowed to cool to room temperature, diluted with H₂O (15 mL), and extracted with CH₂Cl₂ (4×15 mL). The combined organic extracts were dried over MgSO₄ and evaporated in vacuo. The residue was purified by flash chromatography (cyclohexane/EtOAc 8:1) to afford the title com-

pound (85 mg, 56%): ^1H NMR (300 MHz, CDCl_3): δ = 0.91 (m , $5'\text{-H}_3$), 1.22–1.55 (m , $2'\text{-H}_2$, $3'\text{-H}_2$, $4'\text{-H}_2$), superimposed with 1.44 (s , 5-CH_3), 1.63–1.79 (m , $1'\text{-H}_2$), AB signal ($\delta_{\text{A}} = 2.69$, $\delta_{\text{B}} = 3.15$, $J_{\text{AB}} = 18.6$ Hz, A part in addition split by $J_{\text{A},4} = 2.6$ Hz, B part in addition split by $J_{\text{B},4} = 7.0$ Hz, 3-H_2), 5.48 (dd , $J_{4,3\text{-H(B)}} = 7.0$, $J_{4,3\text{-H(A)}} = 2.6$ Hz, 4-H), AA'BB' signal with signal centers at 7.62 and 7.89 ($2 \times 2\text{ArH}$) ppm; proof of the absolute configuration was obtained by X-ray crystal structure analysis.^[14]

(3aR,4S,6aS)-3,3a,4,6a-Tetrahydro-4-pentylfuro[3,4c]pyrazol-6-one (13): a) At 0°C, triethylamine (2.70 mL, 1.97 g, 19.5 mmol, 2.8 equiv) and MeSO_2Cl (0.81 mL, 1.2 g, 11 mmol, 1.5 equiv) were added to a solution of (S,S)-4 (1.20 g, 6.97 mmol) in CH_2Cl_2 (20 mL). After the mixture had been stirred for 30 min, aq. NH_4Cl (15 mL) was added. The mixture was extracted with CH_2Cl_2 (4×40 mL), dried over MgSO_4 and evaporated in vacuo. The residue was purified by flash chromatography (cyclohexane/EtOAc 6:1) to afford (S)-5-pentyl-2-(5H)-furanone (992 mg, 92%): $[\alpha]_{\text{D}} = +91.0$ ($c = 1.2$ in CHCl_3); ^1H NMR (500 MHz, CDCl_3 ; * = interchangeable): δ = 0.88–0.92 (m , $5'\text{-H}_3$), 1.29–1.35 (m , $3'\text{-H}_2$, $4'\text{-H}_2$), 1.38–1.52 (m , $2'\text{-H}_2$), 1.63–1.70 (m , $1'\text{-H}^1$), 1.73–1.80 (m , $1'\text{-H}^2$), 5.04 (dddd, $J_{5,1'} = 7.3^*$, $J_{5,1'\text{-H(2)}} = 5.4^*$, $J_{5,4} \approx J_{5,5} \approx 1.7$ Hz, 5-H), 6.10 (dd , $J_{3,4} = 5.7$, $J_{3,5} = 2.0$ Hz, 3-H), 7.45 (dd , $J_{4,3} = 5.8$, $J_{4,5} = 1.5$ Hz, 4-H) ppm; $t_{\text{r}}(\text{S}) = 34.50$ min, $t_{\text{r}}(\text{R}) = 33.18$ min (100°C, 100 kPa); 94% ee; elemental analysis calcd (%) for $\text{C}_9\text{H}_{14}\text{O}_2$ (154.2): C 70.01, H 9.15; found: C 70.01, H 9.18.

b) At 0°C, diazomethane (0.45 M in diethyl ether, 43 mL, 19.5 mmol, 5.0 equiv) was added to a solution of (S)-5-pentyl-2-(5H)-furanone (600 mg, 3.89 mmol, 94% ee) in diethyl ether (6 mL). The solution was allowed to reach room temperature and was stirred for 15 h. After evaporation in vacuo, the residue was purified by flash chromatography (cyclohexane/EtOAc 5:1) to afford the title compound (730 mg, 96%): IR (film): $\tilde{\nu} = 2960, 2935, 2865, 1770, 1355, 1225, 1205, 1185, 1025, 1000, 945, 905, 765, 755, 730, 720, 695, 655$ cm^{-1} ; elemental analysis calcd (%) for $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_2$ (196.3): C 61.20, H 8.23, N 14.27; found: C 61.27, H 8.08 N 14.09.

(3aS,4R,6aR)-3,3a,4,6a-Tetrahydro-4-pentylfuro[3,4c]pyrazol-6-one (ent-13): a) (R)-5-Pentyl-2-(5H)-furanone (852 mg, 95%) was prepared from (R,R)-4 (1.00 g, 5.80 mmol) by an analogous procedure to that described for 13 (part a): $[\alpha]_{\text{D}} = -94.8$ ($c = 0.9$ in CHCl_3); IR (film): $\tilde{\nu} = 3090, 2955, 2930, 2860, 1750, 1600, 1465, 1330, 1165, 1115, 1100, 1030, 1000, 960, 920, 900, 815$ cm^{-1} ; $t_{\text{r}}(\text{R}) = 29.75$ min, $t_{\text{r}}(\text{S}) = 33.80$ min (140°C, 100 kPa); 97% ee.

b) The title compound (219 mg, 86%) was prepared from (R)-5-pentyl-2-(5H)-furanone (200 mg, 1.30 mmol, 97% ee) by an analogous procedure to that described for 13 (part b): ^1H NMR (500 MHz, CDCl_3): δ = 0.90 (m , $5'\text{-H}_3$), 1.27–1.49 (m , $2'\text{-H}_2$, $3'\text{-H}_2$, $4'\text{-H}_2$), 1.63–1.78 (m , $1'\text{-H}_2$), 2.67 (dddd, $J_{3a,6a} = 9.0$, $J_{3a,3\text{-H(A)}} = 7.9$, $J_{3a,4} = 5.2$, $J_{3a,3\text{-H(B)}} = 2.8$ Hz, 3a-H), 3.92 (dt , $J_{4,1'\text{-H(1)}} = 7.5$, $J_{4,3a} = J_{4,1'\text{-H(2)}} = 5.3$ Hz, 4-H), AB signal ($\delta_{\text{A}} = 4.75$, $\delta_{\text{B}} = 4.83$, $J_{\text{AB}} = 18.5$ Hz, A part in addition split by $J_{\text{A},3a} = 8.2$, $J_{\text{A},6a} = 2.0$ Hz, B part in addition split by $J_{\text{B},3a} = J_{\text{B},6a} = 2.6$ Hz, 3-H₂), 5.52 (dt , $J_{6a,3a} = 9.1$, $J_{6a,3\text{-H(A)}} = J_{6a,3\text{-H(B)}} = 2.2$ Hz, 6a-H) ppm.

(3aS,4S,6aS)-1,3a,4,6a-Tetrahydro-6a-methyl-4-pentylfuro[3,4c]pyrazol-6-one (14) and (3aR,4S,6aS)-3,3a,4,6a-tetrahydro-6a-methyl-4-pentylfuro[3,4c]pyrazol-6-one (15): At -78°C , $n\text{BuLi}$ (2.05 M in hexane, 0.84 mL, 1.73 mmol, 1.7 equiv) was added to a solution of diisopropylamine (0.242 mL, 175 mg, 1.73 mmol, 1.7 equiv) in THF (1.7 mL). After the mixture had been stirred for 30 min, a solution of 13 (200 mg, 1.02 mmol) in THF (1.5 mL) was added and the mixture was stirred for 30 min at -78°C . Methyl iodide (0.13 mL, 0.29 g, 2.0 mmol, 2.0 equiv) was added. The solution was allowed to warm to -40°C and stirred for 18 h. After addition of H_2O the mixture was allowed to reach room temperature, extracted with EtOAc (4×20 mL), and dried over MgSO_4 . Evaporation in vacuo followed by flash chromatography (cyclohexane/EtOAc 6:1) afforded the separated title compounds in a 79:21 ratio (14: 138 mg, 65%; 15: 33 mg, 15%).

14: IR (film): $\tilde{\nu} = 3030, 2960, 2935, 2860, 1770, 1380, 1235, 1230, 1215, 1210, 1200, 1190, 1145, 1115, 1100, 800$ cm^{-1} ; HRMS: m/z calcd for $\text{C}_{11}\text{H}_{18}\text{N}_2\text{O}_2$: 210.137130; found 210.136828; elemental analysis calcd (%) for $\text{C}_{11}\text{H}_{18}\text{N}_2\text{O}_2$ (210.3): C 62.84, H 8.63, N 13.32; found: C 62.78, H 8.82, N 13.19.

15: IR (film): $\tilde{\nu} = 2955, 2935, 2860, 1770, 1550, 1455, 1435, 1375, 1355, 1290, 1255, 1240, 1175, 1110, 1015, 995, 945, 935, 905$ cm^{-1} ; elemental analysis calcd (%) for $\text{C}_{11}\text{H}_{18}\text{N}_2\text{O}_2$ (210.3): C 62.84, H 8.63, N 13.32; found: C 62.94, H 8.75, N 13.05.

(3aR,4R,6aR)-1,3a,4,6a-Tetrahydro-6a-methyl-4-pentylfuro[3,4c]pyrazol-6-one (ent-14) and (3aS,4R,6aR)-3,3a,4,6a-tetrahydro-6a-methyl-4-pentylfuro[3,4c]pyrazol-6-one (ent-15): The title compounds (ent-14: 62 mg, 32%; ent-15: 98 mg, 51%) were prepared in a 42:58 ratio from ent-13 (180 mg, 0.917 mmol) by an analogous procedure to that described for 14 and 15.

ent-14: ^1H NMR (500 MHz, CDCl_3 ; * = interchangeable): δ = 0.90–0.93 (m , $5'\text{-H}_3$), 1.30–1.56 (m , $2'\text{-H}_2$, $3'\text{-H}_2$, $4'\text{-H}_2$), superimposed by 1.54 (s , $1'\text{-H}_3$), 1.63–1.70 (m , $1'\text{-H}^1$), 1.76–1.84 (m , $1'\text{-H}^2$), 3.24 (m , 3a-H), 4.38 (ddd, $J_{4,1'\text{-H(1)}} = 8.6^*$, $J_{4,1'\text{-H(2)}} = 5.9^*$, $J_{4,3a} = 3.1$ Hz, 4-H), 6.03 (brs, NH), 6.70 (d , $J_{3,3a} = 1.1$ Hz, 3-H) ppm.

ent-15: ^1H NMR (500 MHz, CDCl_3): δ = 0.90 (m , $5'\text{-H}_3$), 1.27–1.50 (m , $2'\text{-H}_2$, $3'\text{-H}_2$, $4'\text{-H}_2$), 1.58–1.76 (m , $1'\text{-H}_2$), superimposed with 1.59 (s , $1'\text{-H}_3$), 2.29 (ddd, $J_{3a,3\text{-H(A)}} = 7.5$, $J_{3a,4} = 5.5$, $J_{3a,3\text{-H(B)}} = 2.0$ Hz, 3a-H), 3.78 (ddd, $J_{4,1'\text{-H(1)}} = 7.5$, $J_{4,1'\text{-H(2)}} = J_{4,3a} = 5.5$ Hz, 4-H), AB signal ($\delta_{\text{A}} = 4.71$, $\delta_{\text{B}} = 4.82$, $J_{\text{AB}} = 18.5$ Hz, A part in addition split by $J_{\text{A},3a} = 7.5$ Hz, B part in addition split by $J_{\text{B},3a} = 2.0$ Hz, 3-H₂) ppm.

(5S)-4-Hydroxy-3,5-dimethyl-5-pentyl-2-(3H)-furanone ((S)-18): a) At -78°C , $n\text{-BuLi}$ (2.27 M in THF, 2.47 mL, 5.60 mmol, 2.8 equiv) was added to a solution of diisopropylamine (0.79 mL, 0.57 g, 5.6 mmol, 2.8 equiv) in THF (8 mL). After the mixture had been stirred for 30 min, a solution of (4R,5S)-10 (370 mg, 2.00 mmol, 84% ee) in THF (3 mL) was added and the mixture was stirred for 1 h. A solution of methyl iodide (0.22 mL, 0.51 g, 3.6 mmol, 1.8 equiv) in DMPU (5.4 mL) and THF (7.1 mL) was added to this mixture during 90 min. It was then warmed to -40°C and stirred overnight at this temperature. Aqueous HCl (2%, 30 mL) was added and the mixture was allowed to reach room temperature. After extraction with EtOAc (3×30 mL), the combined organic extracts were washed with aq. NaCl (2×30 mL) and dried over MgSO_4 . Evaporation in vacuo followed by flash chromatography (cyclohexane/EtOAc 3:1) afforded (3R,4R,5S)-4,5-dihydro-4-hydroxy-3,5-dimethyl-5-pentyl-2-(3H)-furanone (295 mg, 74%): $[\alpha]_{\text{D}} = -5.0$ ($c = 0.3$ in CHCl_3); $t_{\text{r}}(3R,4R,5S) = 30.70$ min, $t_{\text{r}}(3S,4S,5R) = 28.51$ min (140°C, 100 kPa); 86% ee; elemental analysis calcd (%) for $\text{C}_{11}\text{H}_{20}\text{O}_3$ (200.3): C 65.97, H 10.07; found: C 65.85, H 9.87.

b) At -78°C , trifluoroacetic anhydride (0.33 mL, 0.49 g, 2.3 mmol, 2.0 equiv) was added to a solution of DMSO (0.29 mL, 0.32 g, 4.1 mmol, 3.5 equiv) in CH_2Cl_2 (6 mL). The resulting mixture was stirred for 30 min. After addition of a solution of (3R,4R,5S)-4,5-dihydro-4-hydroxy-3,5-dimethyl-5-pentyl-2-(3H)-furanone (232 mg, 1.16 mmol, 86% ee) in CH_2Cl_2 (6 mL), the mixture was stirred for 2 h at -78°C . Triethylamine (0.65 mL, 0.47 g, 4.6 mmol, 4.0 equiv) was then added and the mixture was stirred for 14 h. After addition of H_2O (15 mL) the solution was allowed to reach room temperature and was extracted with CH_2Cl_2 (5×20 mL). Drying over MgSO_4 and evaporation in vacuo followed by flash chromatography (cyclohexane/EtOAc 3:1) afforded the title compound (229 mg, 99%): $[\alpha]_{\text{D}} = +0.6$ ($c = 1.1$ in CHCl_3); IR (film): $\tilde{\nu} = 2955, 2930, 2870, 2865, 1720, 1650, 1460, 1405, 1370, 1335, 1310, 1265, 1170, 1105, 1050$ cm^{-1} ; $t_{\text{r}}(\text{S}) = 15.58$ min, $t_{\text{r}}(\text{R}) = 12.69$ min (140°C, 100 kPa); 85% ee; elemental analysis calcd (%) for $\text{C}_{11}\text{H}_{18}\text{O}_3$ (198.2): C 66.64, H 9.15; found: C 66.42, H 9.34.

(5R)-4-Hydroxy-3,5-dimethyl-5-pentyl-2-(3H)-furanone ((R)-18):

a) (3S,4S,5R)-4,5-Dihydro-4-hydroxy-3,5-dimethyl-5-pentyl-2-(3H)-furanone (471 mg, 72%) was prepared from (4S,5R)-10 (611 mg, 3.28 mmol, 80% ee) by an analogous procedure to that described for (S)-18 (part a): $[\alpha]_{\text{D}} = +4.3$ ($c = 0.6$ in CHCl_3); ^1H NMR (500 MHz, CDCl_3): δ = 0.90 (t , $J_{5,4} = 7.0$ Hz, $5'\text{-H}$), 1.25–1.46 (m , $2'\text{-H}_2$, $3'\text{-H}_2$, $4'\text{-H}_2$), superimposed with 1.30 (d , $J_{3\text{-Me},3} = 7.1$ Hz, 3-CH₃) and 1.33 (s , 5-CH₃), 1.63–1.75 (m , $1'\text{-H}_2$), 2.44 (brd , $J_{4\text{-OH},4} = 5.2$ Hz, 4-OH), 2.66 (dq , $J_{3,4} = 9.8$, $J_{3,3\text{-Me}} = 7.1$ Hz, 3-H), 3.86 (dd , $J_{4,3} = 9.9$, $J_{4,4\text{-OH}} = 5.2$ Hz, 4-H) ppm; IR (film): $\tilde{\nu} = 3445, 2955, 2935, 2875, 1755, 1750, 1460, 1385, 1350, 1310, 1255, 1245, 1200, 1125, 1110, 1075, 1040, 945, 620$ cm^{-1} ; $t_{\text{r}}(3S,4S,5R) = 27.13$ min, $t_{\text{r}}(3R,4R,5S) = 30.08$ min (140°C, 100 kPa); 78% ee.

b) The title compound (278 mg, 99%) was prepared from (3*S*,4*S*,5*R*)-4,5-dihydro-4-hydroxy-3,5-dimethyl-5-pentyl-2-(3*H*)-furanone (248 mg, 1.42 mmol, 80% *ee*) by an analogous procedure to that described for (S)-**18** (part b): $[\alpha]_{\text{D}} = -1.0$ ($c = 1.0$ in CHCl_3); $^1\text{H NMR}$ (500 MHz, CDCl_3): $\delta = 0.86$ (m , 5'-H₃), 1.13–1.35 (m , 2'-H₂, 3'-H₂, 4'-H₂), 1.47 (s , 5-CH₃), 1.71–1.86 (m , 1'-H₂), superimposed with 1.73 (s , 3-CH₃), 10.07 (brs , 4-OH) ppm; $t_{\text{R}}(R) = 13.88$ min, $t_{\text{R}}(S) = 16.81$ min (140°C, 100 kPa); 82% *ee*.

(5*S*)-2-Methoxy-3,5-dimethyl-5-pentyl-4-(5*H*)-furanone ((S)-19**):** At room temperature, $\text{Me}_3\text{O}^+\text{BF}_4^-$ (199 mg, 1.35 mmol, 3.0 equiv) was added to a solution of (S)-**18** (89 mg, 0.45 mmol) in CH_2Cl_2 (7 mL). After the mixture had been stirred for 28 h, aq. NaHCO_3 (9 mL) was added. The solution was extracted with CH_2Cl_2 (5×10 mL) and dried over MgSO_4 . After evaporation in vacuo the residue was purified by flash chromatography (cyclohexane/EtOAc 3:1) to afford the title compound (81 mg, 85%): $[\alpha]_{\text{D}} = -54.5$ ($c = 0.5$ in CHCl_3); $t_{\text{R}}(S) = 11.14$ min, $t_{\text{R}}(R) = 11.68$ min (120°C, 100 kPa), 87% *ee*; IR (film): $\tilde{\nu} = 3000, 2960, 2930, 2875, 2865, 1595, 1480, 1455, 1400, 1375, 1195, 1125, 980, 800, 785, 765, 760, 750, 745, 730, 725, 715, 710, 675, 665$ cm^{-1} ; elemental analysis calcd (%) for $\text{C}_{12}\text{H}_{20}\text{O}_3$ (212.3): C 67.95, H 9.43; found: C 68.04, H 9.73.

(5*R*)-2-Methoxy-3,5-dimethyl-5-pentyl-4-(5*H*)-furanone ((R)-19**):** The title compound (173 mg, 87%) was prepared from (R)-**18** (185 mg, 0.933 mmol) by an analogous procedure to that described for (S)-**19**: $[\alpha]_{\text{D}} = +50.6$ ($c = 0.7$ in CHCl_3); $^1\text{H NMR}$ (500 MHz, CDCl_3): $\delta = 0.86$ (m , 5'-H₃), 1.17–1.34 (m , 2'-H₂, 3'-H₂, 4'-H₂), 1.39 (s , 5-CH₃), 1.59 (s , 3-CH₃), 1.69–1.80 (m , 1'-H₂), 4.01 (s , 2-OCH₃); $t_{\text{R}}(R) = 11.58$ min, $t_{\text{R}}(S) = 11.17$ min (120°C, 100 kPa); 81% *ee*.

(3*R*,4*R*,5*R*)-4,5-Dihydro-4-hydroxy-3-methyl-5-pentyl-2-(3*H*)-furanone (20**):** At -78°C , *n*-BuLi was added to a solution of diisopropylamine (1.41 mL, 1.02 g, 10.1 mmol, 5.0 equiv) in THF (7 mL). After the mixture had been stirred for 30 min, a solution of (R,R)-**4** (347 mg, 2.02 mmol, 97% *ee*) in THF (2 mL) was added. The solution was stirred for 1 h at -78°C , then a solution of methyl iodide (1.25 mL, 2.86 g, 20.2 mmol, 10.0 equiv) in THF (7 mL) was added. After the mixture had been stirred for 2 h, a solution of glacial acetic acid (1.7 mL) in THF (7 mL) was added and the mixture was allowed to reach room temperature. Aqueous NaHCO_3 (35 mL) was added and the aqueous phase was extracted with EtOAc (4×30 mL). The combined organic extracts were dried over MgSO_4 and evaporated in vacuo. The residue was purified by flash chromatography (cyclohexane/EtOAc 4:1) to afford the title compound (311 mg, 83%): $[\alpha]_{\text{D}} = +78.3$ ($c = 1.2$ in CHCl_3); $^1\text{H NMR}$ (500 MHz, CDCl_3): $\delta = 0.90$ (m , 5'-H₃), 1.26–1.46 (m , 2'-H₁, 3'-H₂, 4'-H₂), superimposed with 1.30 (d , $J_{3-\text{Me},3} = 7.7$ Hz, 3-CH₃), 1.48–1.58 (m , 2'-H₂), 1.69–1.80 (m , 1'-H₂), 2.42 (d , $J_{4-\text{OH},4} = 4.9$ Hz, 4-OH), 2.61 (qd , $J_{3,3-\text{Me}} = 7.6$, $J_{3,4} = 3.8$ Hz, 3-H), 4.14 (ddd , $J_{4,4-\text{OH}} = J_{4,5} = 4.9$, $J_{4,3} = 3.9$ Hz, 4-H), 4.46 (m , 5-H) ppm; IR (film): $\tilde{\nu} = 3450, 2955, 2930, 2875, 2860, 1760, 1460, 1380, 1355, 1330, 1235, 1205, 1130, 1085, 1045, 1025, 1000, 955$ cm^{-1} .

(3*S*,4*S*,5*S*)-4,5-Dihydro-4-hydroxy-3-methyl-5-pentyl-2-(3*H*)-furanone (*ent*-20**):** The title compound (945 mg, 87%) was prepared from (S,S)-**4** (1.0 g, 5.8 mmol, 94% *ee*) by an analogous procedure to that described for **20**: $[\alpha]_{\text{D}} = -74.6$ ($c = 1.5$ in CHCl_3); elemental analysis calcd (%) for $\text{C}_{10}\text{H}_{18}\text{O}_3$ (186.3): C 64.50, H 9.74; found: C 64.23, H 9.94.

(5*R*)-4-Hydroxy-3-methyl-5-pentyl-2-(3*H*)-furanone (21**):** At -78°C , tri-fluoroacetic anhydride (0.47 mL, 0.70 g, 3.3 mmol, 2.0 equiv) was added to a solution of DMSO (0.41 mL, 0.45 g, 5.8 mmol, 3.5 equiv) in CH_2Cl_2 (9 mL). After the mixture had been stirred for 30 min, a solution of **20** (309 mg, 1.66 mmol) in CH_2Cl_2 (9 mL) was added and the mixture was stirred for 1 h at -78°C . After addition of triethylamine (0.93 mL, 0.67 mg, 6.6 mmol, 4.0 equiv) and stirring for 45 min, H_2O (20 mL) was added and the solution was allowed to reach room temperature. After the mixture had been stirred for 30 min, the aqueous phase was extracted with CH_2Cl_2 (4×30 mL) and the combined organic extracts were dried over MgSO_4 . Evaporation in vacuo followed by flash chromatography (cyclohexane/EtOAc 2:1) afforded the title compound (281 mg, 92%): $[\alpha]_{\text{D}} = +15.6$ ($c = 1.0$ in CHCl_3).

(5*S*)-4-Hydroxy-3-methyl-5-pentyl-2-(3*H*)-furanone (*ent*-21**):** The title compound (776 mg, 94%) was prepared from *ent*-**20** (831 mg, 4.46 mmol) by an analogous procedure to that described for **20**: $[\alpha]_{\text{D}} = -14.5$ ($c = 1.0$ in CHCl_3); $^1\text{H NMR}$ (500 MHz, CDCl_3 ; *, ** = interchangeable): $\delta = 0.87$

(m , 5'-H₃), 1.23–1.46 (m , 2'-H₂, 3'-H₂, 4'-H₂), 1.61 ($dddd$, $J_{\text{gem}} = 14.5$, $J_{1'-\text{H}(1),2'-\text{H}(1)} = 9.8$, $J_{1'-\text{H}(1),5} = 7.8$, $J_{1'-\text{H}(1),2'-\text{H}(2)} = 5.2$ Hz, 1'-H¹), 1.73 (d , $^5J_{3-\text{Me},5} = 1.2$ Hz, 3-CH₃), 1.98 ($dddd$, $J_{\text{gem}} = 14.1$, $J_{1'-\text{H}(2),2'-\text{H}(1)} = 9.6^*$, $J_{1'-\text{H}(2),2'-\text{H}(2)} = 6.1^*$, $J_{1'-\text{H}(2),5} = 3.6$ Hz, 1'-H²), 4.76 (incompletely resolved ddq , $J_{5,1'-\text{H}(1)} = 7.7^{**}$, $J_{5,1'-\text{H}(2)} = 3.4^{**}$, $^3J_{5,3-\text{Me}} = 1.2$ Hz, 5-H), 10.71 (brs , 4-OH) ppm; IR (KBr): $\tilde{\nu} = 2970, 2955, 2925, 2875, 2855, 1710, 1645, 1270, 1230, 1220, 1100, 1080$ cm^{-1} ; elemental analysis calcd (%) for $\text{C}_{10}\text{H}_{16}\text{O}_3$ (184.2): C 65.20, H 8.76; found: C 65.20, H 8.90.

Acknowledgements

This work was financially supported by the Fonds der Chemischen Industrie. We are indebted to Valentina Morgel for her skillful technical assistance and we express our gratitude to Dr. Manfred Keller (Institut für Organische Chemie und Biochemie, Universität Freiburg) for performing the X-ray analyses and to Melanie Junge (Universität Hamburg) for preparing the GLC column for the analysis, the results of which are shown in Figure 1.

- [1] Recent reviews on the chemistry of Δ^2 -butenolides: a) A. Hashem, E. Kleinpeter, *Adv. Heterocycl. Chem.* **2001**, *81*, 107–165; b) N. B. Carter, A. E. Nadany, J. B. Sweeney, *J. Chem. Soc. Perkin Trans. 1* **2002**, 2324–2342.
- [2] Y. Saritas, M. Mekem Sonwa, H. Iznaguen, W. A. König, H. Muhle, R. Mues, *Phytochemistry* **2001**, *57*, 443–457.
- [3] a) M. Junge, W. A. König, *J. Sep. Sci.* **2003**, *26*, 1607–1614; b) V. Schurig, *J. Chromatogr. A* **2001**, *906*, 275–299; c) W. A. König, *Gas Chromatographic Enantiomer Separation with Modified Cyclodextrins*, Hüthig, Heidelberg, **1992**.
- [4] Circular dichroism measurements have been used for determining the absolute configuration of differently substituted Δ^2 -butenolides: A. Gypser, C. Bülow, H.-D. Scharf, *Tetrahedron* **1995**, *51*, 1921–1930; J. K. Gawronski, A. v. Oeveren, H. v. d. Deen, C. W. Leung, B. L. Feringa, *J. Org. Chem.* **1996**, *61*, 1513–1515; V. J. R. V. Mukku, M. Speitling, H. Laatsch, E. Helmke, *J. Nat. Prod.* **2000**, *63*, 1570–1572; T. A. Mansoor, J. Hong, C.-O. Lee, C. J. Sim, K. S. Im, D. S. Lee, J. H. Jung, *J. Nat. Prod.* **2004**, *67*, 721–724; T. Murakami, Y. Morikawa, M. Hashimoto, T. Okuno, Y. Harada, *Org. Lett.* **2004**, *6*, 157–160; ref. [9].
- [5] We are unaware of a method allowing the assignment of absolute configurations to α,β,γ -trisubstituted or $\alpha,\beta,\gamma,\gamma$ -tetrasubstituted Δ^2 -butenolides based on $^1\text{H NMR}$ shift differences observed upon the addition of enantiopure lanthanide shift reagents.
- [6] Reviews: a) B. B. Lohray, *Tetrahedron: Asymmetry* **1992**, *3*, 1317–1349; b) R. A. Johnson, K. B. Sharpless, *Asymmetric Catalysis in Organic Synthesis* (Ed.: I. Ojima), VCH, New York, **1993**, 227–272; c) H. C. Kolb, M. S. VanNieuwenhze, K. B. Sharpless, *Chem. Rev.* **1994**, *94*, 2483–2547; d) G. Poli, C. Scolastico, *Methoden Org. Chem. (Houben-Weyl) 4th ed. 1952–*, Vol. *E21e* (Eds.: G. Helmchen, R. W. Hoffmann, J. Mulzer, E. Schaumann), Thieme, Stuttgart, **1995**, pp. 4547–4598; e) R. A. Johnson, K. B. Sharpless, *Catalytic Asymmetric Synthesis*, 2nd ed. (Ed.: I. Ojima), Wiley-VCH, New York, **2000**, 357–389; f) C. Bolm, J. P. Hildebrand, K. Muniz, *Catalytic Asymmetric Synthesis*, 2nd ed. (Ed.: I. Ojima), Wiley-VCH, New York, **2000**, 399–428.
- [7] C. Harcken, R. Brückner, *Angew. Chem.* **1997**, *109*, 2866–2868; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2750–2752.
- [8] Recent syntheses of enantiomerically pure butenolides include the following: W. Wang, Y. Zhong, G. Lin, *Tetrahedron Lett.* **2003**, *44*, 4613–4616; P. M. Donate, D. Frederico, R. da Silva, M. G. Constantino, G. Del Ponte, P. S. Bonatto, *Tetrahedron: Asymmetry* **2003**, *14*, 3253–3256; S. Koul, B. Singh, S. C. Taneja, G. N. Qazi, *Tetrahedron* **2003**, *59*, 3487–3491; K. Suzuki, K. Inomata, Y. Endo, *Heterocycles* **2003**, *60*, 2743–2748; R. B. Chhor, B. Nosse, S. Sörgel, C. Böhm, M. Seitz, O. Reiser, *Chem. Eur. J.* **2003**, *9*, 260–270; H. Miyabe, M. Ueda, K. Fujii, A. Nishimura, T. Naito, *J. Org. Chem.* **2003**, *68*,

- 5618–5626; A. Samarat, H. Amri, Y. Landais, *Tetrahedron* **2004**, *60*, 8949–8956; P. R. Krishna, V. V. R. Reddy, G. V. M. Sharma, *Synthesis* **2004**, 2107–2114; N. J. Kerrigan, P. C. Hutchison, T. D. Heightman, D. J. Procter, *Org. Biomol. Chem.* **2004**, *2*, 2476–2482; P. V. Ramachandran, D. Pratihar, D. Biswas, A. Srivastava, M. V. R. Reddy, *Org. Lett.* **2004**, *6*, 481–484; L. Thijs, B. Zwanenburg, *Tetrahedron* **2004**, *60*, 5237–5252.
- [9] Recent syntheses of enantiomerically pure Δ^2 -butenolides include the following: a) S. P. Brown, N. C. Goodwin, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2003**, *125*, 1192–1194; b) W. P. Gallagher, R. E. Maleczka, Jr., *J. Org. Chem.* **2003**, *68*, 6775–6779; c) S. Ma, Z. Yu, *J. Org. Chem.* **2003**, *68*, 6149–6152; d) K. Suzuki, K. Inomata, *Tetrahedron Lett.* **2003**, *44*, 745–749; e) T. Nagamitsu, D. Takano, K. Shiomi, H. Ui, Y. Yamaguchi, R. Masuma, Y. Harigaya, I. Kuwajima, S. Omura, *Tetrahedron Lett.* **2003**, *44*, 6441–6444; f) J.-L. Lee, C.-F. Lin, L.-Y. Hsieh, W.-R. Lin, H.-F. Chiu, Y.-C. Wu, K.-S. Wang, M.-J. Wu, *Tetrahedron Lett.* **2003**, *44*, 7833–7835; g) Y. Isegori, Y. Kobayashi, *Synlett* **2004**, *11*, 1941–1944; h) A. Kirschner, P. Langer, U. T. Bornscheuer, *Tetrahedron: Asymmetry* **2004**, *15*, 2871–2874; i) M. Braun, A. Hohmann, J. Rahematpura, C. Bühne, S. Grimme, *Chem. Eur. J.* **2004**, *10*, 4584–4593.
- [10] a) C. Harcken, R. Brückner, E. Rank, *Chem. Eur. J.* **1998**, *4*, 2342–2352 (corrigendum: C. Harcken, R. Brückner, E. Rank, *Chem. Eur. J.* **1998**, *4*, 2390); b) T. Berkenbusch, R. Brückner, *Tetrahedron* **1998**, *54*, 11461–11470; c) T. Berkenbusch, R. Brückner, *Tetrahedron* **1998**, *54*, 11471–11480; d) C. Harcken, R. Brückner, *New J. Chem.* **2001**, *25*, 40–54; e) C. Harcken, R. Brückner, *Synlett* **2001**, 718–721; f) C. Harcken, R. Brückner, *Tetrahedron Lett.* **2001**, *42*, 3967–3971.
- [11] A slightly different preparation of *trans*-**5** from **3** was described by C. García, T. Martín, V. S. Martín, *J. Org. Chem.* **2001**, *66*, 1420–1428 (step 1 in xylene with catalytic piperidine).
- [12] Method: a) N. Ragoussis, *Tetrahedron Lett.* **1987**, *28*, 93–96; b) H. Yamanaka, M. Yokoyama, T. Sakamoto, T. Shiraishi, M. Sagi, M. Mizugaki, *Heterocycles* **1983**, *20*, 1541–1544.
- [13] The enantiopurity of this compound was determined—after dehydration (**4**) or directly (**10**)—by chiral GLC with β -cyclodextrin bonded to dimethylpolysiloxane as a stationary phase.
- [14] CCDC-249458 and CCDC-249459 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [15] A. L. Spek, *J. Appl. Crystallogr.* **2003**, *36*, 7–13.
- [16] Ester **11** (*E*:*Z* = 96:4) was earlier prepared in a different manner: E. J. Corey, R. H. K. Chen, *Tetrahedron Lett.* **1973**, 1611–1614.
- [17] Acid *E*-**7** was obtained in isomerically pure form by conjugate addition of Pent_2CuLi to 2-butyric acid at $-78^\circ\text{C} \rightarrow -15^\circ\text{C}$; method: a) J. Klein, R. M. Turkel, *J. Am. Chem. Soc.* **1969**, *91*, 6186–6187; b) R. M. Carlson, A. R. Oyler, J. R. Peterson, *J. Org. Chem.* **1975**, *40*, 1610–1616; c) Y. Yamamoto, H. Yatagai, K. Maruyama, *J. Org. Chem.* **1979**, *44*, 1744–1746.
- [18] Acid (*Z*)-**7** was obtained in isomerically pure form by conjugate addition (same procedure as in ref. [17]) of Me_2CuLi to 2-octynoic acid; preparation: J. Chenault, J.-F. E. Dupin, *Synthesis* **1987**, 498–499.
- [19] Review: W. E. Bachmann, W. S. Struve, *Org. React.* **1942**, *1*, 38–62.
- [20] Arndt–Eistert homologation of α,β -unsaturated carboxylic chlorides: J. A. Moore, *J. Org. Chem.* **1955**, *20*, 1607–1612. Syntheses of α,β -unsaturated diazoketones from α,β -unsaturated carboxylic chlorides: M. Berenbom, W. S. Fones, *J. Org. Chem.* **1949**, *14*, 1629–1629; P. Ceccherelli, M. Curini, M. C. Marcotullio, O. Rosati, *J. Org. Chem.* **1990**, *55*, 311–315; P. Ceccherelli, M. Curini, M. C. Marcotullio, O. Rosati, E. Wenkert, *J. Org. Chem.* **1991**, *56*, 7065–7070; G. D. Prestwich, I. Ujváry, *J. Labelled Compd. Radiopharm.* **2002**, *2*, 37–38.
- [21] S. Hanessian, P. J. Murray, *J. Org. Chem.* **1987**, *52*, 1170–1171.
- [22] Method: a) H.-M. Shieh, G. D. Prestwich, *J. Org. Chem.* **1981**, *46*, 4319–4321; b) A. R. Chamberlin, M. Dezube, *Tetrahedron Lett.* **1982**, *23*, 3055–3058; c) J. E. Wrobel, B. Ganem, *J. Org. Chem.* **1983**, *48*, 3761–3764.
- [23] Reviews: a) A. J. Mancuso, D. Swern, *Synthesis* **1981**, 165; b) T. T. Tidwell, *Synthesis* **1990**, 857–870; c) T. T. Tidwell, *Org. React.* **1990**, *39*, 297–572.
- [24] a) J. E. Wrobel, B. Ganem, *J. Org. Chem.* **1983**, *48*, 3761–3764; b) K. Matsuno, Y. Sakaguchi, *Heterocycles* **1996**, *43*, 2553–2556; c) K. Matsuno, Y. Sakaguchi, *Chem. Pharm. Bull.* **1997**, *45*, 1620–1625. Otherwise, we are only aware of oxidations of α,β -dihydroxy- γ -butyrolactones leading to α -hydroxytetronic acids (vitamin C type tetronic acids).
- [25] Leading reference for syntheses of enantiopure tetronic acids: D. Tejedor, F. García-Tellado, *Org. Prep. Proced. Int.* **2004**, *36*, 35–59.
- [26] This method was established for $\text{C}^4=\text{O}$ -selective methylations of tetronic acids: F. A. Lakhvich, F. S. Pashkovskii, L. G. Lis, *J. Org. Chem. USSR (Engl. Transl.)* **1992**, *28*, 1295–1301. For $\text{C}^2=\text{OH}$ -versus $\text{C}^4=\text{O}$ -selective methylations, see also: A. Schnedler Wengel, T. Reffstrup, P. M. Boll, *Tetrahedron* **1979**, *35*, 2181–2185; M. O. Garcia, A. D. Rodriguez, *Tetrahedron* **1990**, *46*, 1119–1124; L. Trifonov, J. H. Bieri, R. Prewé, A. S. Dreiding, *Tetrahedron* **1982**, *38*, 397–403.
- [27] For example: A. B. Smith III, E. G. Nolen, Jr., R. Shirai, F. R. Blase, M. Ohta, N. Chida, R. A. Hertz, D. M. Fitch, W. M. Clark, P. A. Sprengeler, *J. Org. Chem.* **1995**, *60*, 7837–7848; Y. Horiguchi, E. Nakamura, I. Kuwajima, *J. Am. Chem. Soc.* **1989**, *111*, 6257–6265; D. Schinzer, C. Allagiannis, S. Wichmann, *Tetrahedron* **1988**, *44*, 3851–3868; L. H. Zalkow, F. X. Merkley, C. Djerassi, *J. Am. Chem. Soc.* **1960**, *82*, 6354–6362.
- [28] B. S. Furniss, A. J. Hannaford, P. W. G. Smith, A. R. Tatchell, *Vogel's Textbook of Practical Organic Chemistry*, 5th ed., Longman, Essex, England, **1989**, 430–432.
- [29] W. C. Still, M. Kahn, A. Mitra, *J. Org. Chem.* **1978**, *43*, 2923–2925.
- [30] E. Pretsch, W. Simon, J. Seibl, T. Clerc, *Spectral data for structure determination of organic compounds*, 2nd ed., Springer, Berlin, **1989**, C90.

Received: November 10, 2004
Published online: February 15, 2005