

Structural Effects of Olefins in the Photooxygenation with Electron-Accepting Sensitizers.
Kinetic Approach to Reactive Intermediates

Satoshi KONUMA, Shin AIHARA, Yasunao KURIYAMA, Hiroaki MISAWA,[†] Ryoichi AKABA,^{††}
Hirochika SAKURAGI,* and Katsumi TOKUMARU*

Department of Chemistry, University of Tsukuba, Tsukuba, Ibaraki 305

Pulsed laser excitation studies on the reactive species in 9-cyanoanthracene(CNA)-sensitized oxygenation of 1,1-diphenyl-2-methylpropene (**1a**) and (*E*)-2,3-diphenyl-2-butene (**1b**) show that the reaction course is mostly governed by competitive quenching of the CNA excited singlet by **1** and oxygen to produce **1**⁺ and ¹O₂, respectively. The results indicate that the reaction courses of tetraphenylethylene and 2,3-dimethyl-2-butene, for example, can be explained by exclusive quenching of sensitizer singlets by the olefin and oxygen, respectively.

There remains controversy in reactive oxygen species in electron-transfer induced oxygenation of olefins.¹⁾ The excited singlet state of cyanoaromatic sensitizers accepts an electron from olefins in acetonitrile to give the olefin radical cations and the sensitizer radical anions, and the subsequent reactions under oxygen afford various types of products depending on the structure of olefins.²⁻⁶⁾ Arylethylenes such as tetraphenylethylene^{2,3)} and (*E*)-stilbene⁴⁾ are reported to afford carbonyl compounds accompanied by oxiranes or other oxygenated products possibly through reactions of the olefin radical cations with superoxide anions (O₂⁻) generated by secondary electron transfer from the acceptor radical anions to oxygen. On the other hand, alkenes exhibit completely different behavior; 2,3-dimethyl-2-butene⁵⁾ and 1,2-dimethylcyclohexene⁶⁾ give almost exclusively hydroperoxides through the ene-reaction with singlet oxygen produced by energy transfer from the sensitizer excited states.


To get deeper insight into the reaction mechanism and the factors controlling the reaction course of electron acceptor-sensitized oxygenation of olefins, we attempted a kinetic approach to determine the reaction rates for various reactive intermediates. 1,1-Diphenyl-2-methylpropene (**1a**) and (*E*)-2,3-diphenyl-2-butene (**1b**) were examined in acetonitrile in the presence of 9-cyanoanthracene (CNA) by laser excitation as well as steady irradiation (for **1a**). 9,10-Dicyanoanthracene-sensitized oxygenation of (*Z*)- and (*E*)-2,3-diphenyl-2-butene occurs exclusively via the singlet-oxygen pathway in carbon tetrachloride; however, in acetonitrile the singlet-oxygen and electron-transfer oxygenations compete with one another.³⁾

Irradiation of CNA (2×10^{-4} M, M = mol dm⁻³) with **1a** (0.01 M) in acetonitrile under oxygen with 405-nm light from a high-pressure mercury lamp yielded benzophenone (**2**), an oxirane (**3**), a hydroperoxide (**4**), and an

[†] Present address: Micro Photoconversion Project, Research Development Corporation of Japan, 15 Morimotocho, Shimogamo, Sakyo-ku, Kyoto 606.

^{††} Present address: Department of Chemistry, Gunma College of Technology, Toriba-machi, Maebashi, Gunma 376.

enal (**5**) (29, 13, 21, and 11%, respectively, based on **1a** consumed).⁷⁾ Hydroperoxide **4** arises undoubtedly from the reaction of **1** with singlet oxygen since, in a control experiment, rose bengal-sensitized oxygenation of **1a** in acetonitrile with visible light almost exclusively afforded **4** in 90% yield. Products **2**, **3**, and **5** can arise from free radical pathways since di-*t*-butyl peroxide- (photolysis at 313 nm) or azobisisobutyronitrile-initiated (thermolysis at 80 °C) autoxidation of **1a** gave **2**, **3**, and **5** (in comparable yields of **2** and **3** with a small amount of **5**). **2** was also afforded in a considerable yield in the anodic oxidation of **1a** (platinum electrodes, 1.65 V vs. SCE) under oxygen in acetonitrile. Olefins often give carbonyl compounds and oxiranes in various pathways. However, the production of hydroperoxide **4** and enal **5** on CNA sensitization clearly indicates the generation of singlet oxygen and allylic radicals, $\text{Ph}_2\text{C}=\text{CMeCH}_2\cdot$, as intermediates; the latter reacts with oxygen to give peroxy radicals, subsequently affording **5** through several steps.




Fluorescence of CNA (lifetime $\tau_S = 16.8$ ns) was effectively quenched by **1a** and **1b** in acetonitrile with rate constants of 6.0×10^8 and $8.0 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, respectively. The estimated free energy change associated with electron transfer from ground-state **1a** or **1b** to excited singlet CNA shows that this process is exothermic.^{8–10)} The CNA fluorescence was also quenched by oxygen with an almost diffusion-controlled rate constant, $1.6 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$. Therefore, under the above oxygenation conditions ($[\text{1a}] = 0.01 \text{ M}$, $[\text{O}_2] \approx 0.01 \text{ M}$), the CNA singlet is quenched ≈ 30 times more effectively by molecular oxygen than by **1a**.

Pulsed laser excitation¹¹⁾ of acetonitrile solutions of CNA and **1a** afforded transient absorptions assignable to $\text{CNA}^{\cdot-}$ ($\lambda_{\text{max}} 590 \text{ nm}$)¹²⁾ and radical cation $\text{1a}^{\cdot+}$ (760–800 nm).¹³⁾ These absorptions decayed with nearly the same rate constants under argon according to second-order kinetics due to their recombination. However, in the presence of oxygen, the decay curves of these absorptions consisted of first- and second-order components ($\text{1a}^{\cdot+} + \text{CNA}^{\cdot-}$ or $\text{O}_2^{\cdot-}$); from the first-order components whose decay rates were linearly dependent on oxygen concentration, the quenching rate constants by oxygen were estimated as 1.1×10^{10} for $\text{CNA}^{\cdot-}$ and $4.8 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ for $\text{1a}^{\cdot+}$. The former value shows an efficient formation of superoxide anions under oxygen, and the latter value is ten times larger than those for stilbene derivatives reported previously;¹⁴⁾ for example, $1.3 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ for (*E*)-stilbene radical cations. Similar studies on **1b** afforded a rate constant of $2.6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ for $\text{1b}^{\cdot+} + \text{O}_2$ reaction.¹⁵⁾ Thus, $\text{1a}^{\cdot+}$ and $\text{1b}^{\cdot+}$ react with oxygen at much higher rates than alkene radical cations such as adamantylideneadamantane radical cations ($5600 \text{ M}^{-1} \text{ s}^{-1}$).¹⁶⁾ These results indicate that the rate constant of reaction of olefin radical cations with oxygen varies in a wide range depending on the structure of olefins.

The quantum yield for intersystem crossing of CNA is very low ($\phi_{\text{isc}} = 0.025$);¹⁷⁾ however, oxygen accelerates intersystem crossing of the CNA excited singlet.¹⁸⁾ The efficiency of the acceleration was measured in acetonitrile by the emission-absorption method;¹⁹⁾ a factor of $k_{\text{isc}}'/k_{\text{isc}}$ was obtained to be 8900 M^{-1} , where k_{isc} is the rate constant for intersystem crossing of the CNA excited singlet and k_{isc}' is a second-order rate constant for oxygen-induced intersystem crossing. Use of $\phi_{\text{isc}} = 0.025$ and $\tau_S = 16.8 \text{ ns}$ affords $k_{\text{isc}} = 1.5 \times 10^6 \text{ s}^{-1}$ and $k_{\text{isc}}' = 1.3 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$. Under the above oxygenation conditions ($[\text{1a}] = 0.01 \text{ M}$, $[\text{O}_2] \approx 0.01 \text{ M}$), the quantum yield for CNA triplet formation is estimated to be nearly 0.3. The CNA triplet has a sufficient excitation energy (174 kJ mol^{-1})²⁰⁾ to excite $^3\text{O}_2$ to $^1\text{O}_2$ (94 kJ mol^{-1}), and is efficiently quenched by oxygen in acetonitrile ($k_q = 2.3 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$), as determined by monitoring its T-T absorption at 430 nm.¹⁷⁾

The production of singlet oxygen on excitation of CNA in the presence of oxygen was confirmed by observing its near-infrared emission at 1.27 μm , and the quantum yield for emission was measured to be ≈ 0.1 under air by using benzophenone as a standard ($\phi_{\Delta}=0.29$ in benzene, where ϕ_{Δ} is the quantum yield for $^1\text{O}_2$ formation on triplet quenching).²¹⁾ The decay of $^1\text{O}_2$ emission was accelerated by the addition of **1a** or **1b**. The rate constants for quenching of $^1\text{O}_2$ by **1a** and **1b** were determined to be 5.0×10^6 and $1.0 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$, respectively, by using benzophenone as a sensitizer for $^1\text{O}_2$ generation under air. These values are much lower than that reported for 2,3-dimethyl-2-butene, one of the most reactive olefins with $^1\text{O}_2$, $2.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$.²²⁾

Scheme 1.²³⁾

The direct observation of the intermediates concerned as well as the formation of various products, some of which are typical for specific pathways, clearly indicates the participation of several pathways in the present oxygenation. It is to be noted that the present olefins, **1a** and **1b**, exhibit intermediate behavior³⁾ between the olefins substituted only by alkyl groups and those substituted only by aromatic groups in the photooxygenation sensitized by electron acceptors. As exemplified above for CNA and **1a** (Scheme 1),²³⁾ the sensitizer singlets interact more rapidly with molecular oxygen than with the substrates, and the intersystem crossing of the sensitizer is very much accelerated by molecular oxygen to result in efficient generation of singlet oxygen. Thus, for olefins highly reactive with singlet oxygen such as alkyl-substituted olefins, singlet oxygen plays an exclusive role in their oxygenation. However, for olefins less reactive with singlet oxygen such as aralkenes, the olefin radical cations (reactive with $\text{O}_2^{\cdot-}$ or O_2) or the species derived from them such as allylic radicals play an important role in the oxygenation. Therefore, it is reasonable to conclude that the reactivity and reaction courses of various olefins in CNA- or 9,10-dicyanoanthracene-sensitized oxygenation are strongly governed by their reactivity with singlet oxygen.

The authors thank to the Ministry of Education, Science and Culture for partial support by Grant-in-Aid for Special Project Research No. 63104001 (K.T.).

References

- 1) a) C. S. Foote, *Tetrahedron*, **41**, 2221 (1985); b) F. D. Lewis, "Photoinduced Electron Transfer," ed by M. A. Fox and M. Chanon, Elsevier, Amsterdam (1988), Part C, pp. 1-69.
- 2) J. Eriksen, C. S. Foote, and T. L. Parker, *J. Am. Chem. Soc.*, **99**, 6455 (1977).

- 3) K. Gollnick and A. Schnatterer, *Photochem. Photobiol.*, **43**, 365 (1986).
- 4) J. Eriksen and C. S. Foote, *J. Am. Chem. Soc.*, **102**, 6083 (1980); L. T. Spada and C. S. Foote, *ibid.*, **102**, 391 (1980); L. E. Manring, M. K. Kramer, and C. S. Foote, *Tetrahedron Lett.*, **25**, 2523 (1984).
- 5) A. Albin and S. Sperti, *Gazzetta Chem. Ital.*, **115**, 227 (1985).
- 6) Y. Araki, D. C. Dobrowolski, T. E. Goynes, D. C. Hanson, Z. Q. Jiang, K. J. Lee, and C. S. Foote, *J. Am. Chem. Soc.*, **106**, 4570 (1984).
- 7) Product yields were determined by GLPC. Those of **4** were determined as the corresponding alcohol after treating the reaction mixtures with triphenylphosphine.
- 8) The free energy change ($-\Delta G^\circ = 7.5\text{--}12\text{ kJ mol}^{-1}$ for CNA singlets) was estimated according to the Rehm-Weller equation⁹⁾ from the following values: $E(\mathbf{1a}^+/\mathbf{1a}) = 1.61\text{ V}$, $E(\mathbf{1b}^+/\mathbf{1b}) = 1.57\text{ V}$,¹⁰⁾ $E(\text{CNA}/\text{CNA}^-) = -1.41\text{ V}$,^{1b)} $E_S(\text{CNA}) = 3.04\text{ eV}$.^{1b)}
- 9) D. Rehm and A. Weller, *Isr. J. Chem.*, **8**, 259 (1970).
- 10) M. Tsuchiya, R. Akaba, S. Aihara, H. Sakuragi, and K. Tokumaru, *Chem. Lett.*, **1986**, 1727.
- 11) Laser flash photolyses were performed by using an XeCl (308 nm) excimer laser (Lambda Physik EMG-101) for benzophenone and by an excimer laser-pumped dye laser (Lambda Physik FL-3002, DPS 406 nm) for CNA, and a pulsed xenon arc (Wacom KXL-151, 150 W) as a monitoring light source. The amplified signal (Hamamatsu R928) was recorded as the time profile of a transmittance change on a storage oscilloscope (Iwatsu TS-8123) and transferred to a personal computer (NEC PC-9801VX21) where it was analyzed.
- 12) H. Hiratsuka, H. Nakamura, Y. Tanizaki, and K. Nakajima, *Bull. Chem. Soc. Jpn.*, **55**, 3407 (1982).
- 13) This absorption was effectively quenched by azulene ($E(\text{Az}^+/\text{Az}) = 0.95\text{ V}$). 1,1-Bis(4-methoxyphenyl)-2-methylpropene (**1c**) also exhibited a transient spectrum in this region, and its behavior to the quenchers was similar to that of **1a**. The spectrum was identified as $\mathbf{1c}^+$ by comparison with that observed on γ -radiolysis of **1c**. We thank Prof. T. Shida, Kyoto University, for measuring the spectrum of $\mathbf{1c}^+$ by γ -radiolysis.
- 14) M. Tsuchiya, T. W. Ebbesen, Y. Nishimura, H. Sakuragi, and K. Tokumaru, *Chem. Lett.*, **1987**, 2121.
- 15) The transient absorption of $\mathbf{1b}^+$ was monitored at 480 nm; H. Suzuki, K. Koyano, T. Shida, and T. Kira, *Bull. Chem. Soc. Jpn.*, **55**, 3690 (1982); T. Shida, "Electronic Absorption Spectra of Radical Ions," Elsevier, Amsterdam (1988), p. 117.
- 16) S. F. Nelsen, D. L. Kapp, R. Akaba, and D. H. Evans, *J. Am. Chem. Soc.*, **108**, 6863 (1986); R. Akaba, S. Aihara, H. Sakuragi, and K. Tokumaru, *J. Chem. Soc., Chem. Commun.*, **1987**, 1262.
- 17) K. Kikuchi, H. Kokubun, and M. Koizumi, *Bull. Chem. Soc. Jpn.*, **41**, 1545 (1968).
- 18) L. E. Manring, C.-I. Gu, and C. S. Foote, *J. Phys. Chem.*, **87**, 40 (1983); D. C. Dobrowolski, P. R. Ogilby, and C. S. Foote, *ibid.*, **87**, 2261 (1983).
- 19) K. Tokumura, K. Kikuchi, and M. Koizumi, *Bull. Chem. Soc. Jpn.*, **46**, 1309 (1973).
- 20) K. Kikuchi, Y. Takahashi, M. Hoshi, T. Niwa, T. Katagiri, and T. Miyashi, *J. Phys. Chem.*, **95**, 2378 (1991).
- 21) A. A. Gorman, I. Hamblett, C. Lambert, A. L. Prescott, M. A. J. Rodgers, *J. Am. Chem. Soc.*, **106**, 4679 (1984); A. A. Gorman and M. A. J. Rodgers, *Chem. Phys. Lett.*, **120**, 58 (1985).
- 22) J. R. Hurst, S. L. Wilson, and G. B. Schuster, *Tetrahedron*, **41**, 2191 (1985).
- 23) The figures in parentheses show the wavelengths of emission or absorption of the intermediates as indicated in text.

(Received July 25, 1991)