

This article was downloaded by: [Laurentian University]

On: 08 December 2014, At: 07:04

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Toward the Development of Metal-Free Synthetic Nucleases: Cleavage of a Model Substrates by 1,4-Diazabicyclo[2.2.2]Octane Derivatives

Lyudmila S. Koroleva^a & Vladimir N. Silnikov^a

^a Novosibirsk Institute of Bioorganic Chemistry, Lavrentev Ave. 8, Novosibirsk, 630090, Russia

Published online: 23 Aug 2006.

To cite this article: Lyudmila S. Koroleva & Vladimir N. Silnikov (2004) Toward the Development of Metal-Free Synthetic Nucleases: Cleavage of a Model Substrates by 1,4-Diazabicyclo[2.2.2]Octane Derivatives, Nucleosides, Nucleotides and Nucleic Acids, 23:6-7, 989-992, DOI: [10.1081/NCN-200026052](https://doi.org/10.1081/NCN-200026052)

To link to this article: <http://dx.doi.org/10.1081/NCN-200026052>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Toward the Development of Metal-Free Synthetic Nucleases: Cleavage of a Model Substrates by 1,4-Diazabicyclo[2.2.2]Octane Derivatives

Lyudmila S. Koroleva* and Vladimir N. Silnikov

Novosibirsk Institute of Bioorganic Chemistry, Novosibirsk, Russia

ABSTRACT

Artificial ribonucleases of A_nBCL series were synthesized by solid-phase method. They consist of a hydrophobic alkyl radical A (n = 3–12 carbon atoms), an “RNA-binding domain” B (bisquaternary salt of 1,4-diazabicyclo[2.2.2]octane), a “catalytic domain” C (histidine residue) and a “linker” L that joins the domains B and C. The effect of the alkyl radical on the catalytic properties of the chemical catalyst was studied using three activated phosphate ester substrates: *p*-nitrophenyl phosphate, bis-*p*-nitrophenyl phosphate, and thymidine-3′-*p*-nitrophenyl phosphate.

Key Words: Metal-free artificial nucleases; *p*-nitrophenyl phosphate derivatives; 1,4-diazabicyclo[2.2.2]octane derivatives.

INTRODUCTION

There is much current interest in the design and development of small compounds that can mimic important enzymatic reactions under physiological conditions. One of the goals of these studies is the preparation of artificial ribonucleases, which cut a specific RNA at the target site and thus are applicable to molecular biology, therapy, and others.

*Correspondence: Lyudmila S. Koroleva, Novosibirsk Institute of Bioorganic Chemistry, Lavrentev Ave. 8, Novosibirsk 630090, Russia; E-mail: kls_2002@mail.ru.

Recently we have developed a series of artificial ribonucleases (ABLC series) containing four functional domains—catalytic domain C, linker group L, RNA-binding di- or polycation B and hydrophobic radical A.^[1] The optimal structure parameters and the role of B, C, L domains were determined in series of experiments.^[2] According to these results, still faster RNA hydrolysis should be possible if all these domains are presented in artificial ribonucleases. However the role and the optimal structure of hydrophobic radical A have remained not studied. In this study, we have synthesized a series of compounds (Fig. 1) with different alkyl residues to optimize the structure of this domain.




Figure 1. Solid-phase synthesis of artificial ribonucleases of ABCL series. 6a–h: a: $n = 2$, b: $n = 3$; c: $n = 5$; d: $n = 6$; e: $n = 8$; f: $n = 9$; g: $n = 11$; h: R—cyclohexyl.

RESULTS AND DISCUSSION

The synthesis of chemical ribonucleases is presented in Fig. 1. Merrifield resin was used as a polymer carrier. Treatment of Merrifield resin with excess N^{α} -Boc- N^{im} -DNP-L-Histidine and EDIPA results in compound 2. Subsequent attachment of the linker group was carried out by N-acylation of amino acid's α -amino groups by 4-bromobutyryl chloride. Compounds 5 were synthesized by alkylation of 1,4-diazabicyclo[2.2.2]octane using different alkyl halides. Mono-quaternary salts of 1,4-diazabicyclo[2.2.2]octane were joined to compound 4. Thus compounds differing in length of alkyl residue were obtained.

Kinetic studies were performed by using three activated phosphate ester substrates: *p*-nitrophenyl phosphate (*p*NPP), bis-*p*-nitrophenyl phosphate (BNPP), and thymidine-3'-*p*-nitrophenyl phosphate (TNPP) which is considered a reference substrate for the cleavage of phosphomono- and phosphodiesteres.^[3] Pseudo-first-order rate constants were determined for the three phosphate ester substrates.

Catalytic activity of RNase mimetics was determined on the surface of polymer carrier. Thus, the influence of micelle formation was excluded. The kinetic measurements were followed spectroscopically, monitoring the release of *p*-nitrophenolate ion at 400 nm. The rate constants were obtained by curve fitting of the initial data of the kinetic curve. Kinetics parameters were calculated by Origin program. Eq. 1 described experimental data in the best way.

$$A(t) = a + b \cdot \exp(t/k') \quad (1)$$

$A(t)$ —absorption of *p*-nitrophenolate at 400 nm, $k' = 1/k_{\text{obs}}$, a , b —constants.

The rate constants of the model substrates hydrolysis as a function of the alkyl fragment length are represented in Fig. 2. The hydrolysis efficiency by artificial ribonucleases depends on substrate structures. Hydrolysis rate of *p*-nitrophenylphosphate weakly depends on structure of alkyl residue of catalyst. The highest value of k_{obs} in this




Figure 2. Dependence hydrolysis rate on the number of carbons in alkyl residue in RNasemimetics.

experiment is observed for compounds A₆BLC and A₁₀BLC. In the case of bis-*p*-nitrophenylphosphate an increase of hydrolysis rate for all catalysts is observed. The influence of alkyl residue length becomes more marked. The most active derivatives are A₆BLC, A₉BLC and A₁₂BLC. The rate constants are changed in the range from $3.9 \cdot 10^{-4} \text{ min}^{-1}$ to $8.3 \cdot 10^{-4} \text{ min}^{-1}$. For thymidine-3'-*p*-nitrophenyl phosphate further growth of efficiency is observed, but in this case compound A₁₀BLC possesses maximum activity. The rate constants are changed in the range from $4.8 \cdot 10^{-4} \text{ min}^{-1}$ to $9.5 \cdot 10^{-4} \text{ min}^{-1}$.

Catalysts of A_nBLC series cleave phosphodiester bonds more efficiently than phosphomonoesters. Even in the case of simple model substrates, the influence of alkyl residue structure on efficiency of cleavage is displayed. In this study conditions excluding micelle formation are used, therefore it can be considered that the role of alkyl residues consists in promoting a suitable spatial complex structure based on hydrophobic interactions.

ACKNOWLEDGMENTS

This work is supported by RFBR (02-04-48664-a), Grants BRHE Rec-008, Wellcome Trust (CRIG 063630) and Russian Science Support Foundation.

REFERENCES

1. Konevets, D.A.; Beck, I.E.; Beloglazova, N.G.; Sulimenkov, I.V.; Sil'nikov, V.N.; Zenkova, M.A.; Shishkin, G.V.; Vlassov, V.V. Artificial ribonucleases: synthesis and RNA cleaving properties of cationic conjugates bearing imidazole residues. *Tetrahedron* **1999**, *55*, 503–512.
2. Konevets, D.A.; Mironova, N.L.; Beck, I.E.; Zenkova, M.A.; Shishkin, G.V.; Vlassov, V.V.; Silnikov, V.N. Chemical ribonucleases: 4. An analysis of the domain structure of chemical ribonucleases based on 1,4-diazabicyclo[2.2.2]octane. *Russ. J. Bioorgan. Chem.* **2002**, *28* (4), 331.
3. Madhavaiah, C.; Verma, S. Kinetic evaluation of metalated diglycine conjugate as a functional mimetic of phosphate ester hydrolase. *Bioconjug. Chem.* **2001**, *12*, 855–860.

Received December 2, 2003

Accepted April 23, 2004