

**284.** *A Study of the Polyhalides. Part III.  
Behaviour in Solution.*

By H. W. CREMER and DONALD R. DUNCAN.

THE polyhalides may be divided into three classes according to their behaviour in solution : (1) Those containing more than one iodine atom, which yield free iodine by dissociation and do not suffer appreciable hydrolysis : these have been carefully investigated by earlier workers and have therefore not been examined in detail again. (2) Those containing one iodine atom, including the salts of the series  $\text{RIBr}_2$ ,  $\text{RIBrCl}$ ,  $\text{RCl}_2$ , and  $\text{RCl}_4$ . In the solid state these yield iodine halides ( $\text{IBr}$ ,  $\text{ICl}$ , or  $\text{ICl}_3$ ) by dissociation. The iodine halides are decomposed by water, and the polyhalides are hydrolysed similarly with formation of iodate, the aqueous solutions containing

little or no iodine halide. The authors have investigated the nature of the equilibria established and the reactions of the solutions in some detail, there being comparatively little published data on the subject. (3) Those containing no iodine; these yield bromine, chlorine, or bromine chloride by dissociation, and suffer hydrolysis, as is shown by their acid reaction. The solutions contain considerable quantities of free halogen.

The experiments now described are concerned mainly with solutions of the second class. They appear to indicate that the main equilibrium established in solution is  $5\text{IBr}_2' + 3\text{H}_2\text{O} \rightleftharpoons 2\text{I}_2 + \text{IO}_3' + 10\text{Br}' + 6\text{H}'$  for a dibromiodide, and similarly for a dichloriodide,  $5\text{IBrCl}' + 3\text{H}_2\text{O} \rightleftharpoons 2\text{I}_2 + \text{IO}_3' + 5\text{Br}' + 5\text{Cl}' + 6\text{H}'$  for a chlorobromiodide, and  $5\text{ICl}_4 + 9\text{H}_2\text{O} \rightleftharpoons \text{I}_2 + 3\text{IO}_3' + 20\text{Cl}' + 6\text{H}'$  for a tetrachloriodide. The establishment of such an equilibrium may be explained as follows, potassium dibromiodide being taken as an example. The compound may be assumed to ionise primarily to  $\text{K}' + \text{IBr}_2'$ ; the anion then dissociates in the same way as in the dry state, *viz.*,  $\text{IBr}_2' \rightleftharpoons \text{Br}' + \text{IBr}$ , and the iodine bromide suffers hydrolysis,  $\text{IBr} + \text{HOH} \rightleftharpoons \text{IOH} + \text{H}' + \text{Br}'$ , the hypoiodous acid finally decomposing in the usual way:  $5\text{IOH} \rightleftharpoons 2\text{I}_2 + \text{H}' + \text{IO}_3' + 2\text{H}_2\text{O}$ . On keeping a solution, a state of equilibrium between the products is established. In this particular case the degree of hydrolysis is small and insufficient to cause precipitation of iodine, but with some polyhalides such precipitation does occur on standing, as in the case of the dichloriodides observed by Reade (J., 1929, 853). Whilst these equations represent the main equilibria, it is probable that traces of other substances may be present (*e.g.*,  $\text{HIBr}_2$ ,  $\text{BrI}_2'$ , etc.). Solutions of chlorobromiodides may contain the ions  $\text{IBr}_2'$  and  $\text{ICl}_2'$  as well as  $\text{IBrCl}'$ ; this might be expected on theoretical grounds, and would account for the tendency of caesium dibromiodide to separate on recrystallisation of the chlorobromiodide. Such complications render it inadvisable to attempt to determine any equilibrium constants, but the relative degrees of hydrolysis may be estimated as described on p. 2040.

## EXPERIMENTAL.

### *Reactions in Solution.*

When a reagent is added to a solution of a polyhalide, if it is capable of reacting with one of the mols. or ions present, thereby removing it from the solution, the equilibrium will be displaced. Hence, towards such a reagent a polyhalide may react as if it were not hydrolysed, whereas, towards a substance which reacts with one of the products of hydrolysis, it may react as if it were completely hydrolysed. Thus the behaviour with different reagents may be used in order to establish the presence or absence of the various ions in a polyhalide solution. In this way, evidence has been obtained for

the presence in  $\text{KIBr}_2$  aq. of  $\text{K}'$ ,  $\text{IBr}_2'$ ,  $\text{I}_2$ ,  $\text{Br}'$ ,  $\text{HIO}$ ,  $\text{IO}_3'$ , and  $\text{H}'$ , and for the almost complete absence of  $\text{K}'''$ ,  $\text{Br}_2$ ,  $\text{IBr}$ ,  $\text{I}'$ ,  $\text{HBrO}$ ,  $\text{BrO}_3'$ , and complex kations such as  $\text{KIBr}'$  and  $\text{KBr}_2'$ . All the reactions investigated confirm the view that the hydrolysis follows the course described above.

(1) *Evidence for the Existence of the Ordinary Metallic Kations in Solution.*—In solutions of a polyhalide the metal gives its normal reactions; thus,  $\text{K}_2\text{SiF}_6$  and  $\text{KClO}_4$  are pptd. as readily from  $\text{KI}_3$  aq. or  $\text{KIBr}_2$  aq. as from  $\text{KI}$  or  $\text{KBr}$  aq. Similarly, simple salts of the bivalent metals may be pptd. from solutions of their polyhalides. None of the reactions affords any evidence for the existence of metallic kations of abnormal valency or of complexes such as  $\text{KIBr}'$ . That the alkali metals are present in solutions of their polyhalides entirely as univalent kations is further supported by their absorption spectra and by the quantitative data given later (p. 2040).

(2) *Evidence for the Existence of Polyhalide Anions.*—Sparingly sol. polyhalides are pptd. whenever a solution of a polyhalide of the same series is added to one of a salt of the metal or base forming the insol. derivative: e.g.,  $\text{KIBr}_2 + \text{NMe}_4\text{NO}_3 = \text{KNO}_3 + \text{NMe}_4\text{IBr}_2(\text{ppt.})$ . Solutions of the compositions  $\text{HIBr}_2$ ,  $\text{HICl}_2$ , and  $\text{HICl}_4$  act as precipitants of  $\text{IBr}_2'$ ,  $\text{ICl}_2'$ , and  $\text{ICl}_4'$  respectively. These double decomp. reactions make it probable that the polyhalide solutions and the corresponding free acids contain the above anions, a conclusion which is confirmed by the absorption spectra and other quant. investigations. Details of these, including evidence for  $\text{IBrCl}'$ , are given later (p. 2040).

(3) *Evidence for the Existence of Free Halogens in Polyhalide Solutions.*—It is well known that solutions of polybromides contain much free  $\text{Br}_2$ , while those of polyiodides contain a small proportion of their  $\text{I}_2$  in the free state. This is shown by the high bromine v. p. of the polybromide solutions, and by the fact that halogens may be extracted from the solutions by, e.g.,  $\text{CCl}_4$ . When this is shaken with an  $M/20$ -solution of  $\text{KIBr}_2$ ,  $\text{KClBrI}$ ,  $\text{KICl}_2$ , or  $\text{KICl}_4$ ,  $\text{I}_2$ , but no appreciable quantity (less than 1%) of  $\text{Br}_2$  or  $\text{Cl}_2$  and no acid passes into the  $\text{CCl}_4$ . Of the substances produced during the hydrolysis of these polyhalides,  $\text{I}_2$  is evidently the only one appreciably sol. in  $\text{CCl}_4$  which is present in more than traces. Since  $\text{Br}_2$ ,  $\text{Cl}_2$ ,  $\text{IBr}$ ,  $\text{ICl}$ , and  $\text{ICl}_3$  are all readily sol. in  $\text{CCl}_4$ , the polyhalide solutions can contain very little of these substances. This is true even of the solutions obtained by shaking  $\text{IBr}$  and  $\text{ICl}$  with  $\text{H}_2\text{O}$ . Table I gives the compositions of the different phases at equilibrium, when 1 g.-mol. of  $\text{IBr}$  is shaken with 2 l. of  $\text{CCl}_4$  and 20 l. of  $\text{H}_2\text{O}$  at  $25^\circ$ . Table II gives the corresponding figures for  $\text{ICl}$ .

TABLE I.

	$\text{I}_2$ .	$\text{Br}_2$ .
Concn. in $\text{H}_2\text{O}$ ...	0.025 <i>N</i>	0.050 <i>N</i>
Concn. in $\text{CCl}_4$ ...	0.227 <i>N</i>	0.000 <i>N</i>
Solid phase .....	100%	0%


TABLE II.

	$\text{I}_2$ .	$\text{Cl}_2$ .
Concn. in $\text{H}_2\text{O}$ ...	0.011 <i>N</i>	0.050 <i>N</i>
Concn. in $\text{CCl}_4$ ...	0.227 <i>N</i>	0.000 <i>N</i>
Solid phase .....	100%	0%


By shaking conc. solutions of the polyhalides with a much larger proportion of  $\text{CCl}_4$ , it may be shown that traces of the other halogens are extracted. Nevertheless, for the purpose of the quant. work described on p. 2040, this small trace may be ignored.

(4) *Evidence for the Existence of Monohalide Ions.*—In ordinary analytical work, the monohalide anions are usually tested for by adding a solution of a

salt of a metal which forms insol. halides. When such tests are performed on solutions of polyhalides, they give the reactions of monohalides; but the halogens are not present entirely as monohalide ions and solutions of  $\text{IBr}_2'$ ,  $\text{ClBrI}'$ ,  $\text{ICl}_2'$ , and  $\text{ICl}_4'$  contain no  $\text{I}'$  ions. This is in accordance with the above theory, and is supported by the following observations: (a)  $\text{HgCl}_2$  gives no ppt. with solutions of these polyhalides. (b) No ppt. is obtained on mixing sat. solutions of  $\text{KIBr}_2$  and  $\text{Pb}(\text{NO}_3)_2$ , although  $\text{PbBr}_2$  is pptd. from more dil. solutions. (c) When  $N/10\text{-AgNO}_3$  is run slowly into a large excess of  $N/10\text{-KIBr}_2$ , the Ag is first pptd. quantitatively as pure AgBr (Found: 0.1857 g. Calc.: 0.1867 g.). Soon, however, free  $\text{I}_2$  begins to be pptd. simultaneously, attaining a max. when about 2 mols. of  $\text{AgNO}_3$  per mol. of  $\text{KIBr}_2$  have been added. The  $\text{I}_2$  is destroyed by an excess of  $\text{AgNO}_3$ , the reaction being complete when 3 mols. of  $\text{AgNO}_3$  have been added. The resultant solution contains 1 mol. of  $\text{KNO}_3$  and 2 mols. of  $\text{HNO}_3$  and the ppt. is a mixture of AgBr, AgI, and  $\text{AgIO}_3$ . These facts are in agreement with the equation



The course of the reaction may be explained as follows: in a solution of a dibromiodide there is an equilibrium



Addition of  $\text{AgNO}_3$  ppts. AgBr; if  $\text{AgIBr}_2$  exists at all, it must be considerably more sol. than AgBr, otherwise it would be pptd.; AgI is not pptd., since there is no  $\text{I}'$  present, and  $\text{AgIO}_3$  is not pptd. because it is 60 times as sol. as AgBr and the concn. of  $\text{IO}_3'$  ion in the solution is only 0.1 of that of the  $\text{Br}'$  ion. The removal of  $\text{Br}'$  from the solution causes the equilibrium to move to the right. Eventually a point is reached when the concn. of  $\text{I}_2$  attains the sat. value, and thereafter it is pptd. along with the AgBr. 2 Mols. of  $\text{AgNO}_3$  are required to ppt. all the  $\text{Br}'$ ; then  $\text{AgIO}_3$  is pptd., 0.2 mol. of  $\text{AgNO}_3$  being required per mol. of original  $\text{KIBr}_2$ , whilst the  $\text{I}_2$  is destroyed by the  $\text{AgNO}_3$ ,  $3\text{I}_2 + 6\text{Ag}' + 3\text{H}_2\text{O} = 5\text{AgI} + 6\text{H}' + \text{AgIO}_3$ , 0.8 mol. of  $\text{AgNO}_3$  being required per mol. of  $\text{KIBr}_2$ . Summation of these equations shows that the completed reaction is represented by the equation already given.

$\text{AgNO}_3$  reacts similarly with  $\text{KClBrI}$  and  $\text{KICl}_2$ . In the former case, the ppt. produced in the first stages of the reaction consists of pure AgBr in quant. yield (Found: 0.0833, 0.0837 g. Calc.: 0.0831 g.); AgCl is not pptd. at first, since it is nearly 15 times as sol. as AgBr.

With  $\text{KICl}_4$  the reaction follows a similar course:  $3\text{KICl}_4 + 15\text{AgNO}_3 + 6\text{H}_2\text{O} = 12\text{AgCl} + \text{AgI} + 2\text{AgIO}_3 + 3\text{KNO}_3 + 12\text{HNO}_3$ . The different stages of the reaction were established quantitatively and follow exactly the same lines as in the case of  $\text{KIBr}_2$ . The odour of the solutions suggests that some substance such as  $\text{HIO}$  or  $\text{HIO}_2$  may be present in appreciable quantity in the intermediate stages of the reactions (see Taylor, J., 1913, **103**, 31). Its presence, however, would not be expected to affect the nature of the final products.

In the above reactions with  $\text{AgNO}_3$ ,  $\text{H}_2\text{O}$  takes part, and the action is evidently one between the  $\text{Ag}'$  ions and the hydrolysis products of the polyhalide, rather than with the polyhalide itself. It follows that in non-aqueous solution, the action, if it occurs at all, must follow a different course. In a solvent such as MeOH, no hydrolysis can occur, and there is no evidence for alcoholysis with formation of  $\text{MeBr}$  and  $\text{IO}_3'$ . Thus, solutions of  $\text{KIBr}_2$  and

$\text{AgNO}_3$ , both in  $\text{MeOH}$ , give a ppt. of  $\text{AgBr}$  and  $\text{AgI}$ , free from  $\text{AgIO}_3$ , and much  $\text{I}_2$  remains in solution even after excess of  $\text{AgNO}_3$  has been added. The authors have investigated in more detail the reaction between  $M/20$ -solutions of  $\text{AgNO}_3$  and  $\text{IBr}$  in  $\text{MeOH}$ , a simpler reaction which follows a similar course. When 1 mol. of  $\text{AgNO}_3$  is run in drop by drop with constant stirring, the  $\text{Ag}$  and  $\text{Br}$  are completely pptd. as  $\text{AgBr}$  (Found : 0.1645, 0.1657 g. Calc. : 0.1643 g.), so that the solution has the composition of  $M/40$ -iodine mononitrate :  $\text{IBr} + \text{AgNO}_3 = \text{AgBr} + \text{I}(\text{NO}_3)$ . The solution is pale orange, but on evaporation at room temp. under reduced pressure it leaves a white solid residue. The product is very deliquescent and is decomposed by  $\text{H}_2\text{O}$  with formation of free  $\text{I}_2$ ,  $\text{HNO}_3$ , and  $\text{HIO}_3$ , probably according to the equation  $5\text{INO}_3 + 3\text{H}_2\text{O} = 2\text{I}_2 + \text{HIO}_3 + 5\text{HNO}_3$ . Addition of another 0.5 mol. of  $\text{AgNO}_3$  to the  $\text{MeOH}$  solution ppts. the  $\text{Ag}$  quantitatively as  $\text{AgI}$  (Found : 0.1755, 0.1748, 0.1755 g. Calc. : 0.1755 g.), and the solution, which is colourless, has a composition corresponding to that of iodine trinitrate :  $2\text{INO}_3 + \text{AgNO}_3 = \text{AgI} + \text{I}(\text{NO}_3)_3$ . Excess of  $\text{AgNO}_3$  produces no further change. Although the compositions of the solutes are those of  $\text{I}(\text{NO}_3)$  and  $\text{I}(\text{NO}_3)_3$ , it cannot be stated with certainty that these two nitrates are actually present, although this seems the simplest explanation of the results. Kappeler (*Ber.*, 1911, **44**, 3496) states that a nitrate or basic nitrate of trivalent  $\text{I}$  is formed when  $\text{HNO}_3$  acts on  $\text{HIO}_3$  under suitable conditions, but he gives no analysis. The high velocity and smoothness of the reaction and the nature of the products suggest that the reaction is ionic, and hence that  $\text{I}$  is capable of forming uni- and ter-valent kations in a non-hydrolysing strongly ionising solvent such as  $\text{MeOH}$  :  $\text{IBr} \rightleftharpoons \text{I}' + \text{Br}'$ ;  $\text{Ag}' + \text{Br}' = \text{AgBr}$ ;  $2\text{I}' \rightleftharpoons \text{I}'' + \text{I}'''$ ;  $\text{Ag} + \text{I}' = \text{AgI}$  (see also p. 2039).

(d)  $\text{TiNO}_3$  gives with  $\text{KIBr}_2$  aq. a ppt. of  $\text{TiBr}$ , free  $\text{I}_2$  being liberated, and the solution giving the reactions of a thallic salt. There is no evidence for the formation of either  $\text{IO}_3'$  or  $\text{I}'$ , and the reaction is probably essentially a reduction of the  $\text{IBr}_2'$  to  $\text{Br}'$  and free  $\text{I}_2$ ,  $2\text{IBr}_2' + \text{Ti}' = \text{Ti}''' + 4\text{Br}' + \text{I}_2$  (compare p. 2036).

(5) *Evidence for the Existence of Halogen Oxyacids or their Anions.*—There is no evidence for the existence of any oxy-acids of  $\text{Cl}$  or  $\text{Br}$  in solutions of polyhalides of the second class (p. 2031), although traces of  $\text{ClO}_3'$  or  $\text{BrO}_3'$  may be formed when their solutions are made alkaline. Reade (*J.*, 1929, 853) has obtained evidence for the presence of  $\text{HIO}$  in solutions of dichloriodides. When a solution of any iodine polyhalide is made alkaline, it contains a high concn. of  $\text{IO}'$ , since the presence of  $\text{OH}'$  ion shifts the equilibrium in this direction (see below). Some polyhalide solutions give a slight ppt. of  $\text{Ba}(\text{IO}_3)_2$  with  $\text{BaCl}_2$  aq. indicating the presence of  $\text{IO}_3'$  in the solution, and di- and tetra-chloriodides sometimes give iodates on attempts to crystallise them from water. Iodate is also formed when the equilibrium is disturbed by the removal of halide or  $\text{H}'$  ions (p. 2034 and below) or of free  $\text{I}_2$  by means of an org. solvent (p. 2040).

(6) *Evidence for the Existence of Hydrogen Ions.*—Solutions of trihalides containing one  $\text{I}$  atom, but not those containing more than one, show a strong acid reaction to indicators; this is in accordance with the hydrolysis theory already advanced (p. 2032); e.g., the acidity of solutions of  $\text{IBr}_2'$  is due to partial hydrolysis :  $5\text{IBr}_2' + 3\text{H}_2\text{O} \rightleftharpoons 2\text{I}_2 + \text{IO}_3' + 10\text{Br}' + 6\text{H}'$ . On adding an excess of  $\text{NaHCO}_3$  to  $\text{KIBr}_2$  aq., the  $\text{H}'$  ions are removed and the hydrolysis is carried to completion. The reaction is approx.  $5\text{KIBr}_2 + 6\text{NaHCO}_3 =$

$3\text{H}_2\text{O} + 6\text{CO}_2 + 2\text{I}_2 + \text{IO}_3' + 10\text{Br}' + 5\text{K}' + 6\text{Na}'$ : it was found that 1.18 mols. of  $\text{NaHCO}_3$  were required to destroy 1 mol. of  $\text{KIBr}_2$  (theory, 1.20) and that the solution contained considerable  $\text{Br}'$ , a smaller concn. of  $\text{IO}_3'$ , and traces only of  $\text{I}_2$ ,  $\text{IO}'$ , and  $\text{BrO}_3'$ , free  $\text{I}_2$  being pptd. and  $\text{CO}_2$  evolved. A similar reaction occurs when  $\text{NaOH}$  aq. or  $\text{KOH}$  aq. is run into  $\text{KIBr}_2$  aq.,  $\text{I}_2$  being pptd. and then redissolved by excess of alkali, the resultant solution containing  $\text{Br}'$ ,  $\text{I}'$ ,  $\text{IO}'$ , and  $\text{IO}_3'$ . On acidifying such a solution, the polyhalide is re-formed. Similar reactions occur with other dibromiodides, and with chlorobromo-, dichloro-, and tetrachloro-iodides. With  $\text{NH}_3$ , impure  $\text{NI}_3$  is pptd. (J., 1930, 2750).

(7) *Action of Reducing Agents on Polyhalide Solutions.*—When a solution of a polyhalide containing 1 I at. is acted upon by a reducing agent, it is first reduced to bromide or chloride, the  $\text{I}_2$  being first precipitated in the free state (e.g.,  $2\text{KIBr}_2 + 2\text{H} = 2\text{KBr} + 2\text{HBr} + \text{I}_2$ ), and then reduced by an excess of a sufficiently strong reducing agent to  $\text{HI}$ . The intermediate pptn. of  $\text{I}_2$  is bound to occur, since monoiodides react with these polyhalides with liberation of  $\text{I}_2$ , e.g.,  $\text{KI} + \text{KIBr}_2 = 2\text{KBr} + \text{I}_2$ ; excess of iodide brings about the formation of polyiodide; in the case of organic polyhalides the polyiodide is often sparingly sol. and remains undissolved. Polyhalide solutions (e.g.,  $\text{KIBr}_2$  aq.) are reduced in the above way by nascent  $\text{H}_2$  (quantitatively),  $\text{Mg}$  powder (even in the absence of acid),  $\text{SO}_2$  (quantitatively in dilute sol.:  $2\text{KIBr}_2 + 2\text{H}_2\text{O} + \text{SO}_2 = 2\text{KBr} + 2\text{HBr} + \text{I}_2 + \text{H}_2\text{SO}_4$ ;  $\text{I}_2 + 2\text{H}_2\text{O} + \text{SO}_2 = 2\text{HI} + \text{H}_2\text{SO}_4$ ),  $\text{H}_2\text{S}$ ,  $\text{Fe}''$  salts, ferrocyanides (oxidised to ferriyanides), etc. Nitrites precipitate  $\text{I}_2$ , slowly if no acid is added but immediately in the presence of acid, presumably according to the equation  $\text{NaNO}_2 + 2\text{KIBr}_2 + \text{H}_2\text{O} = \text{NaNO}_3 + 2\text{KBr} + \text{I}_2 + 2\text{HBr}$ . The reduction does not proceed further in this case, since mono- and poly-iodides are oxidised by acid solutions of nitrites, e.g.,  $2\text{NaNO}_2 + 2\text{KI}_3 + 4\text{HX} = 2\text{NaX} + 2\text{KX} + 2\text{H}_2\text{O} + 2\text{NO} + 3\text{I}_2$ .

The reactions between polyhalides and  $\text{Na}_2\text{S}_2\text{O}_3$  are of particular interest owing to the importance of the tri-iodide-thiosulphate reaction in volumetric analysis. In an ordinary 'iodine' titration, the reaction may be represented by  $\text{I}_3' + 2\text{S}_2\text{O}_3'' = 3\text{I}' + \text{S}_4\text{O}_6''$ . When, on the other hand, free bromine or tribromide solutions are titrated directly with  $\text{Na}_2\text{S}_2\text{O}_3$ , this is oxidised almost entirely to sulphate. Solutions of dibromo- and tetrachloro-iodides give intermediate results, the relative proportions of  $\text{S}_4\text{O}_6''$  and  $\text{SO}_4''$  formed varying largely according to the rate of titration and other conditions.

The quant. oxidation of  $\text{Na}_2\text{S}_2\text{O}_3$  to  $\text{Na}_2\text{S}_4\text{O}_6$  by means of a dibromiodide is represented by the equation (a)  $\text{IBr}_2' + 2\text{S}_2\text{O}_3'' = \text{I}' + 2\text{Br}' + \text{S}_4\text{O}_6''$ , strictly analogous to that for the  $\text{I}_3'$  reaction, and the quant. oxidation to  $\text{SO}_4''$  would follow the equation (b)  $4\text{IBr}_2' + \text{S}_2\text{O}_3'' + 5\text{H}_2\text{O} = 4\text{I}' + 8\text{Br}' + 2\text{SO}_4'' + 10\text{H}'$ . When  $\text{KIBr}_2$  aq. is run slowly into  $\text{Na}_2\text{S}_2\text{O}_3$  aq., 99% of the  $\text{KIBr}_2$  reacts according to equation (a), but when the order of addition is reversed, only about 50% so reacts. If, however, the  $\text{KIBr}_2$  aq. is first saturated with  $\text{KBr}$ , in order to suppress its hydrolysis, 94–97% reacts according to (a). This suggests that the  $\text{IBr}_2'$  ion reacts in the same way as the  $\text{I}_3'$  ion, oxidising the  $\text{Na}_2\text{S}_2\text{O}_3$  to  $\text{Na}_2\text{S}_4\text{O}_6$ , and that it is the  $\text{HIO}_3$  or  $\text{HIO}$  formed by hydrolysis which causes the formation of  $\text{SO}_4''$ . Since the addition of acid also tends to suppress the hydrolysis of dibromiodides, one would expect less  $\text{SO}_4''$  to be formed in an acidified solution, an effect which has been confirmed experimentally. Thus, in a solution containing 8% of  $\text{HCl}$ , reaction (a)

occurred to the extent of 52%, while in one containing 25% of HCl it occurred to the extent of 68%. When the  $\text{Na}_2\text{S}_2\text{O}_3$  aq. is run drop by drop into the  $\text{KIBr}_2$  aq. in the absence of added bromide or acid, with constant stirring, the reaction occurs in well-marked stages. In the first,  $\text{I}_2$  is pptd. and  $\text{SO}_4''$  and acid are formed, apparently according to the equation  $8\text{IBr}_2' + \text{S}_2\text{O}_3'' + 5\text{H}_2\text{O} = 16\text{Br}' + 4\text{I}_2 + 2\text{SO}_4'' + 10\text{H}'$ , the reaction probably occurring *via* the hydrolysis products of the  $\text{KIBr}_2$ . The quantity of  $\text{IBr}_2'$  destroyed and that of acid formed, per c.c. of  $\text{Na}_2\text{S}_2\text{O}_3$  aq. run in, agree with those required by this equation during the early stages of the reaction. The subsequent reaction consists merely in the redissolution of the  $\text{I}_2$  as polyiodide, which oxidises the  $\text{S}_2\text{O}_3''$  to  $\text{S}_4\text{O}_6''$  in the ordinary way, no more sulphate being formed. Although the reaction occurs in stages, the final result is the same as if the reactions (a) and (b) occurred simultaneously and to the same extent,  $4\text{IBr}_2'$  reacting according to (a) and  $4\text{IBr}_2'$  according to (b).

With  $\text{ICl}_4'$  solutions, the reaction producing  $\text{S}_4\text{O}_6''$  occurred to the extent of about 40% (mean of several experiments), and with tribromide or free bromine solutions to about 7%. The exact proportion of  $\text{SO}_4''$  and  $\text{S}_4\text{O}_6''$  formed, however, varies with the method of titration.

In order to determine a polyhalide by  $\text{Na}_2\text{S}_2\text{O}_3$  titration, an excess of KI aq. should first be added to convert all the polyhalide into polyiodide, which can then be titrated quantitatively.

(8) *Action of Oxidising Agents on Polyhalide Solutions.*—Powerful oxidising agents such as acid  $\text{KMnO}_4$  oxidise solutions of dibromiodides and dichloro- and tetrachloro-iodides to iodate, with liberation of  $\text{Br}_2$  or  $\text{Cl}_2$ , but no  $\text{I}_2$ : e.g.,  $\text{KIBr}_2 + 3\text{O} = \text{KIO}_3 + \text{Br}_2$ .

(9) *Action of Halogens on Polyhalide Solutions.*—Halogens may act on polyhalides in aq. solution, causing either addition, substitution, or oxidation of the type just mentioned. For instance,  $\text{Cl}_2$  passed through an aq. solution of a dichloroiodide, first forms tetrachloroiodide, which is converted by excess into  $\text{IO}_3'$ : e.g.,  $\text{KICl}_2 + \text{Cl}_2 = \text{KICl}_4$ ;  $\text{KICl}_4 + \text{Cl}_2 + 3\text{H}_2\text{O} = \text{KIO}_3 + 6\text{HCl}$  (Part I, J., 1931, 1864). Dibromiodides react similarly after the  $\text{Br}_2$  has been liberated (e.g.,  $\text{KIBr}_2 + \text{Cl}_2 = \text{KICl}_2 + \text{Br}_2$ ). From polyiodides  $\text{I}_2$  is liberated. The action of  $\text{Br}_2$  is similar to that of  $\text{Cl}_2$  in all these cases, but  $\text{I}_2$  has little or no action. The reaction  $\text{KIBr}_2 + 2\text{Br}_2 + 3\text{H}_2\text{O} = \text{KIO}_3 + 6\text{HBr}$  was found to be reversible. On mixing  $\text{KIBr}_2$  aq. and  $\text{Br}_2$  aq. in the proportion indicated by this equation, the colour fades considerably but not completely, and the solution gives the reactions of both  $\text{IO}_3'$  and  $\text{IBr}_2'$ . A similar solution is obtained on adding  $\text{HBr}$  aq. to  $\text{KIO}_3$  aq. When a large excess of  $\text{Br}_2$  is used, the product no longer gives the reactions of a dibromiodide (e.g., a ppt. of  $\text{I}_2$  with  $\text{NaHCO}_3$  and of  $\text{NI}_3$  with  $\text{NH}_3$ ), but it gives those of  $\text{IO}_3'$  (e.g., a ppt. with  $\text{BaCl}_2$  aq., not given by the  $\text{KIBr}_2$  aq.) and of  $\text{Br}'$ . Also, as indicated in Table III, the obs. solubility of  $\text{Br}_2$  in  $\text{KIBr}_2$  solutions of different concn., at  $25^\circ$ , agrees with the assumption that oxidation to  $\text{IO}_3'$  is almost complete in solutions satd. with  $\text{Br}_2$ .

TABLE III.

Molarity of $\text{KIBr}_2$ solution .....	0.00	0.013	0.0415	0.063
Solubility of $\text{Br}_2$ (mols./l.), obs. ....	0.20	0.27	0.48	0.58
Solubility of $\text{Br}_2$ (mols./l.), calc. ....	0.20	0.27	0.46	0.58

In calculating the values, it is necessary to allow for the high sol. of  $\text{Br}_2$  in the  $\text{HBr}$  formed during the reaction. The experimental data do not provide any



evidence concerning the existence of tetrabromiodides in aqueous solution; they are unknown in the solid state.

(10) *Action of Stabilisers on Polyhalide Solutions.*—Theoretically, if to a solution of a polyhalide we add one of the substances formed by its hydrolysis (*e.g.*, in the case of a dibromiodide, free  $I_2$ ,  $IO_3'$ ,  $H'$ , or  $Br'$ ), the equilibrium, as given on p. 2032, will be shifted towards the left, *i.e.*, hydrolysis will be suppressed, and the reactions which are due to the presence of hydrolysis products should be inhibited. Owing to its sparing solubility, it is impossible to increase the concn. of  $I_2$  appreciably. The stabilising action of  $IO_3'$  is not very pronounced in the case of a trihalide, since only one  $IO_3'$  ion is produced for every 5 polyhalide ions hydrolysed. The stabilising influence of acidification is far more pronounced, as would be expected from theory, and the influence of acid on the reaction between  $IBr_2'$  and  $S_2O_3''$  is also in accordance with the belief that addition of  $H'$  ions suppresses hydrolysis, just as their removal (addition of alkali) brings about hydrolysis. The stabilising action of acid may also be due, in part, to formation of stable unionised mols. of hydrogen polyhalide, *e.g.*,  $HICl_2$  (see Table V). The addition of monohalide also has a considerable suppressing effect on the hydrolysis, as shown by the results of work on the  $CCl_4$  equilibrium (p. 2041), the absorption spectrum (see below), thiosulphate reaction (p. 2036), etc. Dichloriodide solutions which deposit  $I_2$  on standing are generally stable in the presence of a sufficient excess of  $Cl'$ , while dibromiodide solutions which have been saturated with  $KBr$  do not give a ppt. of  $I_2$  even when made neutral or alkaline, an effect due partly to the suppression of hydrolysis by the  $Br'$  ions and partly to the high solubility of  $I_2$  in  $KBr$  aq. (presumably due to the formation of the ion  $I_2Br'$ ). Addition of  $K$  salts, other than  $KIO_3$  or halide, to a solution of a  $K$  polyhalide has no apparent effect on its absorption spectrum and chemical properties, showing that the  $K$  atom does not enter into the hydrolysis equilibrium.

#### Absorption Spectra.

During our work on these absorption spectra, papers on the subject were published by Gilbert, Goldstein, and Lowry (*J.*, 1931, 1092) and by Gillam and Morton (*Proc. Roy. Soc.*, 1931, *A*, **132**, 152). Since we had independently arrived at almost the same conclusions as these investigators, it is not necessary to discuss the matter in detail, but we record our numerical data in Table IV.

TABLE IV.

$\lambda$ in $\mu\mu$ .	700	650	600	570	550	540	530	520	510			
1. $KIBr_2$ in sat. aq. $KBr$	0.1	0.1	0.6	1.6	2.6	4.2	6.2	8.6	11.5			
2. Dibromiodides .....	1.0	1.1	1.4	3.2	6.2	10.0	12.1					
3. $C_5H_5NIBr$ in $MeOH$	1.3	2.1	3.3	5.7	8.6	11.1						
4. $IBr$ in $MeOH$ .....	1.6	1.8	3	7	10							
5. $KIBr_2$ in aq. starch soln. ....	8	13	26	70	123	165						
6. $IBr$ in $CCl_4$ .....	10	18	28	68	126	168						
$\lambda$ in $\mu\mu$ .	530	520	510	500	490	480	470	460	450	440	430	425
7. $KIBr_2$ in $H_2O$	13	15	18	21	29	47	64	77				
8. $IBr$ in $MeOH$	34	50	77	96	139	179	226	266	289	296		
9. $IBr$ in $CCl_4$ ...	210	275	333	359	357	351	341	300	244	191	138	116

The figures give the value of  $k$  in the equation  $kcd = \log_{10} I_0/I$ . The results were obtained visually, in Expts. 6 and 7 by means of a revolving sector apparatus and in the remainder by using a Nutting photometer. In Expts. 1—4



the solutions were  $M/20$ ; but in the others, more dil. In these dil. solutions the value of  $k$  is not altered greatly by variation of the concn. Within the limits of experimental error, the same values (Expt. 2) were obtained for the dibromiodides of  $K$ ,  $NH_4$ , and  $C_5H_5N \cdot H$  in both  $MeOH$  and  $H_2O$ . The identity in absorption spectrum of the different dibromiodides confirms the view already expressed that the metal or base does not form part of the complex ion, and that the degree of dissociation of the different polyhalides of the same series is independent of the cation. Gilbert, Goldstein, and Lowry observed a difference in intensity of ultra-violet absorption between solutions in  $H_2O$  and in  $EtOH$ , which they attributed to hydrolysis in the former solvent, although the present work does not bear out their ideas as to the nature of the hydrolysis. For instance, the action  $H_2O + RI_3 \rightleftharpoons ROH + HI_3$  does not occur when  $ROH$  is a strong base, while the equilibrium  $ICl_2' \rightleftharpoons I' + Cl_2$  is impossible owing to the action of  $Cl_2$  on the  $I'$  ion (an irreversible reaction producing  $Cl'$  ions and free  $I_2$ ) and is further untenable since the present work indicates definitely that the aq. solutions of dichloriodides contain free  $I_2$  but no free  $Cl_2$ . The fact that no difference is observable in the visible region between solutions of dibromiodides in  $H_2O$  and those in  $MeOH$  may be attributed to similarity in absorption spectrum between the dibromiodide ions and the products of hydrolysis, and to the small degree of hydrolysis of  $IBr_3'$ .

Solvation probably occurs in  $H_2O$  and in  $MeOH$  but not in  $CCl_4$ . The similarities and differences in spectrum are accounted for satisfactorily by assuming the following equilibria, which, from general considerations, would be expected to be established:

(1)  $KIBr_2$  in sat. aq.  $KBr$ : all hydrolysis and dissociation suppressed by the bromide ion, the chief absorbent being  $IBr_2'$ .

(2a) Dibromiodides in  $H_2O$ : see p. 2032.

(2b) Dibromiodides in  $MeOH$ :  $IBr_2' \rightleftharpoons Br' + IBr$ , and the equilibria given in (4) below.

(3)  $C_5H_5NIBr$  in  $MeOH$ :  $C_5H_5NIBr \rightleftharpoons C_5H_5N + IBr$ ; possibly also  $C_5H_5NIBr \rightleftharpoons C_5H_5N'I' + Br'$  (compare Cremer and Duncan, J., 1931, 1861, 2250; Williams, *ibid.*, p. 2783). When this solution is mixed with an equimol. quantity of  $AgNO_3$  in  $MeOH$ , quantitative pptn. of  $AgBr$  occurs, which, however, would be expected according to both these modes of dissociation.

(4)  $IBr$  in  $MeOH$ :  $IBr \rightleftharpoons I' + Br'$ ;  $Br' + IBr \rightleftharpoons IBr_2'$ ; and, to a small extent,  $2IBr \rightleftharpoons I_2 + Br_2$ .

These equilibria must, presumably, also be established in all the above cases where  $IBr$  is formed in  $MeOH$  aq. by dissociation. The simple reaction with  $AgNO_3$  (p. 2034) depends on the insolubility of  $AgBr$ , the equilibria being displaced by removal of  $Br'$  ions.

(5)  $KIBr_2$  and starch: the intense coloration is probably analogous to the blue colour obtained with a tri-iodide and starch.

(6)  $IBr$  in  $CCl_4$ :  $2IBr \rightleftharpoons I_2 + Br_2$ , the ionic reactions not occurring in this solvent.

The free acids  $HIBr_2$ ,  $HICl_2$ , and  $HICl_4$  give absorption spectra similar to those of their salts.

Solutions of  $IBr$  in different solvents show a variation in colour similar to that shown by  $I_2$  solutions, due, no doubt, to variation in solvation and in the equilibrium constants of the above reactions. The solutions in general show

one broad absorption band, and are quite different from the superimposed absorption spectra of  $I_2$  and  $Br_2$ . There is a gradual change in the position of max. absorption, from the green for non-solvating solvents such as ligroin,  $C_6H_6$ ,  $CCl_4$  (500  $\mu\mu$ ),  $C_6H_4Me_2$ , and tetrahydronaphthalene, through the blue for solvents such as  $CCl_4$ , ethylene dichloride and dibromide, and  $C_6H_5\cdot NO_2$ , to the violet for solvating solvents such as glycerol,  $Et_2O$ ,  $AcOH$ ,  $COMe_2$ ,  $EtOH$ , and  $MeOH$  (420  $\mu\mu$ ). For the dibromiodides, where the  $IBr$  is definitely combined, the max. absorption is at 390  $\mu\mu$ , while for the compound with  $C_5H_5N$  it is in the ultra-violet.

*Equilibria between Aqueous Solutions of Polyhalides and Carbon Tetrachloride.*

The effect of shaking  $CCl_4$  with an aq. solution of a polyhalide has already been mentioned (p. 2033). The concn. of  $I_2$  in the  $CCl_4$  at equilibrium is a measure of the relative degrees of hydrolysis of the different polyhalides of the type under investigation, a high value indicating a relatively high concn. of free  $I_2$  in the aq. layer and consequently considerable hydrolysis. For a polyhalide which undergoes considerable hydrolysis with pptn. of  $I_2$ , the addition of a small quantity of  $CCl_4$  merely dissolves some of the pptd.  $I_2$  and does not disturb the equilibrium. In the case of polyhalides which do not of themselves give a ppt. of  $I_2$ ,  $CCl_4$  causes an increase in the degree of hydrolysis, but if the temp., concn. of the aq. solution, and proportion of  $CCl_4$  are fixed, the concns. of  $I_2$  in the  $CCl_4$  in equilibrium with different polyhalides will be in the same order as the degrees of hydrolysis. In the cases given in Table V, the  $CCl_4$  was shaken with 10 times its volume of an  $M/20$ -aqueous solution of each polyhalide at 25° until equilibrium was reached, 3 days sufficing. Col. 1 gives the composition of the solute; the properties of the solution, including the equilibrium with  $CCl_4$ , are independent of the mode of prep. and depend only on the composition and temp., so it is possible to obtain data for solutions of compounds which cannot be isolated in the solid state (e.g.,  $NaIBrCl$  and the free acids). Col. 2 gives the % of the total  $I$  which remains in the aq. layer, col. 3 the % present in the  $CCl_4$  and col. 4 the % pptd. as solid  $I_2$  (calc. by diff.). Col. 5 gives the % hydrolysis, which is assumed to take place according to the simple equations on p. 2032, the intermediate products being neglected; there is, however, probably a considerable concn. of  $HIO$  in the solutions.

TABLE V.

1.	2.	3.	4.	5.	1.	2.	3.	4.	5.
$HICl_4$ .....	96	4	0	13	$NaIBrCl$ .....	53	45	2	61
$KICl_4$ .....	96	4	0	13	$IBr$ .....	50	45	5	65
$ICl_3$ .....	91	9	0	24	$HICl_2$ .....	46	45	9	70
$KIBr_2$ .....	78	22	0	29	$KICl_2$ .....	36	45	19	83
$HIBr_2$ .....	77	23	0	30	$NaICl_2$ .....	34	45	21	86
$NH_4IBr_2$ .....	75	25	0	33	$NH_4ICl_2$ .....	33	45	22	87
$HIBrCl$ .....	64	36	0	47	$ICl$ .....	23	45	32	100
$KIBrCl$ .....	55	45	0	59					

Within the limits of experimental error the figures are the same for all polyhalides belonging to the same series, showing that the metal does not enter into the equilibrium in dil. solution (compare J., 1931, 2249; this vol., p. 2039). These results agree with qual. observations of the stabilities of the solutions; thus,  $M/20$ -solutions of the tetrachloro- and dibromo-iodides,  $ICl_3$ , and the free acids  $HICl_4$ ,  $HIBr_2$ , and  $HIBrCl$  are stable, whereas those of

chlorobromo- and dichloro-iodides deposit  $I_2$  on standing,  $IBr$  and  $ICl$  also being decomposed with separation of  $I_2$ . The polyhalides are all hydrolysed to a smaller extent than the iodine halides from which they are derived; this may also be expressed by the statement that the hydrolysis of the iodine halides is suppressed by increasing the concn. of monohalide ions. This observation is in agreement with the assumption that stable complex ions are formed consisting of a compound of the iodine halide with monohalide ions. It is very difficult on any other assumption to account for this effect, and in particular to account for the fact that chlorides suppress the hydrolysis of  $IBr$ ; although this effect is not very great with dilute solutions, it becomes very pronounced with conc. solutions of chlorides. Neutral salts, other than halides and iodates, have no effect on the hydrolysis of iodine halides, while acids suppress hydrolysis, the suppressing action of acid and iodate being less with the more stable compounds.

The difference between the two chlorides of iodine as regards behaviour towards water is very striking:  $ICl_3$  dissolves to form a clear solution which may be kept indefinitely without separation of  $I_2$ , whereas  $ICl$  is almost completely hydrolysed. In the former case it is possible that hydrolysis may occur without liberation of  $I_2$  (e.g.,  $ICl_3 + H_2O \rightleftharpoons OICl + 2HCl$ , or  $ICl_3 + 2H_2O \rightleftharpoons HIO_2 + 3HCl$ ). In the above investigation, owing to the very small concn. of halogen found in the  $CCl_4$  layer in the case of  $ICl_3$  and the tetrachloroiodides, it is difficult to prove that only  $I_3$  was extracted, as was done in the other cases, and it is possible that the solutions contain  $ICl_3$ , as such or in an ionised condition. There is some evidence for the presence of trivalent iodine kations: thus  $ICl_3$  aq. gives with  $H_2SO_4$  a ppt. which is said to be iodine sulphate (Mellor, "Comprehensive Treatise, etc.," 1922, Vol. II, p. 121). On the other hand, Faull and Baekström (*J. Amer. Chem. Soc.*, 1932, **54**, 620) have shown by a simple transport method that a solution of  $ICl$  in  $HCl$  aq. contains no iodine kations, the  $I$  being present in the anion. Repetition of this experiment with  $ICl_3$  might give interesting results. The difference between  $ICl_3$  and the tetrachloroiodides, on the one hand, and  $ICl$  and the dichloroiodides, on the other, may be due to the existence in aq. solutions of the former of relatively stable trivalent iodine kations, whereas univalent iodine kations are not stable in aqueous solution. Most of the known salts of  $I$  contain this element in the trivalent state.

### Summary.

The polyhalides may be divided into three classes according to their behaviour in aqueous solution, viz., (1) compounds formed from free iodine and monohalides; their solutions contain polyhalide ions, monohalide ions, and free iodine, but no hydrolysis products (iodate, acid, etc.); (2) compounds of iodine halides with monohalides; their solutions contain polyhalide ions, monohalide ions, and the products resulting from the hydrolysis of the iodine halides, but no appreciable concentration of iodine halide itself; (3) compounds of bromine, chlorine, or bromine chloride with monohalides; their solutions contain polyhalide ions, monohalide ions, free halogens, and the hydrolytic products of the last (e.g., bromic acid).

The nature of the equilibria existing in aqueous solutions of

polyhalides of the second class has been established ; their reactions have been examined in detail and measurements made of absorption spectra and of the degree of hydrolysis as indicated by partition experiments with carbon tetrachloride.

Evidence is given for the existence, at least in methyl-alcoholic solution, of iodine mononitrate,  $\text{INO}_3$ , and iodine trinitrate,  $\text{I}(\text{NO}_3)_3$ , and of the ions  $\text{I}'$  and  $\text{I}'''$ . The possibility of the existence of these ions in aqueous solution is also considered.

The nature of the equilibria in solutions of iodine monobromide and its additive compounds ( $\text{KIBr}_2$ ,  $\text{C}_5\text{H}_5\text{NIBr}$ , etc.) in various solvents is also discussed.

KING'S COLLEGE, LONDON.

[Received, May 20th, 1932.]

---