






**Synthesis from D-Glucose of 1,5-Dideoxy-1,5-imino-L-fucitol, a Potent  $\alpha$ -L-Fucosidase Inhibitor**George W. J. Fleet,<sup>a\*</sup> Antony N. Shaw,<sup>a</sup> Stephen V. Evans,<sup>b</sup> and Linda E. Fellows<sup>b\*</sup><sup>a</sup> Dyson Perrins Laboratory, Oxford University, South Parks Road, Oxford OX1 3QY, U.K.<sup>b</sup> Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3DS, U.K.

1,5-Dideoxy-1,5-imino-L-fucitol (**1**), synthesised from methyl  $\alpha$ -D-glucopyranoside, is a potent competitive inhibitor of the hydrolysis of *p*-nitrophenyl  $\alpha$ -L-fucopyranoside catalysed by  $\alpha$ -L-fucosidase (ex. bovine epididymis) causing 50% inhibition of enzymic activity at  $2.5 \times 10^{-8}$  M.

Several polyhydroxylated piperidines and pyrrolidines have been shown to be competitive inhibitors of glycosidases from many sources and are proving useful biochemical tools; several derivatives of 5-amino-5-deoxyglucose (nojirimycin) and of 5-amino-5-deoxymannose have been used as glucosidase and mannosidase inhibitors.<sup>1,2</sup> To date, no compound of this class having fucosidase inhibitory activity has been described; yet glycans containing both D-fucose and L-fucose (**2**) are widespread in nature. In particular,  $\alpha$ -L-fucose is the immunodominant sugar of many complex carbohydrate antigens, and the L-fucose content of some animal glycans is known to change under certain pathological conditions, such as transformation to tumorigenesis.<sup>3</sup> Specific inhibitors of  $\alpha$ -L-fucosidase are likely to find wide application, not only in the investigation of the structure/function relationships of fucose containing glycans, but also in understanding the pathology of inherited disorders characterised by a deficiency of  $\alpha$ -L-fucosidase.<sup>4</sup> This paper reports the synthesis of 1,5-dideoxy-1,5-imino-L-fucitol (**1**) from methyl  $\alpha$ -D-glucopyranoside (**3**); (**1**) is shown to be a very potent competitive inhibitor of bovine epididymis  $\alpha$ -L-fucosidase, but to have no inhibitory action on a range of other glycosidases.

The synthesis of (**1**) from (**3**) requires inversion of configuration at C-2 and C-3, deoxygenation of C-6, and the formation of the piperidine ring between C-1 and C-5 with inversion of configuration at C-5. The protected altrose (**4**), prepared from (**3**) by standard procedures,<sup>5</sup> was benzylated [(benzyl bromide, sodium hydride, tetrabutylammonium iodide in tetrahydrofuran (THF))] to give (**5**), m.p. 91–92 °C (lit.<sup>6</sup> 90–91 °C), in 84% yield. Hydrolysis of the benzylidene acetal by acetic acid:water (4:1) gave diol (**6**) which underwent selective esterification of the primary hydroxy group with toluene-*p*-sulphonyl chloride in pyridine at –20 °C to form (**7**),<sup>†</sup>  $[\alpha]_D^{20} +52^\circ$  (c 0.70, CHCl<sub>3</sub>), in 75% yield.



<sup>†</sup> Satisfactory spectral and/or analytical data were obtained for all new compounds.

Reduction of (7) with lithium aluminium hydride in THF to (8), followed by benzylation of the remaining free hydroxy group, gave methyl 6-deoxy-2,3,4-tri-*O*-benzyl- $\alpha$ -D-altropyranoside (9),  $[\alpha]_{\text{D}}^{20} +81^\circ$  (*c* 0.84,  $\text{CHCl}_3$ ), in 64% yield. Hydrolysis of (9) by trifluoroacetic acid: water (4:1), followed by reduction with sodium borohydride in ethanol, gave the protected 6-deoxy-D-altritol (10), m.p. 74.5–75.5 °C,  $[\alpha]_{\text{D}}^{20} +7.9^\circ$  (*c* 0.88,  $\text{CHCl}_3$ ), in 85% yield [38% yield from (4)]. Conversion into the bis(methanesulphonate) (11) [3 equiv. methanesulphonyl chloride in pyridine, 0 °C], followed by treatment with tetrabutylammonium azide in dimethylformamide (DMF) gave azidomethanesulphonate (12) in 60% yield,  $\nu_{\text{max}}$ . 2095  $\text{cm}^{-1}$  (azide) and  $^1\text{H}$  n.m.r. ( $\text{CDCl}_3$ ) showing H-5 as a quartet of doublets at  $\delta$  5.1. Hydrogenation of (12) in the presence of palladium catalysts gave a mixture of products in which some hydrogenolysis of the benzyl ethers accompanied reduction of the azide; however, treatment with sodium hydrogen telluride<sup>7</sup> smoothly transformed (12) directly to the required piperidine (13),  $[\alpha]_{\text{D}}^{20} -42^\circ$  (*c* 0.80,  $\text{CHCl}_3$ ), in 75% yield. Removal of the benzyl protecting groups from (13) by hydrogenolysis in the presence of palladium black in ethanol gave (1);<sup>‡</sup> the  $^1\text{H}$  n.m.r. spectra of (13) in  $\text{CDCl}_3$  and of (1) in  $\text{D}_2\text{O}$  show that both compounds are in a chair conformation.

The inhibitory action of (1) on the hydrolysis of the corresponding nitrophenyl glycopyranosides catalysed by  $\alpha$ -glucosidase (yeast),  $\beta$ -glucosidase (almonds),  $\alpha$ -galactosidase (green coffee beans),  $\beta$ -galactosidase (*Aspergillus niger*),

$\alpha$ -mannosidase (Jack Bean),  $\beta$ -xylosidase (*Aspergillus niger*), and  $\alpha$ -L-fucosidase was determined. § A concentration of (1) of only  $2.5 \times 10^{-8}$  M was sufficient to cause 50% inhibition of  $\alpha$ -L-fucosidase-catalysed hydrolysis of *p*-nitrophenyl  $\alpha$ -L-fucopyranoside; a Lineweaver–Burk plot shows that (1) is a competitive inhibitor ( $K_i$   $4.8 \times 10^{-9}$  M). In contrast, none of the other enzymes was appreciably inhibited at a concentration of (1) of  $5 \times 10^{-4}$  M. Should this specificity be maintained over a wide range of mammalian enzymes, 1,5-dideoxy-1,5-imino-L-fucitol (1) is likely to prove a research tool of exceptional usefulness.

Received, 9th April 1985; Com. 479

## References

- 1 R. C. Bernotas and B. Ganem, *Tetrahedron Lett.*, 1985, **26**, 1123; T. Niwa, T. Tsuruoka, H. Goi, Y. Kodama, J. Itoh, S. Inouye, Y. Yamada, T. Niida, M. Nobe, and Y. Ogawa, *J. Antibiot.*, 1984, **37**, 1579; Y. Kodama, T. Tsuruoka, T. Niwa, and S. Inouye, *ibid.*, 1985, **38**, 116; S. V. Evans, A. R. Hayman, L. E. Fellows, T. K. M. Shing, A. E. Derome, and G. W. J. Fleet, *Tetrahedron Lett.*, 1985, **26**, 1465.
- 2 S. V. Evans, L. E. Fellows, T. K. M. Shing, and G. W. J. Fleet, *Phytochemistry*, 1985, in the press.
- 3 H. M. Flowers, *Adv. Carbohydr. Chem. Biochem.*, 1981, **39**, 279.
- 4 P. R. Dorling, in 'Lysosomes in Biology and Pathology,' ed. J. T. Dingle, R. T. Dean, and W. Sly, Elsevier, Amsterdam, 1984, p. 347.
- 5 N. K. Richtmeyer, *Methods Carbohydr. Chem.*, 1962, **1**, 107.
- 6 C. L. Stevens, J. P. Dickerson, K. G. Taylor, P. Blumbergs, and P. M. Pillai, *J. Org. Chem.*, 1975, **40**, 2468.
- 7 H. Suzuki and K. Takaoka, *Chem. Lett.*, 1984, 1733.

§ The enzymes and nitrophenyl glycopyranoside substrates were obtained from Sigma. Details of the enzyme assay procedures are given in ref. 2.

‡ Spectroscopic data for (1): an oil,  $[\alpha]_{\text{D}}^{20} -48.8^\circ$  (*c* 0.64,  $\text{H}_2\text{O}$ );  $M + \text{H}^+$  148 ( $\text{NH}_3$ -chemical ionisation);  $^1\text{H}$  n.m.r. (300 MHz) in  $\text{D}_2\text{O}$   $\delta$  0.94 (d,  $\text{CH}_3$ ), 2.22 (dd,  $\text{H}_{1a}$ ), 2.92 (dd,  $\text{H}_{1c}$ ), 3.55 (m,  $\text{H}_2$ ), 3.32 (dd,  $\text{H}_3$ ), 3.64 (m,  $\text{H}_4$ ), 2.67 (qd,  $\text{H}_5$ );  $J(1e, 1a)$  13.0,  $J(1a, 2)$  11.0,  $J(1e, 2)$  5.4,  $J(2, 3)$  9.7,  $J(3, 4)$  3.1,  $J(4, 5)$  1.2,  $J(5, \text{Me})$  6.8 Hz;  $^{13}\text{C}$  n.m.r. (125 MHz) in  $\text{D}_2\text{O}$   $\delta$  75.61 (d,  $\text{CHOH}$ ), 73.06 (d,  $\text{CHOH}$ ), 68.20 (d,  $\text{CHOH}$ ), 53.94 (d,  $\text{CHN}$ ), 49.25 (t,  $\text{CH}_2\text{N}$ ), 16.70 (q,  $\text{CH}_3$ ).