This article was downloaded by: [Heriot-Watt University] On: 06 January 2015, At: 12:36 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

# Microwave-Assisted Efficient One-Pot Synthesis of Nitriles From Aldehydes in the Presence of $P_2O_5/SiO_2$ in Solvent-Free Media

Hossein Eshghi<sup>a</sup> & Zinat Gordi<sup>b</sup>

<sup>a</sup> Department of Chemistry , Ferdowsi University of Mashhad , Mashhad, Iran

<sup>b</sup> Department of Chemistry , Sistan and Baluchestan University , Zahedan, Iran Published online: 21 Dec 2010.

To cite this article: Hossein Eshghi & Zinat Gordi (2005) Microwave-Assisted Efficient One-Pot Synthesis of Nitriles From Aldehydes in the Presence of  $P_2O_5/SiO_2$  in Solvent-Free Media, Phosphorus, Sulfur, and Silicon and the Related Elements, 180:2, 619-623, DOI: <u>10.1080/104265090517460</u>

To link to this article: <u>http://dx.doi.org/10.1080/104265090517460</u>

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u> Phosphorus, Sulfur, and Silicon, 180:619–623, 2005 Copyright © Taylor & Francis Inc. ISSN: 1042-6507 print / 1563-5325 online DOI: 10.1080/104265090517460

# Microwave-Assisted Efficient One-Pot Synthesis of Nitriles From Aldehydes in the Presence of $P_2O_5/SiO_2$ in Solvent-Free Media

**Hossein Eshghi** 

Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, Iran

### Zinat Gordi

Department of Chemistry, Sistan and Baluchestan University, Zahedan, Iran

A rapid and efficient procedure is developed for a one-pot synthesis of nitriles by condensation of aldehydes with hydroxylamine hydrochloride in the presence of  $P_2O_5/SiO_2$  in solvent-free media under microwave irradiation.

Keywords Aldehydes; aldoximes; microwave irradiation; nitriles; phosphorus pentoxide

# INTRODUCTION

Nitriles, important reagents for organic synthesis have been known to chemist for a long time. In recent reports, it was shown that nitriles are very useful starting materials for the synthesis of various bioactive molecules.<sup>1,2</sup> Furthermore, nitriles can be converted to amides, carboxylic acids, amines, ketones, and esters.<sup>3</sup> For example, vanilly-lamine readily obtained by the reduction of vanillylonitrile,<sup>4</sup> could be reacted with acyl chlorides to give capsaicinoids.

Several procedures are available for the one-step conversion of aldehydes into nitriles using different chemical reagents and hydroxylamine hydrochloride.<sup>5.6</sup> However, most the methods suffer from serious drawbacks which include the use of hazardous and expensive or commercially unavailable reagents, long reaction times, low yields, drastic reaction conditions, and tedious workup procedure.

The efficiency of microwave irradiation (MW) in organic synthesis is currently under intensive study.<sup>7</sup> There are just a few publications

Received April 6, 2004; in final form July 29, 2004.

Address correspondence to Hossein Eshghi, Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, 91774-1436, Iran. E-mail: heshghi@hamoon.usb.ac.ir

describing the use of microwave irradiation for preparations of nitriles from aldehydes and hydroxylamine hydrochloride.<sup>8</sup> Our projects have been to develop new synthetic methods using the  $P_2O_5/SiO_2$  in solvent-free media.<sup>9</sup> Having the above facts in mind, we wish to report a very simple and efficient method for one-pot synthesis of nitriles from both alkyl and aryl aldehydes using  $P_2O_5/SiO_2$  and hydroxylamine hydrochloride in solvent-free media under microwave irradiation.

## **RESULTS AND DISCUSSION**

In order to prepare nitriles, various types of aldehydes were mixed with hydroxylamine hydrochloride in the presence of  $P_2O_5/SiO_2$  reagent using microwave irradiation in solvent-free media. In this approach, nitriles were obtained instead of oximes. The general reaction is illustrated in Scheme 1 and the results are reported in Table I. All reactions were performed in one minute. As shown in the Table I several structurally different aldehydes underwent cleanly and remarkably fast the one-pot reaction to the corresponding nitriles. This mild and versatile method which results in yields of 75–95% can also be applied to aldehydes containing other reactive functional groups, such as double bond and hydroxyl group. Even 2-furaldehyde gives a good yield of product in spite of the literature<sup>10</sup> which claims that such a reaction was not successful.

$$\text{RCHO} + \text{NH}_2\text{OH} \cdot \text{HCl} \xrightarrow[\text{MW, 1 min.}]{\text{P}_2\text{O}_5/\text{SiO}_2} \text{RCN} + \text{H}_2\text{O}$$

SCHEME 1 One-pot preparation of nitriles.

In the absence of microwave irradiation, after 10 min grinding of the mixture the corresponding aldoximes were exclusively formed in good yields.<sup>9d,e</sup> This clearly indicates that aldoxime may be an intermediate in the conversion of aldehydes to nitriles. Also, aldoximes are dehydrated to the corresponding nitriles in high yields under the same conditions. The general reaction is illustrated in Scheme 2 and the

SCHEME 2 Conversion of aldoximes to nitriles.

Entry	Aldehyde	Nitrile	Yield (%) <sup>a</sup>	Mp or bp (°C)/ mmHg(lit.)
1	$CH_3CHO$	$CH_3CN$	75	81/760 (80/760) <sup>11</sup>
<b>2</b>	C <sub>3</sub> H <sub>7</sub> CHO	$C_3H_7CN$	78	$113/760\ (110/760)^{11}$
3	$C_5H_{11}CHO$	$C_5H_{11}CN$	78	$160/760\ (162/760)^{11}$
4	$C_6H_{13}CHO$	$C_6H_{13}CN$	75	198/760 (70-71/10) <sup>12</sup>
5	Сно	CN CN	95	190/760 (187/760) <sup>11</sup>
6	ноСно	но-См	80	$113\ (112114)^{10}$
7	ноСно	но-См	89	82-83 (81) <sup>13</sup>
	H <sub>3</sub> CO	H₃CO		
8	(H <sub>3</sub> C) <sub>2</sub> N-CHO	(H <sub>3</sub> C) <sub>2</sub> N-CN	85	74–75 (74–76) <sup>12</sup>
9	O <sub>2</sub> N-CHO		85	146–147 (148) <sup>11</sup>
10	Сно	CN	85	$116117~(117)^{11}$
	O <sub>2</sub> N	O <sub>2</sub> N		
11	СН=СН-СНО	CH=CH-CN	85	$258/760\ (260/760)^{11}$
12	Сно	CN CN	75	$147\ (147{-}148)^{12}$

TABLE I One-Pot Preparation of Nitriles from Aldehydes by Using  $P_2O_5/SiO_2$  and Hydroxylamine Hydrochloride in Solvent-Free Media under Microwave Irradiation

<sup>*a*</sup>IR, <sup>1</sup>H-NMR, and mp or bp confirmed the structures.

results are reported in Table II. As shown in Table II the corresponding nitriles were obtained in 85-95% yields from both alkyl and aryl aldoximes.

In conclusion, this one-pot synthesis of nitriles from aldehydes, without isolation of aldoximes, offers significant improvements over the existing procedures and thus makes available a variety of nitriles and related compounds. Also this simple and readily available reagent affords various nitriles in a short reaction time, with excellent yields,

Entry	Aldoxime	Nitrile	Yield (%) <sup>a</sup>	Mp or bp (°C)/ mmHg(lit.)
1	CH <sub>3</sub> CH=NOH	$CH_3CN$	85	81/760
2	C <sub>3</sub> H <sub>7</sub> CH=NOH	$C_3H_7CN$	88	$(80/760)^{11}$ 113/760 (110/760) <sup>11</sup>
3	C <sub>6</sub> H <sub>13</sub> CH=NOH	$C_6H_{13}CN$	85	(110/760)11 198/760
4		CN/CN	90	$(70-71/10)^{12}$ 190/760 $(187/760)^{11}$
5	но	HO-CN	95	$113 \\ (112-114)^{10}$
6	но	но-СМ	95	82-83 (81) <sup>13</sup>
	H <sub>3</sub> CO	H <sub>3</sub> CO		
7	(H <sub>3</sub> C) <sub>2</sub> N-C=NOH		92	$74-75 \\ (74-76)^{12}$
8		CN CN	85	$\begin{array}{c} 147 \\ (147  148)^{12} \end{array}$

TABLE II Conversion of Aldoximes to Nitriles by Using  $P_2O_5/SiO_2$  in Solvent-Free Media under Microwave Irradiation

<sup>*a*</sup>IR, <sup>1</sup>H-NMR, and mp or bp confirmed the structures.

and good selectivity. Further applications of the  $P_2O_5/SiO_2$  reagent in organic synthesis are under investigation.

### **EXPERIMENTAL**

All melting points recorded are uncorrected open capillary measurements. IR spectra were recorded on a Shimadzu-IR 470 spectrophotometer. <sup>1</sup>H-NMR spectra were recorded on a Bruker-80 and 500 MHz instrument using tetramethylsilane (TMS) as an internal standard. Silica gel 60(230–400 mesh) was obtained from Fluka and was dried in an oven at 120°C for 2 h. Irradiation was carried out in a domestic microwave oven (Electra, 2450 MHz, and 800 W) for an optimized time. The  $P_2O_5/SiO_2$  reagent was obtained according to the early reported procedure.<sup>9</sup>

#### **Preparation of Nitriles**

#### General Procedure

In a typical reaction, a mixture of aldehyde (2 mmol), hydroxylamine hydrochloride (4 mmol) and the  $P_2O_5/SiO_2$  reagent (1 g), or a mixture of aldoxime (2 mmol) and the  $P_2O_5/SiO_2$  reagent (1 g) was ground thoroughly in the mortar. Usually an immediate colour change was observed. The mortar was covered with a watch glass and put inside the microwave oven. The mixture was irradiated for one minute and the completion of the reaction was monitored by TLC. After the completion of the reaction, the mortar was removed from the oven, the mixture was cooled to room temperature and then 10 ml of 5% aqueous HCl was added to the mixture. The resulting solution was extracted with  $CH_2Cl_2$  (2 × 10 ml). The extracts were combined and dried over  $CaCl_2$ . Evaporation of the solvent under vacuum gave nitriles with high purity (based on TLC, <sup>1</sup>H-NMR, IR and melting point). Column chromatography or recrystalization from benzene or benzene-cyclohexane gave pure products.

### REFERENCES

- I. K. Khanna, R. M. Weier, Y. Yu, X. D. Xu, F. J. Koszyk, P. W. Collins, C. M. Koboldt, A. W. Veenhuizen, W. E. Perkins, J. J. Casler, J. L. Masferrer, Y. Y. Zhang, S. A. Gregory, K. Seibert, and P. C. Isakson, *J. Med. Chem.*, 40, 1634 (1997) and literatures citied therein.
- [2] M. Chihiro, H. Nagamoto, I. Takemura, K. Kitano, H. Komatsu, K. Sekiguchi, F. Tabusa, T. Mori, M. Tominnaga, and Y. Yabuuchi, J. Med. Chem., 38, 353 (1995) and literatures citied therein.
- [3] R. F. Smith, J. A. Albright, and A. M. Waring, J. Org. Chem., 31, 4100 (1966).
- [4] J. March, Advanced Organic Chemistry, John Wiley & Sons, New York, pp. 918–919 (1992) and literatures citied therein.
- [5] E. C. Wang and G. J. Lin, *Tetrahedron Lett.*, 39, 4047 (1998) and literatures citied therein.
- [6] K. Tanaka, Solvent-Free Organic Synthesis, Wiley-VCH Verlag Gmbh & Co. KGaA, Weinheim, pp. 294–298 (2003).
- [7] a) S. Caddick, *Tetrahedron*, **51**, 10403 (1995); b) A. Loupy, A. Petit, J. Hamelin, F. Texier-Boullet, P. Jacquault, and D. Mathe, *Synthesis*, 1213 (1998).
- [8] P. Lidstrom, J. Tierney, B. Wathey, and J. Westman, *Tetrahedron*, 57, 9225 (2001).
- [9] a) H. Eshghi, M. Rafie, and M. H. Karimi, Synth. Commun., **31**, 771 (2001); b) H. Eshghi, and Z. Gordi, Synth. Commun., **33**, 2971 (2003); c) H. Eshghi, M. Rafie, Z. Gordi, and M. Bohloli, J. Chem. Research (S), 763 (2003); d) Z. Gordi, M. Sc. Thesis, University of Sistan and Baluchestan, Zahedan, Iran (2000); e) H. Eshghi and Z. Gordi, Phosphorus, Sulfur, and Silicon (2005) in press; f) H. Eshghi and P. Shafieyoon, Phosphorus, Sulfur, and Silicon, **179**, 2149 (2004).
- [10] J. C. Feng, B. Liu, L. Dai, and N. S. Bian, Synth. Commun., 28, 3765 (1998).
- [11] H. Sharghi, M. H. Sarvari, Tetrahedron, 58, 10323 (2002).
- [12] F. E. Chen, H. Fu, G. Meng, Y. Cheng, and Y. X. Lu, Synthesis, 1519 (2000).
- [13] D. G. Desai, S. S. Swami, G. D. Mahale, Synth. Commun., 30, 1623 (2000).