

1,2,5-ortho esters of D-arabinose as versatile arabinofuranosidic building blocks. Concise synthesis of the tetrasaccharidic cap of the lipoarabinomannan of *Mycobacterium tuberculosis*

Toufiq Bamhaoud,^a Sylvie Sanchez^b and Jacques Prandi^{*b}

^a Laboratoire de Chimie Organique et Bioorganique, Université Chouaib Doukkali, BP 20, El Jadida, Maroc

^b Institut de Pharmacologie et de Biologie Structurale du CNRS, 205 route de Narbonne, 31077 Toulouse, France.
E-mail: prandi@ipbs.fr

Received (in Cambridge, UK) 1st February 2000, Accepted 7th March 2000

Published on the Web 31st March 2000


1,2,5-ortho esters of D-arabinose were found to be ideally suited building blocks for the stereoselective formation of α and β arabinofuranosidic linkages after nucleophilic opening of the orthoester with oxygen and sulfur nucleophiles; the tetrasaccharidic cap of the lipoarabinomannan of *Mycobacterium tuberculosis* was then synthesized in a highly convergent manner.

Lipoarabinomannans (LAMs) are highly antigenic polysaccharidic compounds isolated from the cell wall of various species of mycobacteria.¹ They share common structural features, a phosphatidyl *myo*-inositol anchor and a mannan core substituted by an arabinan domain, exclusively composed of D-arabinofuranosidic units.¹ At the ends of this arabinan domain, are found β -D-arabinofuranosides substituted on position 5 by small motifs, called caps, variable on the mycobacterial species.^{2–5} Strong modulations of the biological properties of the LAMs were observed for LAMs isolated from different species of mycobacteria and these variations have been tentatively correlated to the structure of these caps.^{2,4–8}

In light of the importance of these motifs for the biological properties of the LAMs in relation with the immunopathogenicity of *M. tuberculosis*, we now report the expeditious synthesis of **1** (Scheme 1) the major tetrasaccharide found at the ends of the LAMs of *M. tuberculosis* and *M. bovis* BCG in a form suitable for further conjugation with a protein.⁹ Orthoesters **3** and **11** were found to be highly versatile precursors for the elaboration of the two different arabinofuranosidic rings and the construction of the crucial β -arabinofuranosidic linkage. The result is a rapid and convergent synthesis of **1**.

3-*O*-benzyl 1,2,5-benzylidene D-arabinofuranose **3** was readily obtained in 82% yield from the 1,2,5-benzylidene D-arabinofuranose **2** { $[\alpha]_D^{25} -28$ (*c* 0.94, CHCl₃) lit.¹⁰ +28 for the enantiomer} after treatment with benzyl bromide and sodium hydride in dimethylformamide (Scheme 1). SnCl₄ catalyzed opening of orthoester **3** with 5 equiv. of methyl 9-hydroxynonanoate⁹ in CH₂Cl₂ gave the 2-*O*-benzoyl- α -D-arabinofuranoside **4** as the only isolated product in 76% yield. Complete regio- and stereo-selectivity were observed for the double orthoester opening¹¹ and none of the isomeric 5-*O*-benzoyl arabinofuranosidic compound was observed with this catalyst. In an analogous manner, ethyl 2-*O*-benzoyl-3-*O*-benzyl-1-thio- α -D-arabinofuranoside **5** was obtained in 80% yield from the opening of **3** with 1.1 equiv. of ethanethiol under SnCl₄ catalysis, once again with complete stereocontrol of the orthoester opening.^{12,13} Unmasking of the hydroxy group on the 2-position of the arabinofuranosidic ring of thioglycoside **5** for later attachment of the *p*-methoxybenzyl tether¹⁴ was carried out in two steps and gave alcohol **7** in 73% overall yield (Scheme 1).


We also looked at the reactions of 3-*O*-benzyl 1,2,5-dichloroethylidene D-arabinofuranose **11**, available in 90% yield (BnBr, NaH, DMF) from known compound **10**, prepared in only two steps from D-arabinose.¹⁵ As was expected from electronic factors, nucleophilic opening of the orthoester function of **11**



Scheme 1 Reagents and conditions: i, NaH, BnBr, DMF, room temp., 1 h, 82%; ii, 5 equiv. methyl 9-hydroxynonanoate, 0.1 equiv. SnCl₄, MS 4 Å, CH₂Cl₂, room temp., 15 min, 76%; iii, 1.1 equiv. EtSH, 0.1 equiv. SnCl₄, MS 4 Å, CH₂Cl₂, -18 °C, 30 min, 80%; iv, 1.5 equiv. Bu^tPh₂SiCl, 1.5 equiv. imidazole, DMF, room temp., 1 h, 92%; v, 0.1 equiv. MeONa, MeOH, room temp., 79%; vi, 2.5 equiv. BEMP, 2.5 equiv. *p*-MeOBnBr, MeCN, 0 °C to room temp., 1 h, 70%. vii, NaH, BnBr, DMF, room temp., 1 h, 90%; viii, 5 equiv. HOCH₂(CH₂)₇C(O)OMe, 2 equiv. SnCl₄, MS 4 Å, CH₂Cl₂, room temp., 3 h, then MeONa, MeOH, room temp., 1 h, 60%; ix, 1.5 equiv. Bu^tCOCl, 0.1 equiv. DMAP, pyridine, 0 °C, 1.5 h, 80%.

proved to be more difficult than for **3**. 2 Equiv. of SnCl₄ at room temperature were necessary to bring the reaction of **11** with methyl 9-hydroxynonanoate to completion and a mixture of 2-*O*- and 5-*O*-dichloroacetyl α -arabinofuranosides was obtained. α -Arabinofuranoside **12** was isolated in 60% yield after deacylation with sodium methoxide. Selective pivaloylation of the primary hydroxy group of **12** (0 °C, Bu^tCOCl, DMAP, pyridine) gave alcohol **13** ready for glycosylation (80%). Opening of **11** with ethanethiol was also obtained with 1.2 equiv. of EtSH and 0.2 equiv. SnCl₄ at 0 °C, deacylation of the mixture of dichloroacetates as above gave **9** in 64% overall yield from **11**. Selective silylation of the primary position of **9** with TBDPSCl and imidazole afforded 73% of the previously prepared **7**. Finally, the 4-methoxybenzyl group was introduced on the 2-position of **7** with 2.5 equiv. of 4-methoxybenzyl bromide¹⁶ and 2.5 equiv. of 2-*tert*-butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine¹⁷ (BEMP) in acetonitrile to give thioarabinofuranoside **8** in 70% yield.

Coupling of the arabinofuranosidic units and elaboration of the β -arabinofuranosidic linkage were accomplished according to Ogawa's internal aglycon delivery approach.^{14,18} Reaction of 1.1 equiv. of **8** with **13** and 1.5 equiv. of 2,3-dichloro-




Scheme 2 Reagents and conditions: i, 1.1 equiv. **8**, 1 equiv. **13**, 1.5 equiv. DDQ, MS 4 Å, CH₂Cl₂, 0 °C to room temp., 2.5 h, 84%; ii, 1.2 equiv. IDCP, MS 4 Å, CH₂Cl₂, room temp., 1.5 h, 73%; iii, Ac₂O, pyridine, room temp., then 1.2 equiv. Buⁿ₄NF, THF, room temp., 0.5 h, 75%, 2 steps; iv, 1.5 equiv. PhSeH, MS 4 Å, cat. HgBr₂, MeCN, room temp., 2 h, 75%; v, 5.5 equiv. EtSH, MS 4 Å, cat. HgBr₂, MeCN, room temp., 48 h, 79%; vi, MeONa, MeOH, room temp., 16 h, 91%; vii, 1 equiv. **18**, 1.1 equiv. **20**, 1.1 equiv. NIS, 0.1 equiv. TMSOTf, MS 4 Å, CH₂Cl₂, -18 °C to room temp., 71%.

5,6-dicyano-1,4-benzoquinone in CH₂Cl₂ gave the mixed acetal **14** in 84% isolated yield after chromatography (Scheme 2). Intramolecular glycosylation was promoted with 1.2 equiv. of iodonium dicollidine perchlorate (IDCP)¹⁹ in CH₂Cl₂ and gave a rewarding 73% yield of the β-linked disaccharide **15** as the only isolated product. The β-(D) configuration of the new anomeric center was firmly established from the ¹H and ¹³C NMR data ($\delta_{H-1'}$ 5.15, $J_{H-1',H-2'}$ 4.5 Hz; $\delta_{C-1'}$ 101.48).²⁰ Acetylation (acetic anhydride, pyridine) and silyl ether deprotection with tetrabutylammonium fluoride in tetrahydrofuran gave the diarabinofuranoside **16** ready for further elongation (75% from **15**).

The dimannosidic glycosyl donor **21** was efficiently obtained from orthoester **17**²¹ taking advantage of the higher reactivity of selenoglycosides over thioglycosides towards activation.^{22,23} **17** was first opened with 1.5 equiv. of selenophenol²⁴ under HgBr₂ catalysis and gave (α-selenoglycoside **18** in 75% yield (Scheme 2). The known thioglycoside **19**,²⁵ obtained in 79% from **17** after HgBr₂-promoted ethanethiol opening, was deacetylated (MeONa, MeOH) to give glycosyl acceptor **20** in 91% yield. Selective activation of selenoglycoside **18** with 1.1 equiv. of *N*-iodosuccinimide (NIS) and 10% trimethylsilyl trifluoromethanesulfonate (TMSOTf),²⁶ and coupling with 1.1 equiv. of thioglycoside **20** in CH₂Cl₂ at -18 °C gave the expected α-linked dimannosidic compound **21** in 71% yield with complete control of the new anomeric center. **21** was isolated in slightly lower yield (64%) when the original silver trifluoromethanesulfonate/potassium carbonate combination²² was used as promoter for the glycosylation of **20** with **18**.

Final coupling was done by glycosylation of disaccharide **16** with 1.5 equiv. of thiodisaccharide **21** under NIS/catalytic TMSOTf activation (Scheme 3). A 70% yield of tetrasaccharidic compounds **22** was obtained as a 4 : 1 α:β anomeric



Scheme 3 Reagents and conditions: i, 1 equiv. **16**, 1.5 equiv. **21**, 1.7 equiv. NIS, 0.15 equiv. TMSOTf, MS 4 Å, CH₂Cl₂, -18 °C, 1 h, 70%; ii, 0.5 M MeONa, MeOH, room temp., 5 h, 75%; iii, H₂, Pd(OH)₂/C, MeOH, room temp., 1 h, 65%.

mixture on the H-1'' anomeric center. This mixture could not be separated at this stage, so the ester groups were removed with 0.5 M sodium methoxide solution in methanol (75%) and the mixture of triols was chromatographed to give pure **23**. Hydrogenolysis of the benzyl groups [H₂, Pd(OH)₂/C, MeOH] gave the target compound **1**²⁷ in 65% yield.

In conclusion, orthoester derivatives of arabinose and mannose were used for the efficient synthesis of **1**, the tetrasaccharidic cap of the lipoarabinomannan of *M. tuberculosis*, via a convergent route using minimal protecting groups manipulation and selective anomeric activation. Both α- and β-arabinofuranosides were obtained with complete stereocontrol from 3-*O*-benzyl-1,2,5-orthoesters of D-arabinose. Extension of this methodology to the elaboration of other complex poly-arabinofuranosidic structures and synthesis of neoantigens from **1** for biological evaluation are currently under way.

Notes and references

- D. Chatterjee and K. H. Khoo, *Glycobiology*, 1998, **8**, 113.
- T. I. A. Roach, C. H. Barton, D. Chatterjee and J. M. Blackwell, *J. Immunol.*, 1993, **150**, 1886.
- D. Chatterjee, K. Lowell, B. Rivoire, M. R. McNeil and P. J. Brennan, *J. Biol. Chem.*, 1992, **267**, 6234.
- S. Prinzi, D. Chatterjee and P. J. Brennan, *J. Gen. Microbiol.*, 1993, **139**, 2649.
- M. Gilleron, N. Himoudi, O. Adam, P. Constant, A. Venisse, M. Rivière and G. Puzo, *J. Biol. Chem.*, 1997, **272**, 117.
- A. Vercellone, J. Nigou and G. Puzo, *Front. Biosci.*, 1998, **3**, e149.
- P. A. Sieling, D. Chatterjee, S. A. Porcelli, T. I. Prigozy, R. J. Mazzaccaro, Y. Soriano, B. R. Bloom, M. B. Brenner, M. Kronenberg, P. J. Brennan and R. L. Modlin, *Science*, 1995, **269**, 227.
- A. Venisse, J. J. Fournié and G. Puzo, *Eur. J. Biochem.*, 1995, **231**, 440.
- R. U. Lemieux, D. R. Bundle and D. A. Baker, *J. Am. Chem. Soc.*, 1975, **97**, 4076.
- N. K. Kochetkov, A. Y. Khorlin, A. F. Bochkov and I. G. Yazlovetskii, *Izv. Akad. Nauk. SSSR, Ser. Khim.*, 1966, 1966; A. F. Bochkov, Y. V. Voznyi, V. N. Chernetskii, V. M. Dashunin and A. V. Rodionov, *Izv. Akad. Nauk. SSSR, Ser. Khim.*, 1975, 348.
- N. K. Kochetkov, A. F. Bochkov and I. G. Yazlovetsky, *Carbohydr. Res.*, 1969, **9**, 49.
- Other catalysts gave mixtures of isomeric 2-*O*- and 5-*O*-benzoylated arabinofuranosides. For a related orthoester opening, see ref. 13.
- F. Nakatsuko, H. Kamitakahara and M. Hori, *J. Am. Chem. Soc.*, 1996, **118**, 1677.
- Y. Ito and T. Ogawa, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1765; A. Dan, Y. Ito and T. Ogawa, *J. Org. Chem.*, 1995, **60**, 4680.
- Y. Gül Salman, Ö. Makinakan and L. Yüceer, *Tetrahedron Lett.*, 1994, **35**, 9233.
- N. Kornblum, R. A. Smiley, R. K. Blackwood and D. C. Iffland, *J. Am. Chem. Soc.*, 1955, **77**, 6229.
- B. S. Sproat, B. Beijer and A. Iribarren, *Nucleic Acids Res.*, 1990, **18**, 41; C. H. Gotfredsen, J. P. Jacobsen and J. Wengel, *Tetrahedron Lett.*, 1994, **35**, 6941.
- For other approaches to the synthesis of β-arabinofuranosides, see: H. B. Mereyala, S. Hotha and M. K. Gurjar, *Chem. Commun.*, 1998, 685; J. Désiré and J. Prandi, *Carbohydr. Res.*, 1999, **317**, 110.
- P. Smid, G. A. de Ruiter, G. A. Van der Marel, F. M. Rombouts and J. H. Van Boom, *J. Carbohydr. Chem.*, 1991, **10**, 833.
- K. Bock and C. Pedersen, *Adv. Carbohydr. Chem. Biochem.*, 1983, **41**, 27; R. C. Beier and B. P. Mundy, *J. Carbohydr. Chem.*, 1984, **3**, 253.
- N. E. Franks and R. Montgomery, *Carbohydr. Res.*, 1968, **6**, 286.
- S. Mehta and B. M. Pinto, *Tetrahedron Lett.*, 1991, **32**, 4435; *J. Org. Chem.*, 1993, **58**, 3269.
- P. Grice, S. V. Ley, J. Pietruszka and H. W. M. Pripke, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 197.
- W. H. H. Gunther, *J. Org. Chem.*, 1966, **31**, 1202.
- T. Peters, *Liebigs Ann. Chem.*, 1991, 135.
- G. H. Veeneman, S. H. Van Leeuwen, H. Zuurmond and J. H. Van Boom, *J. Carbohydr. Chem.*, 1991, **9**, 783.
- Selected data for **1**. [α]_D²⁵ +18 (c 0.44, water); ¹H NMR (500 MHz, D₂O, 293 K): δ 5.16 (d, 1H, *J* 1.7 Hz, H-1''), 5.11 (d, 1H, *J* 4.5 Hz, H-1'), 5.10 (d, 1H, *J* 2 Hz, H-1), 5.02 (d, 1H, *J* 2 Hz, H-1''). ¹³C NMR (125.72 MHz, D₂O, 293 K): δ 106.09 (C-1), 103.01 (C-1''), 101.05 (C-1'), 98.86 (C-1''), 87.48 (C-2), 80.40 (C-2''), 76.67 (C-2').