

## Interaction of Aromatic Hydrocarbon Excited Singlet States with 2-Azidobiphenyl

BY JOHN S. SWENTON,\* THEODORE J. IKELER, and BRIAN H. WILLIAMS

(Department of Chemistry, The Ohio State University, Columbus, Ohio 43210)

**Summary** Markedly different results are reported for ketone- and aromatic hydrocarbon-sensitized decomposition of 2-azidobiphenyl: the ketones transfer triplet excitation to the azide, and the aromatic hydrocarbons singlet energy.

INTERMEDIATES in the direct and photosensitized decomposition of azides have been studied.<sup>1</sup> From the photochemistry of triarylmethyl azides it was concluded that the same intermediate was formed from the excited singlet and triplet states of the azide.<sup>1g</sup> The conclusion was based on identical product distributions from the direct and hydrocarbon sensitized decompositions of a variety of triarylmethyl azides. We report that the sensitized decomposition of 2-azidobiphenyl (**1**) by ketone sensitizers leads to remarkably different results from those obtained by hydrocarbon sensitizers. The results in this system indicate that aromatic hydrocarbons transfer singlet energy not triplet energy to the ground state azide.

Direct irradiation of  $10^{-2}$ M-solutions of 2-azidobiphenyl in ether, benzene, or isopropyl alcohol leads to the formation of carbazole (68—71%) and 2-azobiphenyl (8—11%).<sup>2</sup> The formation of azo-compound is quenched in the presence of piperylene or oxygen. Sensitized decomposition of (**1**)

in benzene at 3500 Å by acetophenone ( $E_T$  74.6 kcal./mole), *m*-methoxyacetophenone ( $E_T$  72.4 kcal./mole), or benzophenone ( $E_T$  68.5 kcal./mole) gave the azo-compound (**3**)


### Irradiation of 2-azidobiphenyl<sup>a</sup>

Irradiation	% Carbazole	% Azo-compound
Direct irradiation .. ..	68—71	8—11
Acetophenone (1.7 M) .. ..	<2	40
<i>m</i> -Methoxyacetophenone (1.4 M) .. ..	<2	39
Benzophenone (0.9 M) .. ..	<2	41
Triphenylene (0.12 M) .. ..	72	14
Naphthalene (0.2 M) <sup>b</sup> .. ..	67	6
Pyrene (0.02 M) .. ..	95	<1

<sup>a</sup> Solutions  $10^{-3}$  M in benzene; <sup>b</sup> 3000 Å source.

with nearly complete exclusion of carbazole. In marked contrast, photosensitized decomposition of (**1**) by triphenylene, naphthalene, or pyrene led to the production of carbazole as the major product. When the triphenylene sensitized decomposition of (**1**) was carried out in the presence of 0.2M-piperylene, no reduction in the rate of azide disappearance was noted. Since piperylene ( $E_T$  ca. 60 kcal./mole) would be an effective quencher of triphenylene triplets ( $E_T$  66.6 kcal./mole), the triplet state of

triphenylene is not responsible for the sensitized decomposition of (1).



The ketone-sensitized decomposition of (1) is without complication because the low concentration of acceptor

employed ( $10^{-2}\text{M}$ ) and the short lifetime of the ketone singlet state ( $<10^{-10}\text{ sec.}$ )<sup>3</sup> permit only triplet-energy transfer from sensitizer to azide. The high yield of carbazole in the hydrocarbon sensitized runs must be due to singlet-energy transfer to the ground-state azide. Caution should be exercised in employing aromatic hydrocarbons as sensitizers.<sup>4</sup> The observations recorded with pyrene are especially interesting since excited pyrene is functioning as a singlet sensitizer while ground-state pyrene acts as a quencher for azo-compound formation.

We thank the Eli Lilly Company, Indianapolis, for a grant.

(Received, August 26th, 1969; Com. 1308.)

<sup>1</sup> (a) A. Reiser, F. W. Willets, G. C. Terry, V. Williams, and R. Marley, *Trans. Faraday Soc.*, 1968, **64**, 3265; (b) A. Reiser and R. Marley, *ibid.*, p. 1806; (c) L. Horner, A. Christmann, and A. Gross, *Chem. Ber.*, 1963, **96**, 399; (d) L. Horner, G. Bauer, and J. Dörge, *ibid.*, 1965, **98**, 2631; (e) F. D. Lewis and W. H. Saunders, *J. Amer. Chem. Soc.*, 1968, **90**, 7031; (f) *ibid.*, p. 7033; (g) *ibid.*, 1967, **89**, 645; (h) W. H. Saunders and E. A. Caress, *ibid.*, 1964, **86**, 861; (i) A. Misha, S. N. Rice, and W. Lwowski, *J. Org. Chem.*, 1968, **33**, 481; (j) J. S. McConaghy and W. Lwowski, *J. Amer. Chem. Soc.*, 1967, **89**, 4450; (k) J. W. ApSimon and O. E. Edwards, *Canad. J. Chem.*, 1962, **40**, 896; (l) R. Moriarty and M. Rahman, *Tetrahedron*, 1965, **21**, 2877; (m) G. Smolinsky, E. Wasserman, and W. A. Yager, *J. Amer. Chem. Soc.*, 1962, **84**, 3220; (n) *ibid.*, 1964, **86**, 3166.

<sup>2</sup> J. S. Swenton, *Tetrahedron Letters*, 1968, 3421.

<sup>3</sup> F. Wilkinson and J. T. Dubois, *J. Chem. Phys.*, 1963, **39**, 377.

<sup>4</sup> Recently there have been several reports of the involvement of hydrocarbon singlet states where the triplet state had originally been supposed: S. L. Murov, R. S. Cole, and G. S. Hammond, *J. Amer. Chem. Soc.*, 1968, **90**, 2958; R. S. Cole and G. S. Hammond, *ibid.*, p. 2958; P. D. Bartlett and P. S. Engel, *ibid.*, 1968, **90**, 2961.