


Figure 1. Predicted equilibrium geometries for the vinylidene and σ -bonded isomers of $\text{Al-C}_2\text{H}_2$. There are two conformers of the σ -bonded system, with the terminal hydrogen lying trans (the middle structure) or cis (the lowest structure) to the Al atom. Bond distances are in Å.

separated aluminum plus acetylene.¹³ Configuration interaction including all valence electron single and double excitations (3461 configurations) reduces the dissociation energy of the vinylidene complex slightly, to 19.3 kcal. A correction for higher excitations (unlinked clusters)¹⁴ provides our final ab initio prediction of 20.0 kcal for the Al-CCH_2 bond energy. The shorter (by 0.08 Å) Al-C internuclear separation for the vinylidene complex also attests to its stronger Al-C bond. Analogous CI studies of the trans- σ -bonded complex (6739 configurations) yield a binding energy of 7.5 kcal.

The vinylidene structure is inconsistent with Kasai, McLeod, and Watanabe's ESR spectrum; there may appear to be a conflict between theory and experiment. However, the barrier height between vinylidene and acetylene is ~ 8 kcal⁸ and might not be greatly reduced by the presence of the Al atom. If this is the case, then the 1,2-hydrogen shift (eq 4) might not occur



at all at the liquid helium temperatures of Kasai and McLeod.²

The most fascinating feature of the present study is that, while the isolated acetylene-vinylidene reaction is *endothermic* by 40 kcal, the same process becomes *exothermic* in the presence of an Al atom. We suspect that other endothermic reactions such as the methylcarbene-ethylene and methylnitrene-methylenimine rearrangements will also become favorable when complexation to a metal atom, metal cluster, or metal surface⁶ is achieved. This general mechanism for using metal systems to transform endothermic reactions into nearly degenerate rearrangements may play a key role in catalysis and surface chemistry.

Acknowledgment. This research was supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, U.S. Department of Energy, through the Materials and Molecular Research Division of the Lawrence Berkeley Laboratory.

References and Notes

- (1) P. H. Kasai and D. McLeod, *J. Am. Chem. Soc.*, **97**, 5609 (1975).
- (2) P. H. Kasai, D. McLeod, and T. Watanabe, *J. Am. Chem. Soc.*, **99**, 3521 (1977).
- (3) M. J. S. Dewar, *Bull. Soc. Chim. Fr.*, **18C**, 79 (1951).
- (4) W. C. Swope and H. F. Schaefer III, *J. Am. Chem. Soc.*, **98**, 7962 (1976).
- (5) B. R. Brooks and H. F. Schaefer III, *Mol. Phys.*, **34**, 193 (1977).
- (6) H. F. Schaefer III, *Acc. Chem. Res.*, **10**, 287 (1977).
- (7) A. K. Rappe and W. A. Goddard, *J. Am. Chem. Soc.*, **99**, 3966 (1977).
- (8) C. E. Dykstra and H. F. Schaefer III, *J. Am. Chem. Soc.*, **100**, 1378 (1978).
- (9) T. H. Dunning and P. J. Hay, "Modern Theoretical Chemistry", Vol. 3, H. F. Schaefer III, Ed., (Plenum Press, New York, 1977, pp 1-27).
- (10) H. F. Schaefer III, "The Electronic Structure of Atoms and Molecules: A Survey of Rigorous Quantum Mechanical Results", Addison-Wesley, Reading, Mass., 1972.
- (11) F. W. Bobrowicz and W. A. Goddard, ref 9, pp 79-127.
- (12) H. F. Schaefer III, "XeF: Chemically Bound?", Lawrence Berkeley Laboratory Report No. 6298, May 1977.
- (13) The total energies for infinitely separated $\text{Al} + \text{C}_2\text{H}_2$ are -318.6542 (SCF) and -318.8515 hartrees (CI).
- (14) S. R. Langhoff and E. R. Davidson, *Int. J. Quantum Chem.*, **52**, 403 (1978).

Michael Trenary, Mark E. Casida
Bernard R. Brooks, Henry F. Schaefer III*

Department of Chemistry and Lawrence Berkeley Laboratory
University of California, Berkeley, California 94720

Received September 11, 1978

Laser Flash Photolysis with NMR Detection. Microsecond Time-Resolved CIDNP: Separation of Geminate and Random-Phase Processes

Sir:

The time resolution of conventional high resolution NMR spectroscopy is low because of the intrinsic properties of nuclear spin systems. Line-shape analysis for systems undergoing spin exchange overcomes this limitation to some extent.¹ Flow and stopped-flow methods permit direct kinetic measurements, but are at best restricted to reactions occurring over tens of milliseconds.² CIDNP provides information on chemical events taking place on a time scale ranging from 10^{-8} to 10^{-3} s by freezing this information into patterns of nuclear spin polarization within the diamagnetic reaction products.³ This information is conveniently sampled on the time scale of the nuclear spin-lattice relaxation times of the products, that is, seconds. Kinetic analysis of data obtained under such steady-state conditions is, however, highly indirect and is dependent on many theoretical assumptions. We report here examples of time-resolved CIDNP with a resolution of 1×10^{-6} s, a time scale which allows the direct measurement of useful chemical kinetics. Ernst and co-workers have reported qualitatively similar experiments with a time resolution of 2.5×10^{-3} s.⁴ These latter experiments were directed at specific dynamic nuclear spin phenomena^{4a} and illustrate the observation of slow secondary reactions of diamagnetic transients.^{4b}

The experiment, schematically represented in Figure 1, is in principle applicable to any system in which photolysis creates a radical pair. The sample is placed in the probe of a high resolution NMR spectrometer and is subjected to a series of radio frequency pulses randomly modulated in time and phase. This saturation pulse of ~ 50 -ms duration totally destroys any nuclear magnetization thus eliminating all background signals.⁵ The sample is then irradiated by an intense laser flash lasting several nanoseconds. The excited molecules generate radical pairs which induce nuclear magnetization by geminate and random-phase processes. The laser pulse is followed at any desired time interval, τ , by a radio frequency pulse creating a

- (2) C. A. Fyfe, M. Cocivera, and S. W. H. Damji, *Acc. Chem. Res.*, **11**, 277 (1978).
 (3) L. T. Muus, P. W. Atkins, K. A. McLauchlan, and J. B. Pedersen, Eds., "Chemically Induced Magnetic Polarization", Reidel, Dordrecht, 1977.
 (4) (a) S. Schäublin, A. Wokaun, and R. R. Ernst, *Chem. Phys.*, **14**, 285 (1976); (b) *J. Magn. Reson.*, **27**, 273 (1977).
 (5) Under the present operating conditions there is a 25-ms delay between the end of the saturation pulse and the laser flash. This delay arises because the laser charge cycle runs on the 60 Hz mains. Recovery of substrate and/or product signals during this interval is insufficient to exceed the noise level. This delay is in fact exploited by doping the solvent with protio benzene (10% v/v)—enough signal recovery occurs to provide an internal reference (see

- Figure 2) allowing normalization of the CIDNP intensities.
 (6) H.-G. Heine, W. Hartmann, D. R. Kory, J. G. Magyar, C. E. Hoyle, J. K. McVey, and F. D. Lewis, *J. Org. Chem.*, **39**, 691 (1974).
 (7) Appropriate time-resolved CIDEP studies are also applicable to the separation of these processes.³

G. L. Closs,* R. J. Miller

Department of Chemistry, The University of Chicago
 Chicago, Illinois 60637

Received November 27, 1978

Book Reviews*

Aspects of Degradation and Stabilization of Polymers. Edited by H. H. G. JELLINEK (Clarkson College of Technology). Elsevier, Amsterdam, 1978. 690 pp. \$124.50.

Fourteen contributed chapters deal with the properties of polymers relevant to their stability and with their modes of breakdown. Emphasis is on technical applications. Specific topics include mechanical degradation, effect of degradation on mechanical properties, reaction of polymers with pollutant gases, ignition of polymers and flame propagation, degradation processes in ablation, biodegradation, and degradation kinetics.

The Structure of Non-Crystalline Materials. Edited by P. H. GASKELL. Crane, Russak & Co., New York. 1977. 262 pp. \$37.50.

A handsomely printed quarto-size volume containing the Proceedings of the Symposium held in Cambridge, England, 20–23 September, 1976, organized by the Society of Glass Technology. Fifty contributed articles provide a timely panoramic view of the structure and properties of amorphous systems, both experimental and theoretical aspects. A contribution by Nobel laureate N. F. Mott reviews electronic processes in glasses.

High-Power Lasers and Applications (Springer Series in Optical Sciences). Volume 9. Edited by K.-L. KOMPA and H. WALTHER. Springer-Verlag, Berlin. 1978. ix + 228 pp. \$24.00.

Proceedings of the Fourth Colloquium on Electronic Transition Lasers at Munich, June 20–22, 1977. The main topics are high-power VUV, UV, and visible and IR lasers, with applications to nonlinear optics, chemical kinetics, isotope separation, and spectroscopy.

The Reward System in British and American Science. By JERRY GASTON (Southern Illinois University). Wiley/Interscience, New York. 1978. xiii + 204 pp. \$18.95.

Do scientists receive recognition commensurate with their scientific contributions, free from the influence of extraneous social and political factors? The author's study of 600 British and American University physicists, chemists, and biologists indicates that they are indeed fairly rewarded for their productivity, instances to the contrary being statistically insignificant. The most accessible indicators of productivity—which should surprise none of the readers of this Journal—are number of publications and number of citations. A couple of interesting tidbits: Scientists have mean IQ's well above average but, beyond this threshold, there is practically no correlation between IQ and achievement. Also, the later productivity of scientists appears to be substantially influenced by the recognition accorded their early work as new Ph.D.'s.

Coordination Chemistry. Volume 2 (ACS Monograph, No. 174). Edited by A. E. MARTELL (Texas A & M University). American Chemical Society, Washington, D.C. 1978. x + 636 pp. \$90.00.

Volume 2 treats the kinetics and mechanisms of reactions involving coordination compounds whereas Volume 1 (ACS Monograph No. 168, published in 1971) was more concerned with physical and chemical properties. The Monograph contains three chapters: 1. Kinetics and Mechanisms of Complex Formation and Ligand Exchange

(Margerum, Cayley, Weatherburn, and Pagenkopf); 2. Ligand Reactions—The Effect of Metal Complexes on Chemical Processes (Hipp and Busch); 3. Oxidation–Reduction Reactions of Coordination Complexes (Pennington). Chapter 1 focuses on the reactions of the more labile metal ions and their complexes in aqueous solution. Chapter 2 views the known metal-affected reactions from a structural and mechanistic standpoint. Chapter 3, a full-scale review of oxidation–reduction chemistry, includes recent innovations in excited state and intramolecular electron transfer.

Atlas of Metal-Ligand Equilibria in Aqueous Solution. By J. KRAGTEN (University of Amsterdam). Wiley/Halsted, New York. 1978. 781 pp. \$77.50.

A volume in the Ellis Horwood Series in Analytical Chemistry. This atlas gives computer-generated plots of side-reaction coefficients vs. pH for some 45 metals in the presence of 29 common ligands. The graphs for separate metal–ligand combinations are superposable to provide information on mixed ligands. Accompanying plots of pM' vs. pH show the regions of predominance of various species and indicate the conditions under which hydroxides will precipitate and polynuclear hydroxo complexes will form.

Ion-Selective Electrodes. Edited by E. PUNGOR and I. BUZAS. Elsevier, Amsterdam. 1978. iv + 613 pp. \$75.00.

This volume contains the full text of papers presented at a conference held in Budapest, 5–9 September 1977. Seven plenary lectures covered recent trends in ion-selective electrode research, theoretical problems concerning the mechanism of electrode response, electrode construction and standardization techniques, and high-temperature applications. There follow some 48 discussion papers concentrating mainly on analytical applications of ion-selective electrodes. Topics discussed include transport characteristics and dynamical behavior of glass membrane electrodes, enzyme electrodes, liquid membrane electrodes, ion-exchanger electrodes, and incorporation of ion-selective electrodes into automated systems.

The Quest for Absolute Zero. 2nd Edition. By KURT MENDELSSOHN (Oxford). Wiley/Halsted Press, New York. 1977. 281 pp. \$11.50 (paperback).

A charming account of the history of low-temperature research by an important contributor to the field. The book contains lucid semi-technical accounts of cryogenic techniques, thermodynamic and quantum principles near absolute zero, and the phenomena of superconductivity and superfluidity.

Chemistry and Physics of Carbon: A Series of Advances. Volume 13. Edited by P. L. WALKER, JR., and P. A. THROWER. Marcel Dekker, New York. 1977. viii + 295 pp. \$32.50.

The volume consists of two chapters: The Optical Properties of Diamond (G. Davies) and Fracture in Polycrystalline Graphite (J. E. Brocklehurst). The effects of nitrogen impurities in diamond are considered in some detail. Author and subject indexes are appended.

Defects in the Alkaline Earth Oxides. By B. HENDERSON (Trinity College, Dublin) and J. E. WERTZ (University of Minnesota). Wiley/Halsted Press, New York. 1977. v + 159 pp. \$20.00.

* Unsigned book reviews are by the Book Review Editor.