



A Journal of



Accepted Article

Title: ^{11}C labeling: Intracyclic incorporation of carbon-11 into heterocycles

Authors: François Liger, Florence Cadarossane, Thibaut Iecker, Christian Tourvieille, Didier Le BARS, and Thierry Billard

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.201901386

Link to VoR: <http://dx.doi.org/10.1002/ejoc.201901386>

Supported by



WILEY-VCH

COMMUNICATION

¹¹C labeling: Intracyclic incorporation of carbon-11 into heterocycles.

François Liger,^[a] Florence Cadarossanesaib,^[a] Thibaut Iecker,^[a] Christian Tourvieille,^[a] Didier Le Bars,^{[a],[b]} and Thierry Billard^{*[a],[b]}

Abstract: Labeling of heterocycles with carbon-11 is generally performed through peripheral functionalizations and more scarcely inside heterocyclic core. Such less common approach usually requires preliminary multi-step synthesis of reactive species. Herein, a cyclization reaction by direct use of cyclotron-produced [¹¹C]CO₂ is described to obtain various heterocycles intracyclically labeled in only 10 minutes.

Introduction

Positron emission tomography (PET) is a non-invasive medical imaging technique used as diagnostic tool in several pathologies.^[1] PET is also more and more applied in drug design, allowing in vivo quantitative biodistribution and pharmacokinetic studies.^[2] This method requires radiolabeling of molecules with a positron emitting radioisotope.^[3] Among the most useful isotopes, fluorine-18 and carbon-11 are preferentially employed. Even if fluorine-18 is preferred due to its superior radioactive half-life (109 min for fluorine-18 vs 20 min for carbon-11), ¹¹C-molecules are still predominantly used for research purposes. Indeed, it is easier to obtain radiolabeled equivalent of existing drugs by "isotopic replacement" of carbon atom.

The main strategy for compounds labeling is generally peripheral introduction of ¹¹C-substituents.^[3a, 4] Radiolabeling inside the molecules core, in particular of heterocyclic structures, has been scarcely described.

Few intracyclic labelings with carbon-11 have been previously performed with various ¹¹C-reagents:^[5] [¹¹C]phosgene,^[6] [¹¹C]formic acid or derivatives,^[7] [¹¹C]formaldehyde,^[8] [¹¹C]cyanide,^[9] [¹¹C]thiocyanate,^[10] [¹¹C]CS₂^[11] or [¹¹C]CO.^[12] However, all these reagents require their preliminary preparation, often in tedious conditions, which is time consuming. The starting material is generally [¹¹C]CO₂ produced with a cyclotron by ¹⁴N(p, α)¹¹C nuclear reaction. Consequently, direct use of cyclotron-produced [¹¹C]CO₂ could be of interest to obtain intracyclic

radiolabeled heterocycles. Nevertheless, because of low reactivity of CO₂, this approach has so far been little investigated. A first approach has been described by an in situ formation of isocyanates.^[13] [¹¹C]CO₂ has been trapped by a diamine or an amino-alcohol in a first step, followed by a dehydration in a second step, mediated by POCl₃. The resulting isocyanates were then cyclized to provide corresponding imidazolones or oxazolidinones. This method still requires two steps and appears limited in term of substrates scope. More recently, cyclic ureas were synthesized by cyclization of azidoamines with [¹¹C]CO₂, mediated by phosphine.^[14] However, this method is limited to urea compounds. Furthermore, heteroaromatic compounds cannot be directly obtained following these strategies.

Results and Discussion

We have previously described direct ¹¹C-methylation of amines through an in situ metal/carbene-mediated reduction of carbon dioxide.^[15] During this process, the intermediate formation of formamides has been demonstrated. Consequently, we supposed that a similar strategy applied to diamines could lead to cyclic compounds.

This hypothesis has been evaluated with 1,2-phenylenediamine (**1a**) as starting material to obtain the corresponding benzimidazole (**2a**) (Table 1).


Starting from the described conditions for methylation of amines (entry 1), we were delighted to observe the formation of the expected benzimidazole **2a**, with good radiochemical yields (entry 1). However, in this assay, 10 mg of diamine **1a** have been used. Generally, smaller amounts (ca. 20 μmol) of starting material are engaged in radiolabeling reactions. Unfortunately, by decreasing the diamine **1a** quantity (1.1 mg), to match with a more traditional amount, radiochemical yield also dramatically diminished (entry 2). By using a slight excess of ZnCl₂ and **IPr** (1,3-Bis(2,6-diisopropylphenyl)imidazol-2-ylidene), a satisfactory result was reobtained (entry 3). However, an excess of silane appeared to be necessary to observe the expected cyclization (entries 3-4). Some similar cyclizations have also been described with non-radioactive CO₂.^[16] Described conditions did not use any metal catalysts with a reaction time of 24h. Nevertheless, in our case the use of ZnCl₂ appeared to be crucial to observe cyclization in a reaction time of only 20 min, more consistent with carbon-11 half-life (entry 5). The metal certainly contributes to accelerate reaction so fitting with the short time imposed by radioactive decay.

[a] F. Liger, F. Cadarossanesaib, T. Iecker, C. Tourvieille, Dr. D. Le Bars, Prof. Dr. T. Billard
CERMEP-In vivo imaging
59 Bd Pinel, 69677 Lyon, France
E-mail: Thierry.billard@univ-lyon1.fr
www.fmi-lyon.fr

[b] Dr. D. Le Bars, Prof. Dr. T. Billard
Institute of Chemistry and Biochemistry (UMR CNRS 5246),
Univ Lyon, Université Lyon 1, CNRS, CPE, INSA
43 Bd du 11 novembre 1918, 69622 Villeurbanne, France

Supporting information for this article is given via a link at the end of the document. ((Please delete this text if not appropriate))

COMMUNICATION

Table 1. Cyclization reaction with **1a** and [^{11}C]CO $_2$.

Entry	1a (mg)	ZnCl $_2$ (equiv.)	IPr (equiv.)	Silane (equiv.)	2a (%) ^[a]
1	10	0.15	0.15	PhSiH $_3$ (1.00)	68
2	ca. 1.1	0.15	0.15	PhSiH $_3$ (8.50)	27
3	ca. 1.1	1.2	1.2	PhSiH $_3$ (8.50)	52
4	ca. 1.1	1.2	1.2	PhSiH $_3$ (1.60)	-
5	ca. 1.1	-	1.2	PhSiH $_3$ (8.50)	-
6	ca. 1.1	1.2	1.2	PHMS (28.90)	32
7	ca. 1.1	1.2	1.2	Ph $_2$ SiH $_2$ (11.00)	83
8	ca. 1.1	1.2 ^[b]	1.2	Ph $_2$ SiH $_2$ (11.00)	-

[a] Radiochemical yields (RCY) were estimated from trapped [^{11}C]CO $_2$ within the reactor and are decay-corrected from end of [^{11}C]CO $_2$ trapping inside reactor. [b] with FeCl $_3$ instead of ZnCl $_2$.

With PhSiH $_3$ as reducing agent, the cyclization reaction is in competition with methylation of amino groups which can proceed in similar conditions.^[15] Indeed, some side-products arising from this methylation competitive reaction were also observed. To circumvent this drawback, less reducing silanes have been envisaged. If only low yields were obtained with PMHS (Poly(methylhydrosiloxane)), a higher and very good radiochemical yield was observed with Ph $_2$ SiH $_2$ (entries 6-7). In this case, no methylation of amines was detected. Finally, FeCl $_3$ has also been envisaged as catalyst but without any success (entry 8).

Even if a reaction time of 20 min is compatible with carbon-11 radiolabeling, shorter durations are generally expected. Consequently, with optimal conditions (entry 7), reaction time was reduced to 10 min. An important decrease of radiochemical yield has been observed (Table 2, entries 1-2). Microwave activation was deleterious for the reaction, as already observed in the previously described radiomethylation reactions (entry 3).^[15] Doubling catalysts amounts did not bring significant improvement (entry 4). In contrast, an increase of concentration of precursor **1a**,

associated with a decrease of reaction volume, brought positive effects and excellent radiochemical yields were observed (entries 5-7). By switching from 200 μL to 100 μL , no significant improvement was observed. An attempt to reduce reaction time to 5 min led only to a medium result (entry 8). In view of these results, a volume of 200 μL will be generally preferred for further investigations, to avoid potential solubility problems.

Table 2. Effect of reaction time and solvent volume^[a]

Entry	Reaction time (min)	Diglyme volume (μL)	2a (%) ^[b]
1	20	400	83
2	10	400	25
3	10 ^[c]	400	1
4	10 ^[d]	400	38
5	10	300	66
6	10	200	87
7	10	100	90
8	5	100	48


[a] Conditions: **1a** (1.1-2.2 mg), ZnCl $_2$ (1.2 equiv.), IPr (1.2 equiv.), Ph $_2$ SiH $_2$ (11 equiv.), diglyme, 150°C, 10 min. [b] Radiochemical yields (RCY) were estimated from trapped [^{11}C]CO $_2$ within the reactor and are decay-corrected from end of [^{11}C]CO $_2$ trapping inside reactor. [c] under microwave irradiation (100 W). [d] ZnCl $_2$ (2.4 equiv.), IPr (2.4 equiv.).

In these optimal conditions, molar activity of produced benzimidazole **2a** was around 18 GBq/ μmol . To study the scope of this radiolabeling, various aromatic diamines were then studied (Scheme 1).

Various benzimidazoles were obtained with, in general, satisfactory to good radiochemical yields. Substitution onto nitrogen atom does not significantly influence the obtained results (**2a-c**). Whatever the electronic character (electron withdrawing or donor) of aromatic substituents, similar yields were observed (**2a-f**). However, slightly lower yields were observed with **2g** and **2h**, maybe due to a competitive coordination of ZnCl $_2$ catalyst by the carbonyl function (**2g**) or the nitrogen atom of the pyridine core (**2h**). Interestingly, reaction is compatible with functional groups such as ketone (**2g**), ester (**2e**) or bromine (**2d**). In particular, no reduction side-products were detected.


To extend this radiolabeling methodology, 6-membered rings have been also studied. Thus, perimidine **2i** was obtained with similar good yields compared to benzimidazoles. More interestingly, quinazolinone **2j** was also synthesized starting from corresponding 2-aminobenzamide. Because of the weakest nucleophilicity of amide group, the expected cyclic product was formed with only a medium radiochemical yield. However, a good result was achieved by doubling reaction time to 20 min. By decreasing solvent volume to 100 μL , a better yield was also observed in 10 min.

COMMUNICATION




If benzimidazoles represent privileged structure in medicinal chemistry and drugs,^[17] benzothiazoles are also important heterocyclic compounds in drug design and pharmaceutical agents.^[18] This heterocycle is also used in radiotracers development, as illustrated by β -amyloid radioligand $[^{11}\text{C}]\text{PIB}$.^[19] Consequently, our method was extended to aminobenzenethiols (**3**) to obtain, for the first time, intracyclically ^{11}C -labeled benzothiazoles **4** (Scheme 2).

With optimal conditions used for benzimidazoles, only modest radiochemical yield was observed (**4a**). As demonstrated previously (Scheme 1; **2j**), by decreasing solvent volume (100 μL instead of 200 μL), a better result is obtained. Nevertheless, in general, radiochemical yields remain modest, but suitable for workable radiolabelings.



Scheme 2. Cyclizations of aminobenzenethiols with $[^{11}\text{C}]\text{CO}_2$. Radiochemical yields are mean values and were estimated from trapped $[^{11}\text{C}]\text{CO}_2$ within the reactor and are decay-corrected from end of $[^{11}\text{C}]\text{CO}_2$ trapping inside reactor. [a] diglyme volume = 100 μL .


This one-step, late stage, intracyclic labeling has been applied to the first ^{11}C -labeling of theophylline drug^[20] (**2k**), with intracyclic incorporation of radioisotope (Scheme 3).



Scheme 3. Intracyclic labelling of theophylline. Radiochemical yields are mean values and were estimated from trapped $[^{11}\text{C}]\text{CO}_2$ within the reactor and are decay-corrected from end of $[^{11}\text{C}]\text{CO}_2$ trapping inside reactor. [a] diglyme volume = 100 μL .

$[^{11}\text{C}]\text{Theophylline}$ was obtained in a modest radiochemical yield. However, in view of the low reactivity of the starting diamine, the result remains quite satisfactory and demonstrates the usefulness of this method to label elaborated molecules.


During these previous investigations, some reproducibility issues were observed with a batch of **IPr**. Despite various analyses of the different batches of **IPr**, we have not succeeded to determine the reasons of such issues. Consequently, to try to circumvent such potential drawback, a preliminary preparation of **IPr** solutions in diglyme, just before use, was envisaged (Scheme 4). In comparison to commercial **IPr**, similar results were observed with benzimidazole or benzothiazole. This use of home-made solution of **IPr** constitutes a valuable alternative to potential reproducibility issue and we warmly recommend this simple to implement solution to avoid some disappointments during radiolabelings.



Scheme 4. Cyclization reactions with home-made solution of **IPr**. Radiochemical yields are mean values and were estimated from trapped $[^{11}\text{C}]\text{CO}_2$ within the reactor and are decay-corrected from end of $[^{11}\text{C}]\text{CO}_2$ trapping inside reactor.

COMMUNICATION

Finally, in order to demonstrate the effective productivity of this new radiolabeling approach, a full radiosynthesis, with HPLC purification, was performed with perimidine **2i** (Scheme 5). Noteworthy, a home-made solution of **IPr** was preferred.



Scheme 5. Production of perimidine **2i**. The RCY value is mean value and was estimated from trapped $[^{11}\text{C}]\text{CO}_2$ within the reactor and are decay-corrected from end of $[^{11}\text{C}]\text{CO}_2$ trapping inside reactor.

A good radioactive quantity (140 mCi, 5.2 GBq) of $[^{11}\text{C}]$ perimidine (**2i**) was produced with a high radiochemical yield (RCY = 75%) and with a good radiochemical and chemical purity. This result is very encouraging and demonstrates the usefulness of this original radiolabeling method to produce radiotracers.

Conclusions

To conclude, cyclization reactions by direct use of cyclotron-produced $[^{11}\text{C}]\text{CO}_2$ were described to obtain in a late stage and one-pot process, heterocyclic and heteroaromatic compounds intracyclically labeled with carbon-11. This strategy constitutes a proof of concept about direct use of $[^{11}\text{C}]\text{CO}_2$ in “radiocyclization” reactions and should open the way to new developments concerning radiolabeling of heterocyclic molecules.

Experimental Section

Synthesis of $[^{11}\text{C}]$ perimidine (2i**):** 200 µL of freshly prepared **IPr** solution in diglyme (24 µmol, 1.2 eq) was added to a 1 mL vial containing 3.15 mg of 1,8-diaminonaphthalene (20 µmol, 1eq). The resulting solution was sonicated for 2 min and transferred to a 1.1 mL conical vial pre-charged with 3.35 mg of zinc chloride (24.2 µmol, 1.21 eq). After addition of 40 µL of diphenylsilane (0.215 mmol, 11 eq), the vial was sealed and placed into the automated synthesis module 10 min before end of bombardment. When the desired level of radioactivity was reached in cyclotron (nearly 3000 mCi – 111 GBq), reaction with $[^{11}\text{C}]\text{CO}_2$ proceeded according previous general procedure (see SI). After 10 min of heating at 150°C, the reaction vial was cooled at 20°C for 2.5 minutes. Then, 2 x 100 µL of MeOH were sequentially added to the reaction mixture (30 seconds apart) and after further 20 seconds, the whole content of the reaction vial was loaded into a 3 mL HPLC loop. 700 µL of additional MeOH were used for rinsing the reaction vial and loaded into the HPLC loop before injection on semi-preparative HPLC column (Macherey-Nagel Nucleodur C18 HTec column (10µm, 10 x 250mm). HPLC purification was conducted at 3 mL/min with acetate buffer pH 5.2/MeCN (55/45) as mobile phase. Radioactive and UV monitoring at 254 nm allowed the collection of 346–386 mCi (12.8–14.3 GBq, decay-corrected) of purified 1H- $[^{11}\text{C}]$ Perimidine **2i**. Retention time

was 8 minutes. Radiochemical yields were ranging from 70 to 80% decay-corrected.

Acknowledgements ((optional))

This work was supported by European Union (EU) through the project Radiochemistry for Molecular Imaging (project number EU FP7-PEOPLE-2012-ITN-RADIOMI). Frédéric Bonnefoi (CERMEP - in vivo imaging) is acknowledged for his technical assistance. The authors wish to thank the Centre National de la Recherche Scientifique (CNRS), Institut national de la santé et de la recherche médicale (INSERM) and University of Lyon 1 for financial support.

Keywords: Carbon-11 • Heterocycles • Radiochemistry • Radiolabeling • Positron emission tomography

- [1] a) W. Cai, *Mol. Pharm.* **2014**, *11*, 3773-3776; b) G. B. Saha, *Basics of PET Imaging: Physics, Chemistry, and Regulations*, Springer New York, **2010**; c) L. Zimmer, A. Luxen, *Neuroimage* **2012**, *61*, 363-370; d) S. M. Ametamey, M. Honer, P. A. Schubiger, *Chem. Rev.* **2008**, *108*, 1501-1516.
- [2] a) R. Chakravarty, H. Hong, W. Cai, *Mol. Pharm.* **2014**, *11*, 3777-3797; b) C. C. Wagner, O. Langer, *Adv. Drug Del. Rev.* **2011**, *63*, 539-546; c) J. K. Willmann, N. Van Bruggen, L. M. Dinkelborg, S. S. Gambhir, *Nat. Rev. Drug Discov.* **2008**, *7*, 591-607; d) M. Schou, K. Varnäs, *Clin. Transl. Imaging* **2017**, 1-8; e) A. Testa, M. Zanda, C. S. Elmore, P. Sharma, *Mol. Pharm.* **2015**, *12*, 2203-2216.
- [3] a) X. Deng, J. Rong, L. Wang, N. Vasdev, L. Zhang, L. Josephson, S. H. Liang, *Angew. Chem. Int. Ed.* **2019**, *58*, 2580-2605; b) P. W. Miller, N. J. Long, R. Vilar, A. D. Gee, *Angew. Chem. Int. Ed.* **2008**, *47*, 8998-9033; c) Z. Li, P. S. Conti, *Adv. Drug Del. Rev.* **2010**, *62*, 1031-1051.
- [4] a) K. Dahl, C. Hallidin, M. Schou, *Clin. Transl. Imaging* **2017**, *5*, 275-289; b) C. Taddei, A. D. Gee, *J. Labelled Compd. Radiopharm.* **2018**, *61*, 237-251.
- [5] a) B. H. Rotstein, S. H. Liang, M. S. Placzek, J. M. Hooker, A. D. Gee, F. Dolle, A. A. Wilson, N. Vasdev, *Chem. Soc. Rev.* **2016**, *45*, 4708-4726; b) A. Del Vecchio, G. Destro, F. Taran, D. Audisio, *J. Labelled Compd. Radiopharm.* **2018**, *61*, 988-1007.
- [6] a) G. D. Brown, S. K. Luthra, C. S. Brock, M. F. G. Stevens, P. M. Price, F. Brady, *J. Med. Chem.* **2002**, *45*, 5448-5457; b) F. Brady, S. K. Luthra, H.-J. Tochon-Danguy, C. J. Steel, S. L. Waters, M. J. Kensett, P. Landais, F. Shah, K. A. Jaeggi, A. Drake, J. C. Clark, V. W. Pike, *Int. J. Rad. Appl. Instrum. A* **1991**, *42*, 621-628; c) C. Boullais, C. Crouzel, A. Syrota, *J. Labelled Compd. Radiopharm.* **1986**, *23*, 565-567; d) G. A. Brinkman, I. Hass-Lisewska, J. T. Veenboer, L. Lindner, *Int. J. Appl. Radiat. Isot.* **1978**, *29*, 701-702; e) C. Crouzel, G. Mestelan, E. Kraus, J. M. Lecomte, D. Comar, *Int. J. Appl. Radiat. Isot.* **1980**, *31*, 545-548; f) D. Roeda, G. Westera, *Int. J. Appl. Radiat. Isot.* **1981**, *32*, 843-845; g) M. Berridge, D. Comar, C. Crouzel, *Int. J. Appl. Radiat. Isot.* **1983**, *34*, 1657-1658; h) K. Yashio, Y. Katayama, T. Takashima, N. Ishiguro, H. Doi, M. Suzuki, Y. Wada, I. Tamai, Y. Watanabe, *Bioorg. Med. Chem. Lett.* **2012**, *22*, 115-119; i) M. Berridge, D. Comar, C. Crouzel, J. C. Baron, *J. Labelled Compd. Radiopharm.* **1983**, *20*, 73-78; j) D. Roeda, B. Tavitian, C. Coulon, F. David, F. Dolle, C. Fuseau, A. Jobert, C. Crouzel, *Bioorg. Med. Chem.* **1997**, *5*, 397-403; k) K.-I. Seki, K.-I. Nishijima, K. Sanoki, Y. Kuge, M. Takahashi, H. Akizawa, N. Tamaki, L. I. Wiebe, K. Ohkura, *Heterocycles* **2009**, *77*, 1307-1321; l) Y. Takada, M. Ogawa, H. Suzuki, T. Fukumura, *Appl. Radiat. Isot.* **2010**, *68*, 1715-1720; m) G. Roger, F. Dolle, B. De Bruin, X. Liu, L. Besret, Y. Bramouille, C. Coulon, M. Ottaviani, M. Bottlaender, H. Valette, M. Kassiou, *Bioorg. Med. Chem.* **2004**, *12*, 3229-3237; n) F. Dolle, Y. Bramouille, F. Hinnen, S. Demphel,

COMMUNICATION

- P. George, M. Bottlaender, *J. Labelled Compd. Radiopharm.* **2003**, *46*, 783-792; o) C. J. Steel, F. Brady, S. K. Luthra, G. Brown, I. Khan, K. G. Poole, A. Sergis, T. Jones, P. M. Price, *Appl. Radiat. Isot.* **1999**, *51*, 377-388; p) M. Berridge, D. Comar, D. Roeda, A. Syrota, *Int. J. Appl. Radiat. Isot.* **1982**, *33*, 647-651; q) D. Roeda, C. Crouzel, B. Van Zanten, *Radiochem. Radioanal. Lett.* **1978**, *33*, 175-178; r) P. Landais, C. Crouzel, *Appl. Radiat. Isot.* **1987**, *38*, 297-300; s) M. Ogawa, Y. Takada, H. Suzuki, K. Nemoto, T. Fukumura, *Nucl. Med. Biol.* **2010**, *37*, 73-76; t) Y. Bramoulle, D. Roeda, F. Dolle, *Tetrahedron Lett.* **2010**, *51*, 313-316; u) S. Bernard, C. Fuseau, L. Schmid, R. Milcent, C. Crouzel, *Eur. J. Nucl. Med.* **1996**, *23*, 150-156; v) Z. Chen, W. Mori, X. Zhang, T. Yamasaki, P. J. Dunn, G. Zhang, H. Fu, T. Shao, Y. Zhang, A. Hatori, L. Ma, M. Fujinaga, L. Xie, X. Deng, H. Li, Q. Yu, J. Rong, L. Josephson, J.-A. Ma, Y. Shao, S. Tomita, M.-R. Zhang, S. H. Liang, *Eur. J. Med. Chem.* **2018**, *157*, 898-908.
- [7] a) D. Roeda, F. Hinnen, F. Dolle, *J. Labelled Compd. Radiopharm.* **2003**, *46*, 1141-1149; b) W. R. Banks, T. J. Tewson, G. A. Digenis, *Appl. Radiat. Isot.* **1990**, *41*, 719-725; c) U. Heiser, D. Ramsbeck, H.-U. Demuth (Probiobio AG, Germany), US20120301398A1, **2012** [*Chem. Abstr.* 158:1161]; d) M. Schou, C. Halldin, *J. Labelled Compd. Radiopharm.* **2012**, *55*, 460-462.
- [8] a) R. Neelamegam, T. Hellenbrand, F. A. Schroeder, C. Wang, J. M. Hooker, *J. Med. Chem.* **2014**, *57*, 1488-1494; b) D. Roeda, H. T. Sipilä, Y. Bramoulle, J. D. Enas, F. Vaufrey, F. Dolle, C. Crouzel, *J. Labelled Compd. Radiopharm.* **2002**, *45*, 37-47; c) M. W. Nader, S. K. Zeisler, A. Theobald, F. Oberdorfer, *Appl. Radiat. Isot.* **1998**, *49*, 1599-1603; d) M. Van Der Mey, A. D. Windhorst, R. P. Klok, J. D. M. Herscheid, L. E. Kennis, F. Bischoff, M. Bakker, X. Langlois, L. Heylen, M. Jurzak, J. E. Leysen, *Bioorg. Med. Chem.* **2006**, *14*, 4526-4534; e) M. Hanyu, Y. Takada, H. Hashimoto, K. Kawamura, M.-R. Zhang, T. Fukumura, *J. Pept. Sci.* **2013**, *19*, 663-668; f) M. Saeed, T. J. Tewson, C. E. Erdahl, A. Kohen, *Nucl. Med. Biol.* **2012**, *39*, 697-701; g) J. M. Hooker, M. Schoenberger, H. Shchlieferstein, J. S. Fowler, *Angew. Chem. Int. Ed.* **2008**, *47*, 5989-5992; h) G. Berger, M. Maziere, J. Sastre, D. Comar, *J. Labelled Compd. Radiopharm.* **1980**, *17*, 59-71; i) D. Roeda, F. Dollé, C. Crouzel, *J. Labelled Compd. Radiopharm.* **1997**, *40*, 725-726; j) E. H. Abraham (USA), WO2003099342A1, **2003** [*Chem. Abstr.* 140:2418]; k) W. B. Mathews, Y. Nakamoto, E. H. Abraham, U. Scheffel, J. Hilton, H. T. Ravert, M. Tatsumi, P. A. Rausero, B. J. Traugher, A. Y. Salikhova, R. F. Dannals, R. L. Wahl, *Mol. Imaging Biol.* **2005**, *7*, 203-208.
- [9] a) M. Ponchant, F. Hinnen, S. Demphel, C. Crouzel, *Appl. Radiat. Isot.* **1997**, *48*, 755-762; b) S. J. Lee, J. S. Fowler, D. Alexoff, M. Schueller, D. Kim, A. Nauth, C. Weber, S. W. Kim, J. M. Hooker, L. Ma, W. Qu, *Org. Biomol. Chem.* **2015**, *13*, 11235-11243; c) P. S. Kruijer, P. J. Van Leuffen, J. D. M. Herscheid, *Appl. Radiat. Isot.* **1996**, *47*, 611-616.
- [10] T. Haywood, S. Cesarec, S. Kealey, C. Plisson, P. W. Miller, *MedChemComm* **2018**, *9*, 1311-1314.
- [11] T. Haywood, S. Kealey, S. Sanchez-Cabezas, J. J. Hall, L. Allott, G. Smith, C. Plisson, P. W. Miller, *Chem. Eur. J.* **2015**, *21*, 9034-9038.
- [12] a) T. Kihlberg, F. Karimi, B. Langstrom, *J. Org. Chem.* **2002**, *67*, 3687-3692; b) F. Karimi, T. Kihlberg, B. Langstrom, *J. Chem. Soc., Perkin Trans. 1* **2001**, 1528-1531; c) O. Lindhe, P. Almqvist, M. Kaagedal, S.-A. Gustafsson, M. Bergstroem, D. Nilsson, G. Antoni, *Int. J. Mol. Imaging* **2011**, 694179, 694176 pp; d) S. Kealey, C. Plisson, T. L. Collier, N. J. Long, S. M. Husbands, L. Martarello, A. D. Gee, *Org. Biomol. Chem.* **2011**, *9*, 3313-3319; e) M. W. Nader, F. Oberdorfer, *Tetrahedron Lett.* **2011**, *52*, 2309-2311; f) J. Verbeek, J. Eriksson, S. Syvaenen, M. Labots, E. C. M. De Lange, R. A. Voskuyl, M. P. J. Mooijer, M. Rongen, A. A. Lammertsma, A. D. Windhorst, *EJNMMI Res.* **2012**, *2*, 36, 11 pp; g) S. Kealey, S. M. Husbands, I. Bennacef, A. D. Gee, J. Passchier, *J. Labelled Compd. Radiopharm.* **2014**, *57*, 202-208; h) T. Cornilleau, H. Audrain, A. Guillemet, P. Hermange, E. Fouquet, *Org. Lett.* **2015**, *17*, 354-357.
- [13] a) N. Vasdev, O. Sadovski, A. Garcia, F. Dolle, J. H. Meyer, S. Houle, A. A. Wilson, *J. Labelled Compd. Radiopharm.* **2011**, *54*, 678-680; b) A. Schirbel, M. H. Holschbach, H. H. Coenen, *J. Labelled Compd. Radiopharm.* **1999**, *42*, 537-551.
- [14] A. Del Vecchio, F. Caille, A. Chevalier, O. Loreau, K. Horkka, C. Halldin, M. Schou, N. Camus, P. Kessler, B. Kuhnast, F. Taran, D. Audisio, *Angew. Chem., Int. Ed.* **2018**, *57*, 9744-9748.
- [15] F. Liger, T. Eijsbouts, F. Cadarossanesaib, C. Tourvieille, D. Le Bars, T. Billard, *Eur. J. Org. Chem.* **2015**, 6434-6438.
- [16] a) O. Jacquet, C. Das Neves Gomes, M. Ephritikhine, T. Cantat, *ChemCatChem* **2013**, *5*, 117-120; b) X. Gao, B. Yu, Z. Yang, Y. Zhao, H. Zhang, L. Hao, B. Han, Z. Liu, *ACS Catal.* **2015**, *5*, 6648-6652.
- [17] a) Y. Bansal, O. Silakari, *Biorg. Med. Chem.* **2012**, *20*, 6208-6236; b) R. S. Keri, A. Hiremathad, S. Budagumpi, B. M. Nagaraja, *Chem. Biol. Drug Des.* **2015**, *86*, 19-65.
- [18] a) R. S. Keri, M. R. Patil, S. A. Patil, S. Budagumpi, *Eur. J. Med. Chem.* **2015**, *89*, 207-251; b) S. Tariq, P. Kamboj, M. Amir, *Arch. Pharm.* **2019**, *352*, 1800170.
- [19] a) J. P. Holland, S. H. Liang, B. H. Rotstein, T. L. Collier, N. A. Stephenson, I. Greguric, N. Vasdev, *J. Labelled Compd. Radiopharm.* **2014**, *57*, 323-331; b) N. S. Mason, C. A. Mathis, W. E. Klunk, *J. Labelled Compd. Radiopharm.* **2013**, *56*, 89-95.
- [20] A. Brayfield, *Martindale: The Complete Drug Reference*, Pharmaceutical Press, London, UK, **2014**.


COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Labeling inside. Intracyclic labeling with carbon-11 of heterocycles has been performed directly from cyclotron-produced [^{11}C]CO₂. Many heterocycles have been obtained via a de novo cyclization in only 10 min, without the preliminary transformation of [^{11}C]CO₂

**Radiochemistry***

*François Liger, Florence Cadarossanesaib, Thibaut Iecker, Christian Tourvieille, Didier Le Bars, Thierry Billard**

Page No. – Page No.

^{11}C labeling: Intracyclic incorporation of carbon-11 into heterocycles.

*one or two words that highlight the emphasis of the paper or the field of the study