

Communication

Enantioselective Synthesis of anti-1,2-Diamines by Cu-Catalyzed Reductive Couplings of Azadienes with Aldimines and Ketimines

Xinxin Shao, Kangnan Li, and Steven J. Malcolmson

J. Am. Chem. Soc., **Just Accepted Manuscript** • Publication Date (Web): 18 May 2018

Downloaded from <http://pubs.acs.org> on May 18, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Enantioselective Synthesis of *anti*-1,2-Diamines by Cu-Catalyzed Reductive Couplings of Azadienes with Aldimines and Ketimines

Xinxin Shao, Kangnan Li, and Steven J. Malcolmson*

Department of Chemistry, Duke University, Durham, NC 27708, United States

Supporting Information Placeholder

ABSTRACT: Here we report highly efficient and chemoselective azadiene–imine reductive couplings catalyzed by (Ph-BPE)Cu–H that afford *anti*-1,2-diamines. In all cases, reactions take place with either aldimine or ketimine electrophiles to deliver a single diastereomer of product in >95:5 er. The products' diamines are easily differentiable, facilitating downstream synthesis.

The catalytic enantioselective preparation of vicinal diamines is an important goal in synthetic chemistry owing to the large number of pharmaceuticals, natural products, and chiral ligands that contain this motif.¹ Although several approaches to this moiety have been reported by a number of researchers, significant shortcomings in scope or the ability to differentiate the products' two amino groups constrain their utility (Scheme 1). One major strategy has utilized intermolecular olefin diamination² to afford either the *anti*- or *syn*-1,2-diamines.³ In nearly all such cases, the two introduced amino groups have identical substituents, making their differentiation challenging to achieve.⁴ Another strategy has employed N-substituted enolates or nitroalkanes in Mannich-type reactions;^{5–7} either diastereomer may be selectively formed. However, in the former case, the requirement of an electron withdrawing group reduces the scope of diamines that may be prepared. In the latter, nitro group reduction is needed to secure the diamine.⁸ In both cases, when tetrasubstituted amine-containing stereogenic centers are formed, one of that center's other substituents has been limited to a carbonyl-like group.


To address these limitations, we sought to develop a method that would unite two N-containing reagents via catalytic enantioselective C–C bond formation such that: 1) a greater diversity of 1,2-diamines, including those with N-containing tetrasubstituted stereogenic centers, might be garnered; 2) the nitrogen groups of the products would be easily differentiated in order to assist in subsequent derivatizations; and 3) either free amine could be obtained without the need for harsh reducing conditions. We envisioned that the reductive coupling of 2-azadienes⁹ and suitably activated imines could allow us to realize this goal (Scheme 1). However, catalytic enantioselective reductive couplings with imines are rare.^{10,11} Successful implementation of our proposed strategy would require high catalyst efficiency and control over diastereo-, and enantio-, and chemoselectivity for the desired C–

C bond formation (versus imine reduction¹²).


Within the last several years, enantioselective Cu-catalyzed reductive couplings^{13,14} of unsaturated hydrocarbons with various C-electrophiles has rapidly emerged as an effective way for preparing myriad chemical motifs, often comprised of vicinal stereogenic centers. Vinylarenes,^{10c,d,15} vinylboronic esters,¹⁶ allenes,^{11e,17} and conjugated enynes¹⁸ and dienes^{15g,18} have comprised the substrates for these processes, yet none has established vicinal heteroatom-substituted stereogenic centers. Our recent disclosure of the Cu-catalyzed reductive coupling of 2-azadienes and ketones shows the promise these reagents hold for achieving such a goal.⁹ In this work, we demonstrate that 2-azadienes participate in chemoselective Cu-catalyzed reductive couplings with N-diphenylphosphinoyl (Dpp) imines. Both aldimines and ketimines react to furnish *anti*-1,2-diamines¹⁹ with vicinal stereogenic centers, in most cases as a single stereoisomer. The two N-groups of the products, one an imine and the other a phosphinamide, are readily discriminated, enabling their subsequent divergent elaboration.

Scheme 1. Catalytic Enantioselective Methods for Preparing Vicinal Diamines and Proposed Strategy


■ Olefin Diamination Methods



■ Mannich & Nitro-Mannich Reactions




■ Proposal: 2-Azadiene–Imine Reductive Couplings



We initially explored addition of terminal 2-azadiene **1a** to

Dpp-alimine **2a** (Table 1). Optimal conditions employ 3.0 equivalents of azadiene, DMMS as the reducing agent, *t*-BuOH as additive, a Cu-based catalyst with (*S,S*)-Ph-BPE as the ligand, and 5 °C (ice bath) reaction temperature (entry 1). After 1 h under these conditions, the desired diamine **3a** can be obtained in 82% yield, solely as the *anti*-diastereomer and as a single enantiomer. Accompanying **3a** is ca. 15% reduction product **4a**. Utilizing imine activating groups other than Dpp (e.g., Ts, Boc, etc.) results in <2% conversion to **3a** (entry 2).^{10d,12} Conducting the reaction at 22 °C results in poorer chemoselectivity, delivering more of the unwanted **4a**; however, stereoselectivity remains unaffected (entry 3). Omitting *t*-BuOH not only lowers catalyst efficiency but also adversely affects chemoselectivity (entry 4), similar to observations made by the Buchwald lab in styrene–imine couplings.^{10d} The identity of the alcohol additive is also critical for the selective formation of **3a** (entry 5). Although an *i*-Pr-BPE–Cu complex fails to furnish any product (entry 6), switching to *i*-Pr-DuPHOS generates **3a** in 40% yield, 10:1 dr, and 99:1 er but accompanied by a substantial quantity of **4a** (entry 7). A QuinoxP-derived catalyst, although highly selective for C–C bond formation coupling over imine reduction (8:1), generates **3a** in only 4:1 dr (entry 8).²⁰

Table 1. Impact of Reaction Conditions in Azadiene–Imine Reductive Couplings^a



entry	variation from the "standard" conditions	3a:4a ^b	yield of 3a		
			(%) ^c	dr ^b	er ^d
1	none	5:1	82	>20:1	>99:1
2	Ts, Boc, <i>et al.</i> instead of P(O)Ph ₂	–	–	–	–
3	22 °C	2:1	63	>20:1	>99:1
4	no <i>t</i> -BuOH	0.8:1	14	>20:1	>99:1
5	MeOH instead of <i>t</i> -BuOH	1.5:1	50	>20:1	>99:1
6	<i>i</i> -Pr-BPE instead of Ph-BPE	–	<5	–	–
7	<i>i</i> -Pr-DuPHOS instead of Ph-BPE	0.9:1	40	10:1	99:1
8	QuinoxP instead of Ph-BPE	8:1	59/16 ^e	4:1	99:1


^aReaction under N₂ with 0.1 mmol imine **2a**. ^bDetermined by 400 MHz ¹H NMR spectroscopy of the unpurified mixture. ^cIsolated yield of purified **3a**. ^dDetermined by HPLC analysis of purified **3a** major diastereomer. ^eYield of the major/minor diastereomer of **3a**.

Several aldimines undergo coupling with azadiene **1a**, leading to *anti*-diamines **3** as a single diastereomer (Table 2). In most cases, only a single enantiomer of product is generated. A variety of aryl-substituted imines participate in the reaction with the more electron rich substrates affording the highest yields (64–93%, entries 1–5, 13, 15). Halogen substituents are tolerated with diamines **3g–i** and **3o** isolated in 55–73% yield (entries 6–8, 14). More electron poor imines also yield the desired diamines **3j–l** (entries 9–11) but in somewhat diminished yields (41–59%). The observed trend is due to increasingly competitive imine reduction as the imine partner becomes more electron deficient;²¹ however, boronic ester **3m** is isolated in 75% yield as a single stereoisomer (entry 12). Notably,

more sterically hindered aldimines do not affect reaction efficiency: *o*-tolyl **3p** is furnished in 89% yield (entry 15). Heteroaryl aldimines can be coupled efficiently with the terminal azadiene as well to generate diamines **3q–s** in 83–94% yield (entries 16–18).

Unsaturated imines also undergo efficient coupling with azadiene **1a** (entries 19–20). Allylic amine **3t**, bearing a trisubstituted olefin, is formed in 71% yield. The less hindered cinnamyl **3u** is isolated in 61% yield but the reductive coupling also affords ca. 15% of saturated diamine. An alkyl-substituted imine leads to 52% yield of saturated diamine **3v** in >99:1 er (entry 21). An alkynyl aldimine failed to deliver the desired diamine product.


Table 2. Aldimine Scope in Couplings with Azadiene 1a



entry	product, R	yield (%) ^b	er ^c
1	3b , 4-Me ₂ NC ₆ H ₄	96	>99:1
2	3c , 4-MeOC ₆ H ₄	89	>99:1
3	3d , 4-F ₂ HCOCC ₆ H ₄	64	>99:1
4	3e , 4-MeSC ₆ H ₄	82	>99:1
5	3f , 4-(N-pyrazolyl)C ₆ H ₄	84	>99:1
6	3g , 4-FC ₆ H ₄	73	>99:1
7	3h , 4-ClC ₆ H ₄	62	>99:1
8	3i , 4-BrC ₆ H ₄	64	>99:1
9	3j , 4-MeO ₂ CC ₆ H ₄	59	>99:1
10	3k , 4-F ₃ CC ₆ H ₄	44	>99:1
11 ^d	3l , 4-NCC ₆ H ₄	41	>99:1
12 ^e	3m , 4-(pin)BC ₆ H ₄	75	>99:1
13	3n , 2-naphthyl	85	>99:1
14	3o , 3-BrC ₆ H ₄	55	>99:1
15	3p , 2-MeC ₆ H ₄	89	>99:1
16	3q , 4-furyl	93	>99:1
17	3r , 3-thiophenyl	83	>99:1
18	3s , 3-indolyl(NMe)	94	98.5:1.5
19	3t , C(Me)CHPh	71	>99:1
20 ^f	3u , CHCHPh	61	>99:1
21	3v , CH ₂ CH ₂ Ph	52	>99:1

^aReaction under N₂ with 0.2 mmol imine **2**. Dr measured by 400 MHz ¹H NMR spectroscopy of the unpurified mixture. ^bIsolated yield of purified **3**. ^cDetermined by HPLC analysis of purified **3**. ^d5.0 equiv **1a**. ^e4.0 equiv **1a**. ^fFormed as a 4:1 mixture of **3u**:**3v**; yield of isolated **3u**.


We also examined the coupling of 4-substituted 2-azadienes with aldimines to deliver diamines comprised of α -alkyl groups other than methyl. As typified in eq. 1, the added steric hindrance of azadiene **1b** leads to slower Cu–H insertion and a more competitive reduction, which adversely affects the



diamine yield. An electron rich aldimine, such as **2c**, and an extended reaction time (6 h) are required to obtain good yield of **3w**. Increasing to 5.0 equivalents of azadiene improves the reaction as well with the product then isolated in 56% yield (versus 42% with 3.0 equiv **1b**), >20:1 dr, and >99:1 er.

We next sought to test whether azadiene couplings with ketimines would enable the synthesis of 1,2-diamines wherein one stereogenic center is fully substituted. Reactions that form such motifs wherein both amines are bound to stereogenic centers, each with a variety of substituents, are rare and challenging to achieve. Therefore, we were pleased to find that terminal azadiene **1a** reacts with aryl-alkyl and diaryl ketimines to generate diamines **6a–h** in 84–92% yield (Table 3). With the exception of the less electrophilic, electron rich imine **5b** (entry 2), which requires higher temperature and longer reaction time, reactions proceed efficiently at 5 °C within 2 h. Transformations occur with >98% chemoselectivity for the reductive coupling regardless of imine identity. Remarkably, in all cases, the diamines are obtained in >20:1 dr (entries 1–7) and with high enantioselectivity. Notably, in addition to tolerating several aryl groups, the coupling is also permissible with longer chain alkyl groups (entry 7). The sterically encumbered benzophenone imine reacts smoothly to give diamine **6h** in 86% yield (entry 8).

Table 3. Couplings of Azadiene 1a with Ketimines^a



entry	product, Ar, R	yield (%) ^b	er ^c
1	6a , Ph, Me	90	96.5:3.5
2 ^d	6b , 4-MeOC ₆ H ₄ , Me	84	96:4
3	6c , 4-MeO ₂ CC ₆ H ₄ , Me	92	>99:1
4	6d , 4-F ₃ CC ₆ H ₄ , Me	89	>99:1
5	6e , 3-BrC ₆ H ₄ , Me	91	>99:1
6	6f , 3-ClC ₆ H ₄ , Me	87	>99:1
7	6g , Ph, (CH ₂) ₂ Ph	89	99:1
8	6h , Ph, Ph	86	>99:1


^aReaction under N₂ with 0.1 mmol imine **5**. ^{b-c}See Table 2. ^dReaction at 22 °C for 24 h.

Furthermore, ketimines also participate in reductive couplings with 4-substituted 2-azadienes, proceeding with ca. 60–70% chemoselectivity to furnish diamines **6i–r** as a single diastereomer and with high enantioselectivity (Table 4). Despite the steric congestion, reactions proceed to completion within 6 h at 5 °C. Product yields are improved with 10 mol % catalyst loading.²² Variation of the ketimine's aryl substituent, including both electron rich and electron poor arenes, is tolerated in couplings with azadiene **1b** (entries 1–5). The azadiene may contain several functional groups, including heterocycles, ethers, esters, and halides that are preserved in the products (44–63% yield, entries 6–10). The versatility of the reaction partners should enable the assembly of a range of complex molecules from these diamine building blocks.

The developed azadiene–imine reductive couplings have the advantage that the amines within the products are readily differentiated (Scheme 2) as one is doubly protected as an

imine (red) and the other mono-protected as the phosphinamide (blue). Either may be transformed to the amine by hydrolysis (i.e., without reduction). These qualities allow for selective functionalization of either product nitrogen. For example, deprotonation of the phosphinamide N–H of **3a** enables alkylation to deliver **7** in 84% yield while retaining the imine. Alternatively, the imine may be hydrolyzed under mildly acidic conditions and the resulting free amine then functionalized, such as in the formation of benzyl carbamate **8** (85% yield over two steps). The phosphinamide may then be cleaved with stronger acid, enabling functionalization of the liberated amine: phosphinamide **8** is converted to *t*-butyl carbamate **9** in 80% yield (two steps).


Table 4. Addition of 4-Substituted Azadienes to Ketimines^a



entry	product, Ar, R	yield (%) ^b	er ^c
1	6i , Ph, (CH ₂) ₂ Ph	62	96:4
2	6j , 4-MeO ₂ CC ₆ H ₄ , (CH ₂) ₂ Ph	56	99:1
3	6k , 4-ClC ₆ H ₄ , (CH ₂) ₂ Ph	62	>99:1
4	6l , 3,4-dioxolatoC ₆ H ₃ , (CH ₂) ₂ Ph	59	96:4
5	6m , 2-naphthyl, (CH ₂) ₂ Ph	72	98.5:1.5
6	6n , Ph, <i>n</i> -Bu	54	95.5:4.5
7	6o , Ph, (CH ₂) ₂ (3-thiophenyl)	63	97.5:2.5
8	6p , Ph, (CH ₂) ₃ OTBS	53	96.5:3.5
9	6q , Ph, (CH ₂) ₄ OBz	56	97:3
10	6r , Ph, (CH ₂) ₄ Cl	44	96:4

^aReaction under N₂ with 0.2 mmol imine **5**. ^{b-c}See Table 2.

Scheme 2. Utilizing the Products' Differentiated Amines



In this work, we have shown that reductive couplings of 2-azadienes with imines are an efficient and highly stereoselective way to construct vicinal diamines, several of which are difficult to access through existing protocols and have not before succumbed to enantioselective synthesis. The methodology

represents a rare example of enantioselective reductive couplings of imines as well as Cu-catalyzed reductive couplings to set vicinal heteroatom-substituted stereogenic centers. Our future efforts will concentrate on the further development of azadienes and their applications to chiral amine synthesis.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, analytical data for new compounds, and X-ray crystallographic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*steven.malcolmson@duke.edu

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We are grateful for financial support of this research from the NIH (GM124286), ACS Petroleum Research Fund (56575-DN1), and Duke University. We thank Dr. Roger Sommer (NC State) for X-ray crystallographic analysis.

REFERENCES

- (1) For a review, see: Lucet, D.; Le Gall, T.; Mioskowski, C. "The Chemistry of Vicinal Diamines," *Angew. Chem., Int. Ed.* **1998**, *37*, 2580.
- (2) For a review on enantioselective olefin diamination, see: (a) Zhu, Y.; Cornwall, R. G.; Du, H.; Zhao, B.; Shi, Y. "Catalytic Diamination of Olefins via N-N Bond Activation," *Acc. Chem. Res.* **2014**, *47*, 3665. For examples of enantioselective intermolecular reactions, see: (b) Du, H.; Yuan, W.; Zhao, B.; Shi, Y. "Catalytic Asymmetric Diamination of Conjugated Dienes and Triene," *J. Am. Chem. Soc.* **2007**, *129*, 11688. (c) Du, H.; Zhao, B.; Shi, Y. "Catalytic Asymmetric Allylic and Homoallylic Diamination of Terminal Olefins via Formal C-H Activation," *J. Am. Chem. Soc.* **2008**, *130*, 8590. (d) Cornwall, R. G.; Zhao, B.; Shi, Y. "Catalytic Asymmetric Synthesis of Cyclic Sulfamides from Conjugated Dienes," *Org. Lett.* **2013**, *15*, 796. (e) Muñiz, K.; Barreiro, L.; Romero, R. M.; Martínez, C. "Catalytic Asymmetric Diamination of Styrenes," *J. Am. Chem. Soc.* **2017**, *139*, 4354.
- (3) For ring-forming (intramolecular) enantioselective cases, see: (a) Sequeira, F. C.; Turnpenny, B. W.; Chemler, S. R. "Copper-Promoted and Copper-Catalyzed Intermolecular Alkene Diamination," *Angew. Chem., Int. Ed.* **2010**, *49*, 6365. (b) Ingalls, E. L.; Sibbald, P. A.; Kaminsky, W.; Michael, F. E. "Enantioselective Palladium-Catalyzed Diamination of Alkenes Using N-Fluorobenzenesulfonimide," *J. Am. Chem. Soc.* **2013**, *135*, 8854. (c) Turnpenny, B. W.; Chemler, S. R. "Copper-Catalyzed Alkene Diamination: Synthesis of Chiral 2-Aminomethyl Indolines and Pyrrolidines," *Chem. Sci.* **2014**, *5*, 1786. (d) Mizar, P.; Laverny, A.; El-Sherbini, M.; Farid, U.; Brown, M.; Malmedy, F.; Wirth, T. "Enantioselective Diamination with Novel Chiral Hypervalent Iodine Catalysts," *Chem. - Eur. J.* **2014**, *20*, 9910. (e) Fu, S.; Yang, H.; Li, G.; Deng, Y.; Jiang, H.; Zeng, W. "Copper(II)-Catalyzed Enantioselective Intramolecular Cyclization of N-Alkenylureas," *Org. Lett.* **2015**, *17*, 1018. (f) Wang, F.-L.; Dong, X.-Y.; Lin, J.-S.; Zeng, Y.; Jiao, G.-Y.; Gu, Q.-S.; Guo, X.-Q.; Ma, C.-L.; Liu, X.-Y. "Catalytic Asymmetric Radical Diamination of Alkenes," *Chem.* **2017**, *3*, 979.
- (4) For intermolecular diaminations where the two amines may be differentiated, see: (a) Simmons, B.; Walji, A. M.; MacMillan, D. W. C. "Cycle-Specific Organocascade Catalysis: Application to Olefin Hydroamination, Hydro-oxidation, and Amino-oxidation, and to Natural Product Synthesis," *Angew. Chem., Int. Ed.* **2009**, *48*, 4349. (b) Fu, R.

Zhao, B.; Shi, Y. "Synthesis of (+)-CP-99,994 via Pd(o)-Catalyzed Asymmetric Allylic and Homoallylic C-H Diamination of Terminal Olefin," *J. Org. Chem.* **2009**, *74*, 7577.

(5) For a review on Mannich reactions with N-substituted nucleophiles, see: (a) Arrayás, R. G.; Carretero, J. C. "Catalytic Asymmetric Direct Mannich Reaction: A Powerful Tool for the Synthesis of α,β -Diamino Acids," *Chem. Soc. Rev.* **2009**, *38*, 1940. For enantioselective examples, see: (b) Kobayashi, S.; Yazaki, R.; Seki, K.; Yamashita, Y. "The Fluorenone Imines of Glycine Esters and Their Phosphonic Acid Analogues," *Angew. Chem., Int. Ed.* **2008**, *47*, 5613. (c) Hernández-Toribio, J.; Arrayás, R. G.; Carretero, J. C. "Direct Mannich Reaction of Glycinate Schiff Bases with N-(8-Quinoyl)sulfonyl Imines: A Catalytic Asymmetric Approach to anti- α,β -Diamino Esters," *J. Am. Chem. Soc.* **2008**, *130*, 16150. (d) Kano, T.; Sakamoto, R.; Akakura, M.; Maruoka, K. "Stereocontrolled Synthesis of Vicinal Diamines by Organocatalytic Asymmetric Mannich Reaction of N-Protected Aminoacetaldehydes: Formal Synthesis of (-)-Agelastatin A," *J. Am. Chem. Soc.* **2012**, *134*, 7516. (e) Zhang, W.-Q.; Cheng, L.-F.; Yu, J.; Gong, L.-Z. "A Chiral Bis(betaine) Catalyst for the Mannich Reaction of Azlactones and Aliphatic Imines," *Angew. Chem., Int. Ed.* **2012**, *51*, 4085. (f) Lin, S.; Kawato, Y.; Kumagai, N.; Shibasaki, M. "Catalytic Asymmetric Mannich-Type Reaction of N-Alkylidene- α -Aminoacetonitrile with Ketimines," *Angew. Chem., Int. Ed.* **2015**, *54*, 5183. (g) Kondo, M.; Nishi, T.; Hatanaka, T.; Funahashi, Y.; Nakamura, S. "Catalytic Enantioselective Reaction of α -Aminoacetonitriles Using Chiral Bis(imidazoline) Palladium Catalysts," *Angew. Chem., Int. Ed.* **2015**, *54*, 8198. (h) Kano, T.; Kobayashi, R.; Maruoka, K. "Versatile In Situ Generated N-Boc-Imines: Application to Phase-Transfer-Catalyzed Asymmetric Mannich-Type Reactions," *Angew. Chem., Int. Ed.* **2015**, *54*, 8471.

(6) For enantioselective vinylogous Mannich reactions to prepare this motif, see: (a) Ranieri, B.; Curti, C.; Battistini, L.; Sartori, A.; Pinna, L.; Casiraghi, G.; Zanardi, F. "Diastereo- and Enantioselective Catalytic Vinylogous Mukaiyama-Mannich Reactions of Pyrrole-Based Silyl Dienolates with Alkyl-Substituted Aldehydes," *J. Org. Chem.* **2011**, *76*, 10291. (b) Silverio, D. L.; Fu, P.; Carswell, E. L.; Snapper, M. L.; Hoveyda, A. H. "N-Substituted Tertiary and O-Substituted Quaternary Carbon Stereogenic Centers by Site-, Diastereo- and Enantioselective Vinylogous Mannich Reactions," *Tetrahedron Lett.* **2015**, *56*, 3489.

(7) For examples of enantioselective nitro-Mannich reactions, see: (a) Yamada, K.-i.; Harwood, S. J.; Gröger, H.; Shibasaki, M. "The First Catalytic Asymmetric Nitro-Mannich-Type Reaction Promoted by a New Heterobimetallic Complex," *Angew. Chem., Int. Ed.* **1999**, *38*, 3504. (b) Yamada, K.-i.; Moll, G.; Shibasaki, M. "The First Enantioselective and Diastereoselective Catalytic Nitro-Mannich Reaction: A New Entry to Chiral Vicinal Diamines," *Synlett* **2001**, *SI*, 980. (c) Knudsen, K. R.; Risgaard, T.; Nishiwi, N.; Gothelf, K. V.; Jørgensen, K. A. "The First Catalytic Asymmetric Aza-Henry Reaction of Nitronates with imines: A Novel Approach to Optically Active β -Nitro- α -Amino Acid- and α,β -Diamino Acid Derivatives," *J. Am. Chem. Soc.* **2001**, *123*, 5843. (d) Nugent, B. M.; Yoder, R. A.; Johnston, J. N. "Chiral Proton Catalysis: A Catalytic Enantioselective Direct Aza-Henry Reaction," *J. Am. Chem. Soc.* **2004**, *126*, 3418. (e) Yoon, T. P.; Jacobsen, E. N. "Highly Enantioselective Thiourea-Catalyzed Nitro-Mannich Reactions," *Angew. Chem., Int. Ed.* **2005**, *44*, 466. (f) Singh, A.; Yoder, R. A.; Shen, B.; Johnston, J. N. "Chiral Proton Catalysis: Enantioselective Brønsted Acid Catalyzed Additions of Nitroacetic Acid Derivatives as Glycine Equivalents," *J. Am. Chem. Soc.* **2007**, *129*, 3466. (g) Trost, B. M.; Lupton D. W. "Dinuclear Zinc-Catalyzed Enantioselective Aza-Henry Reaction" *Org. Lett.* **2007**, *9*, 2023. (h) Singh, A.; Johnston, J. N. "A Diastereo- and Enantioselective Synthesis of α -Substituted syn- α,β -Diamino Acids," *J. Am. Chem. Soc.* **2008**, *130*, 5866. (i) Uruguchi, D.; Koshimoto, K.; Ooi, T. "Chiral Ammonium Betaines: A Bifunctional Organic Base Catalyst for Asymmetric Mannich-Type Reaction of α -Nitrocarboxylates," *J. Am. Chem. Soc.* **2008**, *130*, 10878. (j) Davis, T. A.; Wilt, J. C.; Johnston, J. N. "Bifunctional Asymmetric Catalysis: Amplification of Brønsted Basicity Can Orthogonally Increase the Reactivity of a Chiral Brønsted Acid," *J. Am. Chem. Soc.* **2010**, *132*, 2880. (k) Handa, S.; Gnanadesikan, V.; Matsunaga, S.; Shibasaki, M. "Heterobimetallic Transition Metal/Rare Earth Metal Bifunctional Catalysis: A Cu/Sm/Schiff Base Complex for Syn-Selective Catalytic Asymmetric

Nitro-Mannich Reaction," *J. Am. Chem. Soc.* **2010**, *132*, 4925. (l) Sprague, D. J.; Singh, A.; Johnston, J. N. "Diastereo- and Enantioselective Additions of α -Nitro Esters to Imines for *anti*- α,β -Diamino Acid Synthesis with α -Alkyl-Substitution," *Chem. Sci.* **2018**, *9*, 2336.

(8) For other enantioselective approaches to vicinal diamines, see: (a) Ooi, T.; Sakai, D.; Takeuchi, M.; Tayama, E.; Maruoka, K. "Practical Asymmetric Synthesis of Vicinal Diamines through the Catalytic Highly Enantioselective Alkylation of Glycine Amide Derivatives," *Angew. Chem., Int. Ed.* **2003**, *42*, 5868. (b) Kitagawa, O.; Yotsumoto, K.; Kohriyama, M.; Dobashi, Y.; Taguchi, T. "Catalytic Asymmetric Synthesis of Vicinal Diamine Derivatives through Enantioselective N-Alkylation Using Chiral π -Allyl Pd-Catalyst," *Org. Lett.* **2004**, *6*, 3605. (c) Trost, B. M.; Fandrick, D. R.; Brodmann, T.; Stiles, D. T. "Dynamic Kinetic Asymmetric Allylic Amination and Acyl Migration of Vinyl Aziridines with Imido Carboxylates," *Angew. Chem., Int. Ed.* **2007**, *46*, 6123. (d) Arai, K.; Lucarini, S.; Salter, M. M.; Ohta, K.; Yamashita, Y.; Kobayashi, S. "The Development of Scalable Multidentate Niobium Complexes as Catalysts for the Highly Stereoselective Ring Opening of *meso*-Epoxides and *meso*-Aziridines," *J. Am. Chem. Soc.* **2007**, *129*, 8103. (e) Yu, R.; Yamashita, Y.; Kobayashi, S. "Titanium(IV)/Tridentate BINOL Derivative as Catalyst for *meso*-Aziridine Ring-Opening Reactions: High Enantioselectivity, Strong Positive Non-Linear Effect and Structural Characterization," *Adv. Synth. Catal.* **2009**, *351*, 147. (f) Macdonald, M. J.; Hesp, C. R.; Schipper, D. J.; Pesant, M.; Beauchemin, A. M. "Highly Enantioselective Intermolecular Hydroamination of Allylic Amines with Chiral Aldehydes as Tethering Catalysts," *Chem. - Eur. J.* **2013**, *19*, 2597. (g) Wu, B.; Gallucci, J. C.; Parquette, J. R.; RajanBabu, T. V. "Bimetallic Catalysis in the Highly Enantioselective Ring-Opening Reactions of Aziridines," *Chem. Sci.* **2014**, *5*, 1102. (h) Uraguchi, D.; Kinoshita, N.; Kizu, T.; Ooi, T. "Synergistic Catalysis of Ionic Brønsted Acid and Photosensitizer for a Redox Neutral Asymmetric α -Coupling of N-Arylaminoethanes with Aldimines," *J. Am. Chem. Soc.* **2015**, *137*, 13768. (i) Izumi, S.; Kobayashi, Y.; Takemoto, Y. "Catalytic Asymmetric Synthesis of *anti*- α,β -Diamino Acid Derivatives," *Org. Lett.* **2016**, *18*, 696. (j) Chai, Z.; Yang, P.-J.; Zhang, H.; Wang, S.; Yang, G. "Synthesis of Chiral Vicinal Diamines by Silver(I)-Catalyzed Enantioselective Aminolysis of N-Tosylaziridines," *Angew. Chem., Int. Ed.* **2017**, *56*, 650. (k) Dumoulin, A.; Bernadat, G.; Masson, G. "Enantioselective Three-Component Amination of Enecarbamates Enables the Synthesis of Structurally Complex Small Molecules," *J. Org. Chem.*, **2017**, *82*, 1775. (l) Mwenda, E. T.; Nguyen, H. N. "Enantioselective Synthesis of 1,2-Diamines Containing Tertiary and Quaternary Centers through Rhodium-Catalyzed DYKAT of Racemic Allylic Trichloroacetimidates," *Org. Lett.* **2017**, *19*, 4814. (m) Perrotta, D.; Wang, M.-M.; Waser, J. "Lewis Acid Catalyzed Enantioselective Desymmetrization of Donor-Acceptor *meso*-Diaminocyclopropanes," *Angew. Chem., Int. Ed.* **2018**, *57*, 5120.

(9) Li, K.; Shao, X.; Tseng, L.; Malcolmson, S. J. "2-Azadienes as Reagents for Preparing Chiral Amines: Synthesis of 1,2-Amino Tertiary Alcohols by Cu-Catalyzed Enantioselective Reductive Couplings with Ketones," *J. Am. Chem. Soc.* **2018**, *140*, 598.

(10) For enantioselective examples see: (a) Ngai, M.-Y.; Barchuk, A.; Krische, M. J. "Enantioselective Iridium-Catalyzed Imine Vinylation: Optically Enriched Allylic Amines via Alkyne-Imine Reductive Coupling Mediated by Hydrogen," *J. Am. Chem. Soc.* **2007**, *129*, 12644. (b) Zhou, C.-Y.; Zhu, S.-F.; Wang, L.-X.; Zhou, Q.-L. "Enantioselective Nickel-Catalyzed Reductive Coupling of Alkynes and Imines," *J. Am. Chem. Soc.* **2010**, *132*, 10955. (c) Ascic, E.; Buchwald, S. L. "Highly Diastereo- and Enantioselective CuH-Catalyzed Synthesis of 2,3-Disubstituted Indolines," *J. Am. Chem. Soc.* **2015**, *137*, 4666. (d) Yang, Y.; Perry, I. B.; Buchwald, S. L. "Copper-Catalyzed Enantioselective Addition of Styrene-Derived Nucleophiles to Imines Enabled by Ligand-Controlled Chemoselective Hydrocupration," *J. Am. Chem. Soc.* **2016**, *138*, 9787. For a related redox neutral process, see: (e) Oda, S.; Franke, J.; Krische, M. J. "Diene hydroaminomethylation via Ruthenium-Catalyzed C-C Bond Forming Transfer Hydrogenation: Beyond Carbonylation," *Chem. Sci.* **2016**, *7*, 136.

(11) For non-enantioselective examples, see: (a) Townes, J. A.; Evans, M. A.; Queffelec, J.; Taylor, S. J.; Morken, J. P. "Stereoselective Synthesis of *trans* β -Lactams through Iridium-Catalyzed Reductive Coupling of Imines and Acrylates," *Org. Lett.* **2002**, *4*, 2537. (b) Kong,

J.-R.; Cho, C.-W.; Krische, M. J. "Hydrogen-Mediated Reductive Coupling of Conjugated Alkynes with Ethyl (N-Sulfinyl)iminoacetates: Synthesis of Unnatural α -Amino Acids via Rhodium-Catalyzed C-C Bond Forming Hydrogenation," *J. Am. Chem. Soc.* **2005**, *127*, 11269. (c) Komanduri, V.; Grant, C. D.; Krische, M. J. "Branch-Selective Reductive Coupling of 2-Vinyl Pyridines and Imines via Rhodium Catalyzed C-C Bond Forming Hydrogenation," *J. Am. Chem. Soc.* **2008**, *130*, 12592. (d) Zhu, S.; Lu, X.; Luo, Y.; Zhang, W.; Jiang, H.; Yan, M.; Zeng, W. "Ruthenium(II)-Catalyzed Regioselective Reductive Coupling of α -Imino Esters with Dienes," *Org. Lett.* **2013**, *15*, 1440. (e) Liu, R. Y.; Yang, Y.; Buchwald, S. L. "Regiodivergent and Diastereoselective CuH-Catalyzed Allylation of Imines with Terminal Allenes," *Angew. Chem., Int. Ed.* **2016**, *55*, 14077. (f) Lee, K. N.; Lei, Z.; Ngai, M.-Y. " β -Selective Reductive Coupling of Alkenylpyridines with Aldehydes and Imines via Synergistic Lewis Acid/Photoredox Catalysis," *J. Am. Chem. Soc.* **2017**, *139*, 5003. For related redox neutral processes, see: (g) Schmitt, D. C.; Lee, J.; Dechert-Schmitt, A.-M. R.; Yamaguchi, E.; Krische, M. J. "Ruthenium Catalyzed Hydroaminoalkylation of Isoprene via Transfer Hydrogenation: Byproduct-Free Prenylation of Hydatonins," *Chem. Commun.* **2013**, *49*, 6096. (h) Chen, T.-Y.; Tsutsumi, R.; Montgomery, T. P.; Volchkov, I.; Krische, M. J. "Ruthenium-Catalyzed C-C Coupling of Amino Alcohols with Dienes via Transfer Hydrogenation: Redox-Triggered Imine Addition and related Hydroaminoalkylations," *J. Am. Chem. Soc.* **2015**, *137*, 1798. (i) Oda, S.; Sam, B.; Krische, M. J. "Hydroaminomethylation Beyond Carbonylation: Allene-Imine Reductive Coupling by Ruthenium-Catalyzed Transfer Hydrogenation," *Angew. Chem., Int. Ed.* **2015**, *54*, 8525.

(12) For enantioselective Cu-catalyzed imine hydrosilylation, see: Lipshutz, B. H.; Shimizu, H. "Copper(I)-Catalyzed Asymmetric Hydrosilylations of Imines at Ambient Temperatures," *Angew. Chem., Int. Ed.* **2004**, *43*, 2228.

(13) For Cu-H reviews, see: (a) Rendler, S.; Oestreich, M. "Polishing a Diamond in the Rough: "Cu-H" Catalysis with Silanes," *Angew. Chem., Int. Ed.* **2007**, *46*, 498. (b) Lipshutz, B. H. "Rediscovering Organocopper Chemistry Through Copper Hydride. It's All About the Ligand," *Synlett* **2009**, 509. (c) Jordan, A. J.; Lalic, G.; Sadighi, J. P. "Coinage Metal Hydrides: Synthesis, Characterization, and Reactivity," *Chem. Rev.* **2016**, *116*, 8318.

(14) For reviews of catalytic reductive couplings with metals other than Cu, see: (a) Montgomery, J. "Nickel-Catalyzed Reductive Cyclizations and Couplings," *Angew. Chem., Int. Ed.* **2004**, *43*, 3890. (b) Hassan, A.; Krische, M. J. "Unlocking Hydrogenation for C-C Bond Formation: A Brief Overview of Enantioselective Methods," *Org. Process Res. Dev.* **2011**, *15*, 1236. (c) Standley, E. A.; Tasker, S. Z.; Jensen, K. L.; Jamison, T. F. "Nickel Catalysis: Synergy between Method Development and Total Synthesis," *Acc. Chem. Res.* **2015**, *48*, 1503. (d) Nguyen, K. D.; Park, B. Y.; Luong, T.; Sato, H.; Garza, V. J.; Krische, M. J. "Metal-Catalyzed Reductive Coupling of Olefin-Derived Nucleophiles: Reinventing Carbonyl Addition," *Science* **2016**, *354*, No. aah5133. (e) Kim, S. W.; Zhang, W.; Krische, M. J. "Catalytic Enantioselective Carbonyl Allylation and Propargylation via Alcohol-Mediated Hydrogen Transfer: Merging the Chemistry of Grignard and Sabatier," *Acc. Chem. Res.* **2017**, *50*, 2371.

(15) (a) Saxena, A.; Choi, B.; Lam, H. W. "Enantioselective Copper-Catalyzed Reductive Coupling of Alkenylazaarenes with Ketones," *J. Am. Chem. Soc.* **2012**, *134*, 8428. (b) Wang, Y.-M.; Bruno, N. C.; Placeres, A. L.; Zhu, S.; Buchwald, S. L. "Enantioselective Synthesis of Carbo- and Heterocycles through a CuH-Catalyzed Hydroalkylation Approach," *J. Am. Chem. Soc.* **2015**, *137*, 10524. (c) Wang, Y.-M.; Buchwald, S. L. "Enantioselective CuH-Catalyzed Hydroallylation of Vinylarenes," *J. Am. Chem. Soc.* **2016**, *138*, 5024. (d) Bandar, J. S.; Ascic, E.; Buchwald, S. L. "Enantioselective CuH-Catalyzed Reductive Coupling of Aryl Alkenes and Activated Carboxylic Acids," *J. Am. Chem. Soc.* **2016**, *138*, 5821. (e) Friis, S. D.; Pirnot, M. T.; Buchwald, S. L. "Asymmetric Hydroarylation of Vinylarenes Using a Synergistic Combination of CuH and Pd Catalysis," *J. Am. Chem. Soc.* **2016**, *138*, 8372. (f) Zhou, Y.; Bandar, J. S.; Buchwald, S. L. "Enantioselective CuH-Catalyzed Hydroacylation Employing Unsaturated Carboxylic Acids as Aldehyde Surrogates," *J. Am. Chem. Soc.* **2017**, *139*, 8126. (g) Gui, Y.-Y.; Hu, N.; Chen, X.-W.; Liao, L.-L.; Ju, T.; Ye, J.-H.; Zhang, Z.; Li, J.; Yu,

1 D.-G. "Highly Regio- and Enantioselective Copper-Catalyzed Reduc-
2 tive Hydroxymethylation of Styrenes and 1,3-Dienes with CO₂," *J. Am.*
3 *Chem. Soc.* **2017**, *139*, 17011. (h) Gribble, Jr., M. W.; Guo, S.; Buchwald,
4 S. L. "Asymmetric Cu-Catalyzed 1,4-Deaeromatization of Pyridines and
5 Pyridazines without Preactivation of the Heterocycle or Nucleophile,"
6 *J. Am. Chem. Soc.* **2018**, *140*, 5057.

7 (16) (a) Han, J. T.; Jang, W. J.; Yun, J. "Asymmetric Synthesis of
8 Borylalkanes via Copper-Catalyzed Enantioselective Hydroallylation,"
9 *J. Am. Chem. Soc.* **2016**, *138*, 15146. (b) Lee, J.; Torker, S.; Hoveyda, A.
10 H. "Versatile Homoallylic Boronates by Chemo- S_N2'-, Diastereo- and
11 Enantioselective Catalytic Sequence of Cu-H Addition to Vinyl-
12 B(pin)/Allylic Substitution," *Angew. Chem., Int. Ed.* **2017**, *56*, 821.

13 (17) Tsai, E. Y.; Liu, R. Y.; Yang, Y.; Buchwald, S. L. "A Regio- and
14 Enantioselective CuH-Catalyzed Ketone Allylation with Terminal Al-
15 lenes," *J. Am. Chem. Soc.* **2018**, *140*, 2007.

(18) Yang, Y.; Perry, I. B.; Lu, G.; Liu, P.; Buchwald, S. L. "Copper-
Catalyzed Asymmetric Addition of Olefin-Derived Nucleophiles to
Ketones," *Science* **2016**, *353*, 144.

(19) For azaallyl anion additions to imines that afford *syn*-1,2-dia-
mines, see: (a) Chen, Y.-J.; Seki, K.; Yamashita, Y.; Kobayashi, S. "Cat-
alytic Carbon-Carbon Bond-Forming Reactions of Aminoalkane De-
rivatives with Imines," *J. Am. Chem. Soc.* **2010**, *132*, 3244. (b) Matsu-
moto, M.; Harada, M.; Yamashita, Y.; Kobayashi, S. "Catalytic Imine-
Imine Cross-Coupling Reactions," *Chem. Commun.* **2014**, *50*, 13041.

(20) For additional screening data, see the Supporting Information.

(21) See the Supporting Information for further details.

(22) For example, with 5 mol % Cu and 6 mol % Ph-BPE, diamine
6i in Table 4, entry 1 is isolated in only 50% yield after 12 h.

Insert Table of Contents artwork here

