This article was downloaded by: [University of California, San Diego] On: 04 June 2015, At: 18:39 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: <a href="http://www.tandfonline.com/loi/lcar20">http://www.tandfonline.com/loi/lcar20</a>

# SYNTHESIS OF 2-(TRIMETHYLSILYL)ETHYL $\alpha$ -d-MANNOPYRANOSIDES REVISITED

Rina Saksena<sup>a</sup>, Jian Zhang<sup>a</sup> & Pavol Kováč<sup>b</sup> <sup>a</sup> NIDDK, LMC, National Institutes of Health, Bethesda, MD, 20892, U.S.A. <sup>b</sup> NIDDK, LMC, National Institutes of Health, Bethesda, MD, 20892, U.S.A. Published online: 16 Aug 2006.

To cite this article: Rina Saksena , Jian Zhang & Pavol Kováč (2002) SYNTHESIS OF 2-(TRIMETHYLSILYL)ETHYL  $\alpha$ -d-MANNOPYRANOSIDES REVISITED, Journal of Carbohydrate Chemistry, 21:6, 453-470

To link to this article: http://dx.doi.org/10.1081/CAR-120016846

## PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

JOURNAL OF CARBOHYDRATE CHEMISTRY Vol. 21, No. 6, pp. 453–470, 2002

## SYNTHESIS OF 2-(TRIMETHYLSILYL)ETHYL α-d-MANNOPYRANOSIDES REVISITED

Rina Saksena, Jian Zhang, and Pavol Kováč\*

NIDDK, LMC, National Institutes of Health, Bethesda, MD 20892, USA

#### ABSTRACT

Glycosylation of 2-(trimethylsilyl)ethanol with various ethyl 1-thioglycosides, which were activated with *N*-iodosuccinimide and silver triflate, was studied. The starting thioglycosides, some prepared for the first time, were obtained conventionally from the corresponding  $\alpha$ -1-acetates. When  $\beta$ -1acetates were more readily available, these were converted to the  $\alpha$ -anomers by anomerization, prior to the glycosylation. Using ethyl 1-thioglycosides as glycosyl donors, especially those bearing a pivaloyl or a nonparticipating group at O-2, the corresponding 2-(trimethylsilyl)ethyl  $\alpha$ -D-mannopyranosides were obtained in excellent yields.

*Key Words*: 2-(Trimethylsilyl)ethyl glycosides; Glycosylation; NIS/AgOTf; Anomerization

#### **INTRODUCTION**

The importance of 2-(trimethylsilyl)ethyl (SE) glycosides in the chemical synthesis of oligosaccharides lies in the fact that they are stable during many protecting group manipulations, yet the glycosidic 2-(trimethylsilyl)ethyl linkage can be cleaved under conditions which leave interglycosidic linkages in oligosaccharides (essentially) intact. A further desirable property of 2-(trimethylsilyl)ethyl glycosides is that they can be converted to an array of other useful glycosyl derivatives.<sup>[1–4]</sup> While

453

DOI: 10.1081/CAR-120016846 Copyright © 2002 by Marcel Dekker, Inc. 0732-8303 (Print); 1532-2327 (Online) www.dekker.com

<sup>\*</sup>Corresponding author.

454

#### SAKSENA, ZHANG, AND KOVÁČ

good yields of 2-(trimethylsilyl)ethyl glycosides of most common sugars can be obtained conventionally, synthesis of 2-(trimethylsilyl)ethyl  $\alpha$ -D-mannopyranosides is notoriously problematic.<sup>[1,5]</sup>

The formation of 2-(trimethylsilyl)ethyl  $\alpha$ -D-mannopyranosides from per-Oacylated glycosyl chlorides has been studied.<sup>[5]</sup> Fully acylated mannopyranosyl chlorides having different acyl groups at O-2 were treated with 2-(trimethylsilyl)ethanol (SEOH) under the conditions of silver trifluoromethanesulfonate (triflate, AgOTf) mediated Koenigs-Knorr reaction. The highest yield (72%) of the desired product was obtained from 3,4,6-tri-O-acetyl-2-O-benzoyl-α-D-mannopyranosyl chloride, prepared from 1,3,4,6-tetra-O-acetyl-2-O-benzoyl- $\beta$ -D-mannopyranose (26).<sup>a</sup> Since the yield of the fully benzoylated SE glycoside 11 from the corresponding glycosyl chloride was only 48%, the author<sup>[5]</sup> concluded that it was not the presence of the benzoyl group at O-2 alone, but "a proper ensemble of the protecting groups in a mannosyl donor" that minimizes side reactions and, as a result, affects the yield of 2-(trimethylsilyl)ethyl  $\alpha$ -Dmannopyranosides. The same communication reported that, unlike with D-mannose, a high yield of the corresponding 2-(trimethylsilyl)ethyl α-glycoside could be obtained from the fully benzoylated L-rhamnose. Accordingly, we have been able to prepare 2-(trimethylsilyl)ethyl 4-azido-3-O-benzyl-2-O-benzoyl-4,6-dideoxy-α-D-mannopyranoside (2),<sup>[6]</sup> from the corresponding glycosyl chloride, in 82% yield. However, the need for a large quantity of 2, in connection with our synthesis of perosaminecontaining oligosaccharides suitable for conjugation,<sup>[7]</sup> rendered even this yield not quite satisfactory. In an attempt to improve the yield, we have glycosylated SEOH using ethyl 4-azido-3-O-benzyl-2-O-benzoyl-4,6-dideoxy-1-thio-α-D-mannopyranoside (1) as a glycosyl donor and obtained the SE glycoside 2 in virtually theoretical yield. That result prompted us to undertake the present study, and here we describe the thioglycoside approach to making a number of useful 2-(trimethylsilyl)ethyl α-D-mannopyranosides. Because of the convenience<sup>[8]</sup> of its use, silver AgOTf/N-iodosuccinimide (NIS) reagent<sup>[9]</sup> was used as the promoter.

#### **RESULTS AND DISCUSSION**

Because of their versatility, thioglycosides have become very popular glycosyl donors in glycoside synthesis. Nevertheless, to our knowledge, thioglycosides have not been employed in syntheses of SE  $\alpha$ -D-mannopyranosides. The ethyl 1-thio-D-mannopyranosides used during this work were conveniently prepared from the corresponding  $\alpha$ -1-*O*-acetyl derivatives. In the case of **30** and **31**, the preparation was first carried out from the more readily available  $\beta$ -1-acetates **26** and **27**. When the yields of **30** and **31** were not quite satisfactory we turned to conversions of the  $\alpha$ -acetates **28** and **29**, as it has been reported<sup>[10]</sup> that the use of  $\alpha$ -anomers resulted in shorter reaction times and cleaner conversions to thioglycosides. In our situation, however, the benefit from the use of the  $\alpha$ -compounds was unimpressive. The  $\alpha$ -1-acetates **28** and **29**, were conveniently prepared from the  $\beta$ -anomers **26** and **27** by treatment with a mixture of AcOH, Ac<sub>2</sub>O and H<sub>2</sub>SO<sub>4</sub>,

<sup>&</sup>lt;sup>a</sup>No physical constants or spectral data were reported for this substance, and the compound, as well as its HO-2-free precursor, are erroneously shown (Ref. [5]) as  $\alpha$ -anomers.

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

455

which is generally used to effect acetolysis. An equilibrium mixture was formed, containing the  $\alpha$ - and  $\beta$ -anomers in a ratio of ~93:7 (NMR). The pure  $\alpha$ -anomers were then obtained by chromatography, which also gave an unresolved mixture of anomers containing a considerable amount of the  $\beta$ -anomer. The economy of preparation of the  $\alpha$ anomers could be improved by repeated treatment of this unresolved mixture of anomers with the acetolysis reagent (see above), to give product showing the same  $\alpha$ : $\beta \simeq 93:7$ ratio (NMR).



In the quest for improving the yield of **2** we treated ethyl 4-azido-2-*O*-benzoyl-4,6-dideoxy-1-thio- $\alpha$ -D-mannopyranoside (**1**) with the NIS/AgOTf reagent and compared the yield of the desired compound **2** with that obtained from 4-azido-2-*O*-benzoyl-4,6-dideoxy- $\alpha$ -D-mannopyranosyl chloride.<sup>[6]</sup> Monitoring the reaction by thinlayer chromatography (TLC) showed that 1.4 and 1.2 equivalents of NIS and AgOTf, respectively, was necessary to drive the reaction to completion, and that the conversion was a one-product reaction. After processing (see Experimental) and chromatography, the desired SE glycoside **2** was obtained in virtually theoretical yield.

The above results suggested that high yields of other SE mannopyranosides might be generally possible to obtain by way of 1-thioglycosides. Therefore, to ascertain if the thioglycoside functionality alone caused the dramatic change of the situation, glycosylation of SEOH with ethyl 2,3,4,6-tetra-*O*-acetyl-1-thio- $\alpha$ -D-mannopyranoside (4) was carried out (since we do not deem this reaction to be preparatively useful, this experiment is not described in the Experimental). Examination of the reaction mixture by TLC showed that several products were formed, and that the desired SE glycoside (7) was present in only a relatively small amount. The chromatographic mobility observed for two of the byproducts isolated from this conversion indicated that they were the hemiacetal (8), and 2-(trimethylsilyl)ethyl 3,4,6-tri-*O*-acetyl-1-thio- $\alpha$ -D-mannopyranoside (9). This corresponded to the previous observation<sup>[5]</sup> that products of partial deacetylation are formed during reactions of acetylated glycosyl donors with SEOH (cf., also formation of **20** during conversion **15**  $\rightarrow$  **19**). Conventional acetylation of **9** gave the peracetate **7** (TLC, NMR).

To imitate the substitution pattern at positions 1 and 2 in **1**, which gave the optimum yield of the desired product, reaction of ethyl 2,3,4,6-tetra-*O*-benzoyl-1-thio- $\alpha$ -D-mannopyranoside (**10**) with SEOH was examined next. Compound **10** was prepared from ethyl 1-thio- $\alpha$ -D-mannopyranoside<sup>[11,12]</sup> (**6**), here obtained crystalline for the first time. The NIS/triflic aciD-promoted<sup>[9]</sup> reaction of **10** with SEOH (not described in the Experimental) was rather sluggish and yielded a multicomponent mixture. The same conversion mediated with NIS/AgOTf<sup>[9]</sup> was virtually instantaneous, but 3.5 equivalents of the promoter were required to leave only trace of the starting thioglycoside unchanged. Analysis of the crude product by TLC showed the presence of two pro-

## 456

 $\mathbf{P}_{\mathrm{M}}$ 

## SAKSENA, ZHANG, AND KOVÁČ

*Table 1.* Amounts of Reagents Used (mmol) in Reactions of 1 mmol of the Starting Thioglycoside with 2 mmols of 2-(Trimethylsilyl)ethanol, and Yields (%) of Products

Thioglycoside	Product	NIS	AgOTf	Yield
N3 BnO SEt	N3 Bro OSE	1.2	1.4	$\sim 100$
1 Aco Aco SEt	2 AcO AcO ACO OSE	1.5	1.6	$\sim 20\%$
4 OBz Bzo Bzo SEt	7 OBz BzO BzO OSE	3.5	3.5	52.4
10 Pivo Pivo Pivo SEt	11 Pivo Pivo Pivo Pivo OSE	3.5	3.5	89
13 OBn Bno Aco	14 Bno Aco OSE	1.5	1.5	88.4
15 OBn BnO BnO BnO	19 Bno Bro Bzo OSE	1.5	1.5	85.2
17 OBn Bno Pivo SEt	22 Bno Pivo Bno OSE	1.5	1.9	87
18	23 Bno Pivo Bno OSE			3
	24			

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

457

	Table 1. Continued			
Thyoglycoside	Product	NIS	AgOTf	Yield
Aco Bzo SEt	ACO BZO OSE	2	2.2	73
30 OAc Aco Pivo SEt	32 OAc Aco Pivo	1.3	1.2	71
31 OBn BnO BnO BnO SEt	33 Bno Bno OSE	1.2	1.2	66
34	35 OBn BnO BnO OSE			32
	36			

ducts. After chromatography, the less polar product was shown to be the desired compound 11, obtained in 52.4% yield. NMR and MS analysis of the more polar material showed that it was the hemiacetal 12. The low yield of 11 from 10, which compared well with that obtained<sup>[5]</sup> from the corresponding glycosyl chloride, indicated that, as with 4 (*vide supra*), there was no advantage in the use of thioglycoside. Neither did the use of a thioglycoside as a glycosyl donor change the situation in the conversion  $30 \rightarrow 32$ . Improved yield of the corresponding SE glycoside was not obtained, compared to that from the glycosyl chloride.<sup>[5]</sup>

It has been suggested<sup>[5]</sup> that it is the hydrolysis of orthoesters, which are intermediates in glycoside formation from 2-*O*-acylated glycosyl donors, that causes the low yield of SE mannosides. Kunz<sup>[13]</sup> found that orthoester formation during glycosylation is suppressed when 2-*O*-pivaloyl glycosyl donors are used, unlike with other 2-*O*-acylated glycosyl donors. When we treated the fully pivaloylated 1-thioglycoside **13** with 2-(trimethylsilyl)ethanol (see Experimental), the desired, hitherto unknown, SE glycoside **14** was virtually the only product formed and, after chromatography, it was isolated crystalline in excellent yield of 87%. In view of the above it was somewhat disappointing to find that the yield of SE glycoside **33** from the 2-*O*-pivaloylated thioglycoside **31**, was not improved compared to that from **30**.

The potential utility of 2-(trimethylsilylethyl) 3,4,6-tri-O-benzyl- $\alpha$ -D-mannopyranoside (20) for making some (1  $\rightarrow$  2)-linked manno-oligosaccharides prompted us to examine glycosylation of SEOH with ethyl 2-O-acyl-3,4,6-tri-O-benzyl- $\alpha$ -1-thio-D-

## 458

2

## SAKSENA, ZHANG, AND KOVÁČ

mannopyranosides 15, 17, and 18. The yields of the corresponding SE glycosides from these glycosyl donors was high (>80%). The best yield was obtained from the conversion  $18 \rightarrow 23$ , which was virtually a one product reaction, yielding 87% of 23, after chromatography, along with a little of the corresponding  $\beta$ -anomer, 2-(trimethylsilyl)ethyl 2-O-pivaloyl-3,4,6-tri-O-benzyl-β-D-mannopyranoside 24.

		R <sup>4</sup> O <b>-</b> F	0R <sup>6</sup> 0 R <sup>2</sup> 0	$\mathbf{r}^{R^1}$		
	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	$\mathbf{R}^4$	R <sup>6</sup>
3	OAc	Н	Ac	Ac	Ac	Ac
4	SEt	Н	Ac	Ac	Ac	Ac
5	Н	SEt	Ac	Ac	Ac	Ac
6	SEt	Н	Н	Н	Н	Н
7	OSE	Н	Ac	Ac	Ac	Ac
8	Н	ЭН	Ac	Ac	Ac	Ac
9	OSE	Н	OH	Ac	Ac	Ac
10	SEt	Н	Bz	Bz	Bz	Bz
11	OSE	Н	Bz	Bz	Bz	Bz
12	HO	ЭН	Bz	Bz	Bz	Bz
13	SEt	Н	Piv	Piv	Piv	Piv
14	OSE	Н	Piv	Piv	Piv	Piv
15	SEt	Н	Ac	Bn	Bn	Bn
16	SEt	Н	Н	Bn	Bn	Bn
17	SEt	Н	Bz	Bn	Bn	Bn
18	SEt	Н	Piv	Bn	Bn	Bn
19	OSE	Н	Ac	Bn	Bn	Bn
20	OSE	Н	OH	Bn	Bn	Bn
21	HO	ЭН	Ac	Bn	Bn	Bn
22	OSE	Н	Bz	Bn	Bn	Bn
23	OSE	Н	Piv	Bn	Bn	Bn
24	Н	OSE	Piv	Bn	Bn	Bn
25	Н	Ac	Н	Ac	Ac	Ac
26	Н	Ac	Bz	Ac	Ac	Ac
27	Н	Ac	Piv	Ac	Ac	Ac
28	Ac	Н	Bz	Ac	Ac	Ac
29	Ac	Н	Piv	Ac	Ac	Ac
30	SEt	Н	Bz	Ac	Ac	Ac
31	SEt	Н	Piv	Ac	Ac	Ac
32	OSE	Н	Bz	Ac	Ac	Ac
33	OSE	Н	Piv	Ac	Ac	Ac
34	SEt	Н	Bn	Bn	Bn	Bn
35	OSE	Н	Bn	Bn	Bn	Bn
36	Н	OSE	Bn	Bn	Bn	Bn
37	Н	SEt	Bz	Ac	Ac	Ac
38	Н	SEt	Piv	Ac	Ac	Ac

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

459

Finally, we deemed it interesting to examine the formation of SE glycosides from the fully benzylated mannosyl donor, ethyl 2,3,4,6-tetra-*O*-benzyl-1-thio- $\alpha$ -D-mannopyranoside (34). Since it contains only alkyl protecting groups, orthoester formation during SE-glycoside formation can not occur. The conversion  $34 \rightarrow 35 + 36$  was high yielding, but produced the SE glycosides with impaired stereoselectivity.

Outcome of the reactions described above, together with the results of others<sup>[1,5]</sup> confirm that formation of SE  $\alpha$ -D-mannopyranosides is not a simple process. Although it is not possible to make a simple generalization, the following observations have been made. Firstly, the use of thioglycosides as glycosylating reagents for SEOH alone can be beneficial, compared to the use of glycosyl chlorides.<sup>[5,6]</sup> For example, in the conversion  $1 \rightarrow 2$ , the use of ethyl 1-thioglycoside as the glycosyl donor, even with the 2-O-benzoyl protection, resulted in a one-product reaction and a high yield of the desired product. On the other hand, the use of thioglycosides had no significant effect on the yield of 7 and 11 (see Experimental), compared to the use of glycosyl chlorides.<sup>[5,6]</sup> Secondly, variously protected ethyl 1-thio  $\alpha$ -D-mannopyranosides show different reactivity when treated with the NIS/AgOTf reagent. Therefore, the amount of the NIS/ AgOTf reagent needed to drive the reaction to completion varied with the substitution pattern, and it had to be determined for each ethyl 1-thioglycoside (Table 1). Presumably, the same holds also for other 1-thioglycosides. Thirdly, a comparison of yields of the corresponding SE glycosides obtained from thioglycosides 4 and 13, respectively, 52.4 and 89%, seems to suggest that formation and subsequent hydrolysis of orthoesters during the formation of SE glycosides<sup>[5]</sup> is a factor that influences the yield of SE mannopyranosides. It is, however, important to note in this context that glycosylation of SEOH with 2-O-acetylated glycosyl donors does not necessarily result in a poor yield of SE mannopyranosides (cf. the conversion  $15 \rightarrow 19$ ). Neither does the presence of the O-pivaloyl group at position 2 in a glycosyl donor guarantee excellent yield of the corresponding SE mannopyranoside (cf. the conversion  $31 \rightarrow 33$ ). Thus, besides simple orthoester formation and its hydrolysis, other phenomena must be involved, which affect the yields of SE  $\alpha$ -mannopyranosides. The results here described suggest that these yields are affected not only by the nature of the acyl group at O-2, but also by its presence/absence at position 6. This can be confidently concluded from the yields of conversions of the tri-O-benzylated donors 15, 17, and 18, which all gave very good yields of the corresponding SE glycosides, as well as from the comparison in the yields of SE  $\alpha$ -glycosides obtained from the mannoside **10** (52.4 or 48%<sup>[5]</sup>), rhamnose<sup>[5]</sup> (86%), **1** ( $\sim$ 100%), or the corresponding glycosyl chloride<sup>[6]</sup> (82%).

#### EXPERIMENTAL

**General methods.** Unless stated otherwise, optical rotations were measured at ambient temperature for solutions in chloroform  $(c \sim 1)$ , with a Perkin–Elmer automatic polarimeter, Model 341. All reactions were monitored by TLC on Silica gel 60 coated glass slides (Whatman or Analtech). The judgment expressed in Results and Discussion concerning the quality of individual reactions is based mainly on TLC and not on the actual yields. Thus, a reaction is occasionally described as a "one-product reaction," even though the yield reported for that reaction is <100%, which reflects manipulative losses and not incomplete reactions or formation of by-products. Column chromatography was performed by gradient elution from columns of silica gel. Solvent

	Table 2.	1H NMR Chemica	l Shifts <sup>a</sup> and Peak	t Multiplicities for	or Compounds 1,5	-7,10-14,16,18,19,22-2	4, and 26–38		460
Compound	H-1	H-2	H-3	H-4	H-5	H-6	$SCH_2CH_3$	$SCH_2CH_3$	)
$1^{\mathrm{b}}$	5.34d	5.63dd	3.85dd	3.55t	4.20 - 3.90 m	1.38d	2.72-2.52m	1.28t	
л; Л	4.80d	5.52dd	5.08dd	5.26t	3.72ddd	4.28dd, 4.14dd	2.74q	1.31t	
<b>6</b> d	5.29bd	4.02dd	3.75, dd, po	3.63t	3.97ddd, po	3.86dd, 3.73dd, po	2.72-2.55m	1.24t	
7°	4.80d	5.17dd	5.33	5.23t	3.97ddd	4.23dd, 4.08dd			
10	5.58bd	$\sim 5.82$ dd,po	5.84dd, po	6.16	4.84ddd	4.69dd, 4.53dd	2.84-2.66m	1.35t	
$11^{\mathrm{f}}$	5.13d	5.68dd	5.94dd	6.09t	$\sim$ 4.44m, po	4.70m, 4.48dd, po			
<b>12</b> <sup>g</sup>	5.53d	5.75dd	6.02dd	6.19t	4.68dt	4.78dd, 4.43dd			
$13^{\rm h}$	5.24bd	5.34dd	5.30dd	5.51t	4.44ddd	4.23dd, 4.12dd	2.75-2.57m	1.31t	
<b>14</b> <sup>i</sup>	4.79d	5.20dd	5.39dd	5.47t	4.05ddd	4.20dd, 4.13dd			
<b>16</b> <sup>j</sup>	5.38d	4.08dd	3.83dd, po	3.90t	4.16ddd	3.79dd, po, 3.67dd	$2.72 - 2.50 \mathrm{m}$	1.27t	
<b>16</b> <sup>k</sup>	5.53bd	4.13dd	3.94dd	4.08t	4.41mpo	3.82dd, 3.69dd	2.52–2.27m	1.07t	
<b>18</b> <sup>1</sup>	5.30d	5.44bt	$3.95 - 3.87 \mathrm{m}$	3.95 - 3.87 m	$4.20 - 4.13 \mathrm{m}$	3.82dd, 3.71dd	2.73–2.53m	1.28t	
<b>18</b> <sup>m</sup>	5.41d	5.74	4.10dd	4.16t	4.41m, po	3.84dd, 3.66dd	2.45-2.25m	1.04t	
<b>19</b> <sup>n</sup>	4.94d	5.69dd	4.22dd	4.14t	4.07m	3.83dd, 3.74dd			
$22^{\circ}$	5.01d	5.94dd	$4.35 - 4.30 \mathrm{m}$	$4.35 - 4.30 \mathrm{m}$	4.11ddd	3.92dd, 3.78dd			
<b>23</b> <sup>p</sup>	4.82d, po	5.31dd	3.97dd	3.85 - 3.69 m	3.85 - 3.69 m	$3.85 - 3.69 \mathrm{m}$			
24 <sup>q</sup>	4.50d, po	5.56dd	3.65dd	3.82–3.72m	3.47ddd	3.82–3.72m			S
$26^{\circ}$	6.00d	5.75dd	5.28dd	5.46t	3.92ddd	4.36dd, 4.22dd			SAI
$27^{\rm s}$	5.89d	5.45dd	5.13dd	5.33t	<b>3.83ddd</b>	4.29dd, 4.16dd			KSI
28 <sup>t</sup>	6.41d	5.81dd	5.73dd	5.89t	4.10ddd	4.40dd, 4.16dd			EN
<b>29</b> <sup>u</sup>	6.08d	5.26dd	5.34dd	5.43t	4.06ddd	4.27dd, 4.12dd			A,
$30^{\vee}$	5.44bd	5.59dd	5.38dd	5.49t	4.46ddd	4.35dd, 4.16dd	2.78–2.59m	1.33t	ZH
<b>31</b> <sup>w</sup>	5.28d, po	5.34dd	5.25dd, po	5.37t, po	4.39ddd	4.30dd, 4.12dd	2.76–2.57m	1.32t	IAI
32 <sup>×</sup>	4.87bd	5.90-5.86m	5.90-5.86m	5.90-5.86m	4.17m	4.43dd, 4.32dd			NG
<b>33</b> <sup>y</sup>	4.83d	5.20dd	5.38-5.29m	5.38–5.29m	4.01ddd	4.24dd, 4.14dd			, A
$34^{z}$	5.50d	3.94dd	4.07dd	4.30t	$\sim$ 4.42m	3.86dd, 3.73dd	2.58–2.35m	1.12t	NI
<b>35</b> <sup>aa</sup>	5.07d	$\sim$ 3.94, po	4.22dd	4.33t	4.15ddd	4.01–3.91m, 3.85dd			) K
$36^{\mathrm{bb}}$	4.39d	3.87dd, po	3.50dd, po	3.87t, po	$\sim$ 3.45m, po	3.82dd, 3.74dd			0
37 <sup>cc</sup>	4.89d	5.76dd	5.18dd	5.37t	3.78ddd	4.33dd, 4.20dd	2.75q	1.30t	VÁ
$38^{\rm dd}$	4.81d	5.50dd	5.06dd	5.29t	3.72ddd	4.28dd, 4.15dd	2.74q	1.30t	Ċ

MARCEL DEKKER, INC. • 270 MADISON AVENUE • NEW YORK, NY 10016 ©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

46

, .

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

461

mixtures slightly less polar than those used for TLC were used at the onset of elution. Chemical ionization mass spectra (CIMS, positive or negative ion) were obtained on a Finnigan 4000 quadrupole mass spectrometer with ammonia reagent gas. Fast atom bombardment (FAB) mass data were obtained using a JEOL SX-102 mass spectrometer

Notes to Table 2: <sup>a</sup>Measured in CDCl<sub>3</sub>, unless stated otherwise; d, doublet; t, triplet; q, quartet; m, multiplet: b, broad; po, partially overlapped; \*not determined due to overlapping signals.  $^{b}\delta_{PhCH_{2}}$  4.73, 4.55 (2 d  $^{2}J$  11.2 Hz). <sup>c</sup>δ<sub>COCH</sub>, 2.19s, 2.08s, 2.05s, 1.99s. <sup>d</sup>Measured in D<sub>2</sub>O.  $^{e}\delta_{OCH,CH,Si}$  3.76ddd J 6.2 and 9.7;  $\delta_{OCH,CH,Si}$  3.51ddd;  $\delta_{CH,CH,Si}$  0.94m;  $\delta_{(CH,),Si} \sim 0.0s$ ;  $\delta_{COCH,CH,Si}$ 2.12s, 2.06s, 2.01s, 1.96s.  $^{t}\delta_{CH,CH,Si}$  3.95dt, po, J 10.6 Hz;  $\delta_{CH,CH,Si}$  3.68dt, po, J 10.0 Hz;  $\delta_{CH,CH,Si}$  1.09m;  $\delta_{(CH_{3})}$ Si 0.06s. <sup>g</sup>Data for the  $\alpha$ -anomer. Measured after addition of D<sub>2</sub>O.  $^{h}\delta_{C(CH_{3})_{3}}$  1.28s, 1.24s, 1.17s, 1.12s.  $\delta_{\text{OCH,CH,Si}}$  3.82, 3.80 (2 t, J 10.0);  $\delta_{\text{OCH,CH,Si}}$  3.56, 3.54 (2 t, J 10.0);  $\delta_{\text{CH,CH,Si}}$  0.99m;  $\delta_{\text{(CH,)Si}}$  $\sim 0.03$ s;  $\delta_{C(CH_2)}$ , 1.27, 1.24, 1.16, 1.12, 4 s.  $^{J}\delta_{PhCH_{2}}$  4.84–4.48m. <sup>k</sup>Measured in C<sub>6</sub>D<sub>6</sub>.  $\delta_{PhCH_2}$  4.89, 4.59, 2d, <sup>2</sup>J 11.3 Hz; 4.53, 4.40, 2d, <sup>2</sup>J 12.0 Hz; 4.28s.  $^{1}\delta_{PhCH_{2}}$  4.85–4.46m;  $\delta_{COCH_{2}}$  1.23s. <sup>m</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  5.03–4.34m;  $\delta_{COCH_3}$  1.25s. <sup>n</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  5.05–4.38m; the signal of OCH<sub>a</sub>CH<sub>2</sub>Si appears as 2 t ( $\delta$  3.83, 3.76), po with the H-6a signal; the signal of OCH<sub>b</sub>CH<sub>2</sub>Si appears as 2 t at  $\delta$  3.39 and 3.37, J 9.7 Hz;  $\delta_{CH_2CH_2Si}$  $0.94 - 0.81 \text{m}; \ \delta_{(CH_2),Si} \sim -0.04 \text{s}.$ <sup>o</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  4.81–4.41m; the signal of OCH<sub>a</sub>CH<sub>2</sub>Si appears as 2 t ( $\delta$  3.84, 3.82), po with the H-6b signal; the signal of OCH<sub>b</sub>CH<sub>2</sub>Si appears as 2 t at  $\delta$  3.41 and 3.38, J 9.3 Hz;  $\delta_{CH_2CH_2Si}$ 0.95–0.80m;  $\delta_{(CH_3)_3}Si \sim -0.03s$ .  ${}^{P}\delta_{PhCH_{2}}$  4.83–4.44m; the signal of OCH<sub>a</sub>CH<sub>2</sub>Si is part of m at 3.85–3.69;  $\delta_{CH_{2}CH_{2}Si}$  3.49, 3.47 (2 t, J 9.8 Hz);  $\delta_{C(CH_3)_3}$  1.19s;  $\delta_{CH_2CH_2Si}$  0.89m;  $\delta_{(CH_3)_3}Si - 0.02s$ . <sup>q</sup>δ<sub>PhCH</sub>, 4.87–4.44m; δ<sub>OCH CH<sub>2</sub>Si</sub> 3.99, 3.97 (2 t, J 9.3 Hz); δ<sub>OCH-CH<sub>2</sub>Si</sub> 3.58, 3.56 J 9.3 Hz); δ<sub>CH-CH<sub>2</sub>Si</sub> 0.93ddd, po, J 7.1 and 8.3;  $\delta_{C(CH_3)_3}$  1.24s;  $\delta_{(CH_3)_3Si}$  0.01s. <sup>r</sup>δ<sub>COCH<sub>3</sub></sub> 2.15s, 2.07s, 2.05s, 1.99s. <sup>s</sup>δ<sub>COCH<sub>3</sub></sub> 2.09s, 2.08s, 2.05s, 1.99s. <sup>t</sup>Measured in C<sub>6</sub>D<sub>6</sub>.  $\delta_{COCH_2}$  1.79s, 1.68s, 1.63s, 1.50s.  ${}^{u}\delta_{COCH_{3}}$  2.18s, 2.09s, 2.05s, 1.99s;  $\delta_{C(CH_{3})_{3}}$  1.28s. <sup>v</sup>δ<sub>COCH<sub>2</sub></sub> 2.13s, 2.07s, 1.97s.  $^{\rm w}\delta_{\rm COCH_2}$  2.09s, 2.05s, 1.97s, 1.99s;  $\delta_{\rm C(CH_2)_2}$  1.28s. <sup>x</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{COCH_1}$  1.82s, 1.68s, 1.61s;  $\delta_{OCH_2}$  CH<sub>2</sub>Si, 3.69m;  $\delta_{OCH_2}$  CH<sub>2</sub>Si, 3.30m;  $\delta_{CH_2}$  CH<sub>2</sub>Si 0.89m;  $\delta_{(CH_2)_2}$ Si ~ 0.01s.  ${}^{y}\delta_{COCH_{3}}$  2.09s, 2.04s, 1.97s;  $\delta_{OCH_{4}CH_{3}Si}$  3.81ddd;  $\delta_{OCH_{4}CH_{3}Si}$  3.57ddd;  $\delta_{CH_{2}CH_{3}Si}$  0.99m;  $\delta_{(CH_{4})_{3}Si}$ 0.04s;  $\delta_{C(CH_2)_2}$  1.27. <sup>z</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{CH,Ph}$  4.99–4.34m, partially overlapped. <sup>aa</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  5.03–4.48m; the signal of OCH<sub>a</sub>CH<sub>2</sub>Si is part of m at 4.01–3.83; the signal of OCH<sub>b</sub>CH<sub>2</sub>Si appears as m at 3.53–3.44;  $\delta \delta_{CH,CH,Si}$  0.96–0.90m;  $\delta_{(CH_2),Si} \sim 0.16s$ .  $^{bb}\delta_{PhCH_2}$  5.02–4.39m;  $\delta_{OCH_a}CH_2Si$  4.12–4.03m; the signal of  $OCH_bCH_2Si$  is part of m at 3.55– 3.43;  $\delta_{CH_2CH_2Si}$  1.0t;  $\delta_{(CH_3)_3Si} \sim 0.03s$ . <sup>cc</sup>δ<sub>COCH<sub>3</sub></sub> 2.12s, 2.05s, 1.97s.  $^{dd}\delta_{\text{COCH}_3}$  2.08s, 2.04s, 1.96s.

## 462

## SAKSENA, ZHANG, AND KOVÁČ

utilizing a glycerol matrix. Assignments of NMR signals (Tables 2–4), obtained at 300 MHz for <sup>1</sup>H and 75 MHz for <sup>13</sup>C at 25°C, were made by first-order analysis of spectra and, when feasible, the assignments were supported by APT and/or DEPT experiments, homonuclear decoupling and/or homo- and heteronuclear 2-dimensional correlation spectroscopy. The commercial software supplied with the spectrometers (Varian Gemini or Varian XL 300) was used. The  $\beta$ -anomeric configuration of substances was ascertained by <sup>1</sup>H–<sup>13</sup>C coupling constants<sup>[14]</sup>  $J_{C-1,H-1}$ . Solutions in organic solvents were dried with anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated at 40°C/2kPa.

*Table 3.* <sup>1</sup>H NMR Coupling Constants<sup>a</sup> for Compounds 1, 5–7, 10–14, 16, 18, 19, 22–24, and 26–38

	Coupling Constants (Hz)						
Compound	J <sub>1,2</sub>	$J_{2,3}$	$J_{3,4}$	$J_{4,5}$	$J_{5,6a}$	$J_{5,6b}$	$J_{6a,6b}$
1	1.6	3.2	9.9	9.9	6.2		
5	1.1	3.5	10.1	10.1	6.1	2.6	12.4
6	$\sim 1.2$	3.3	$\sim 9.8$	$\sim 9.8$	2.7	6.4	12.5
7	1.4	3.1	10.2	10.2	6.0	2.3	12.5
10	$\sim 1.0$	3.1	$\sim 10.0$	$\sim 10.0$	2.6	4.5	12.4
11	1.8	3.3	$\sim 10.3$	$\sim 10.3$	*	5.0	$\sim 13$
12	1.8	3.1	10.2	$\sim 10$	2.6	3.5	12.2
13	1.6	3.3	9.9	10.1	4.4	1.9	12.5
14	1.7	3.1	9.9	9.6	4.4	2.0	12.5
16	1.4	3.1	9.3	9.3	4.5	1.9	11.1
<b>16</b> <sup>b</sup>	1.5	3.6	9.2	9.2	4.8	1.7	10.9
18	1.5	*	*	*	4.3	1.9	10.9
<b>18</b> <sup>b</sup>	1.8	2.9	9.2	9.2	4.6	1.5	11.0
<b>19</b> <sup>b</sup>	1.8	3.3	9.2	$\sim 9.2$	4.6	1.7	10.8
22 <sup>b</sup>	1.8	2.6	*	9.7	4.6	1.7	10.8
23	1.9	3.3	9.1	*	*	*	*
24	$\sim 1.0$	3.0	9.3	9.3	2.3	4.3	*
26	0.9	3.5	10.3	$\sim 10$	5.5	2.3	13.0
27	1.2	3.2	10.1	9.8	5.0	2.2	12.7
28	2.0	3.3	10.2	10.2	4.7	2.5	12.3
29	2.1	3.0	10.0	10.0	4.0	2,4	12.2
30	1.8	3.5	10.0	10.0	5.0	2.4	12.3
31	$\sim 1.6$	3.1	10.0	9.9	4.7	2.3	12.2
32	$\sim 1.5$	*	*	*	4.9	2.5	12.1
33	1.9	2.9	*	9.9	5.3	2.6	12.4
<b>34</b> <sup>b</sup>	1.4	3.2	10.0	10.0	5.2	1.6	10.9
35 <sup>b</sup>	$\sim 1.4$	3.1	9.3	9.6	*	3.8	$\sim 10.7$
36	0.8	3.1	9.3	9.3	2.0	5.9	10.6
37	1.1	3.5	10.0	10.1	5.5	2.5	12.2
38	0.9	3.3	10.0	10.0	5.6	2.2	12.7

<sup>a</sup>Measured in CDCl<sub>3</sub>, unless stated otherwise.

<sup>b</sup>Measured in C<sub>6</sub>D<sub>6</sub>.

\*Not determined due to overlap, or broadening of signal, or second order effect.

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

463

NIS and AgOTf were purchased from Aldrich Chemical Company. Before use, NIS was purified by crystallization from dioxane-CCl<sub>4</sub>, and dried at  $80^{\circ}$ C/2 kPa for 2 h, mp  $201-202^{\circ}$ C. Ethyl 2,3,4,6-tetra-O-acetyl-1-thio- $\alpha$ , $\beta$ -D-mannopyranoside was obtained as described,<sup>[15]</sup> except that the reaction was performed at room temperature. Essentially the same anomeric mixture was formed within 3-4 h as that described<sup>[15]</sup> with the reaction time 6–7 days. Chromatography and crystallization yielded the  $\alpha$ -anomer 4, mp 106– 107°C (from EtOH), ref. 16, mp 107-108°C. The β-anomer 5 had mp 160-161°C (from EtOH) and  $[\alpha]_{\rm D} - 64^{\circ}$  (c 0.9, CHCl<sub>3</sub>), in accord with literature values reported by Fried and Waltz.<sup>[12]</sup> The NMR data, hitherto not reported, are in Tables 2-4. Ethyl 2,3,4,6tetra-O-benzoyl-1-thio- $\alpha$ -D-mannopyranoside (10) was obtained from 5 as described by Sarbajna and Roy.<sup>[11]</sup> Ethyl 2,3,4,6-tetra-O-benzyl-1-thio-α-D-mannopyranoside (**34**) was obtained from **6** as described by Helander et al.<sup>[16]</sup> 1,3,4,6-Tetra-O-acetyl-β-D-mannopyranose (25) was prepared as reported by Bovin et al.<sup>[17]</sup> Ethyl 2-O-benzoyl-3,4,6-tri-Obenzyl-1-thio- $\alpha$ -D-mannopyranoside (17) was prepared according to Zhang et al.<sup>[18]</sup> Ethyl 2-O-acetyl-3,4,6-tri-O-benzyl-1-thio- $\alpha$ -D-mannopyranoside (15) was obtained as described.<sup>[19]</sup> The compound crystallized readily from isopropyl ether containing a few drops of hexane, mp 51-53°C, ref. 19 mp 50-51°C. Deacetylation gave ethyl 3,4,6-tri-*O*-benzyl-1-thio- $\alpha$ -D-mannopyranoside (**16**).<sup>[18]</sup>

General procedure for preparation of SE glycosides (for amounts of reagents and yields of products, see Table 1). To a solution of the thioglycoside (1 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added 2-(trimethylsilyl) ethanol and molecular sieves (powder, 4 Å, 0.5 g), and the mixture was stirred for 15 min. N-Iodosuccinimide was added, which resulted in slight pink color development. The stirring was continued for another 15 min, when AgOTf (0.1 M in toluene, freshly prepared and stored in the dark) was added dropwise. Red color developed immediately upon the addition, and kept developing intermittently as the AgOTf solution was added, and it finally changed to yellow, indicating that all starting material that had been activated with NIS was consumed. TLC showed complete conversion of the starting material into the faster moving SE glycoside and, occasionally, to more polar product(s). When conversion of the starting material was incomplete, more NIS was added, followed by "titration" with the solution of AgOTf. The reaction mixture was terminated by addition of triethylamine (equimolar to AgOTf). The bright yellow precipitate formed was filtered through a pad of Celite directly into a separating funnel containing excess of 2:1 (v/v) sodium thiosulfate (10%)-sodium bicarbonate (satd) solution. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>, the organic phase was dried, concentrated, and the crude product was chromatographed. 2-(Trimethylsilyl)ethyl glycosides that were prepared following the above protocol are listed below. In all glycosylations of SEOH, except for the conversions  $1 \rightarrow 2$ , and  $34 \rightarrow 35 + 36$  a product more polar than the corresponding 1-thioethyl glycoside, presumably<sup>[5]</sup> the product of hydrolysis of the starting glycosyl donor, was formed.

Ethyl 4-azido-2-*O*-benzoyl-3-*O*-benzyl-4,6-dideoxy-1-thio- $\alpha$ -D-mannopyranoside (1). Benzoyl chloride (0.12 mL, 1.02 mmol) was added to a solution of ethyl 4azido-3-*O*-benzyl-4,6-dideoxy-1-thio- $\alpha$ -D-mannopyranoside<sup>[20]</sup> (220 mg, 0.68 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) containing pyridine (0.17 mL, 2.03 mmol), and the mixture was stirred for 3 h at room temperature. TLC (5:1 hexane-EtOAc) then showed that the reaction was

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

## 464

#### SAKSENA, ZHANG, AND KOVÁČ

Compound	C-1	C-2	C-3	C-4	C-5	C-6
1 <sup>b</sup>	82.31	69.76	76.45	64.55	67.51	18.57
5 <sup>°</sup>	82.54	70.39	71.81	65.79	76.58	62.75
<b>6</b> <sup>d</sup>	84.40	72.00	71.26	67.28	73.22	61.01
<b>7</b> <sup>e</sup>	96.99	69.81	69.13	66.27	68.4	62.63
<b>10</b> <sup>f</sup>	82.35	72.14	70.50	67.07	69.23	62.88
11 <sup>g</sup>	97.10	70.77	70.11	67.08	68.82	63.10
12 <sup>h</sup>	92.28	70.96	69.89	66.83	68.73	62.70
<b>13</b> <sup>i</sup>	82.47	71.06	69.82	65.32	69.31	62.08
14 <sup>j</sup>	97.22	69.75	69.40	65.32	68.86	62.19
<b>16</b> <sup>k</sup>	83.35	69.85	80.42	74.49	71.39	68.83
<b>16</b> <sup>1</sup>	84.24	70.57	81.50	75.60	72.72	70.07
<b>18</b> <sup>m</sup>	82.45	69.91	78.77	74.38	71.68	68.91
<b>18</b> <sup>n</sup>	83.31	70.84	79.89	75.42	73.06	70.02
<b>19</b> °	98.26	69.73	79.30	75.58	72.78	70.25
<b>22</b> <sup>p</sup>	98.18	70.17	79.37	75.44	72.93	70.26
<b>23</b> <sup>q</sup>	97.18	68.31	78.42	74.32	71.24 <sup>r</sup>	69.11
24 <sup>s</sup>	98.56	67.71	80.57	74.30	75.46	69.15
<b>26</b> <sup>t</sup>	90.48	68.70	70.75	65.31	73.02	61.97
$27^{\mathrm{u}}$	90.38	70.75	65.16	67.83	72.88	61.83
<b>28</b> <sup>v</sup>	91.30	69.79	69.91	66.25	71.65	62.52
<b>29</b> <sup>w</sup>	90.57	67.90	69.09	65.24	70.50	61.86
<b>30</b> <sup>x</sup>	82.24	71.64	69.60	66.31	68.90	62.36
<b>31</b> <sup>y</sup>	82.17	70.68	69.77	65.97	68.73	62.16
32 <sup>z</sup>	97.43	70.93	69.93	66.82	69.43	62.86
<b>33</b> <sup>aa</sup>	96.86	69.30	69.42	65.94	68.31	62.36
<b>34</b> <sup>bb</sup>	82.77	77.89	81.64	76.10	73.39	70.34
35 <sup>cc</sup>	98.49	76.49	81.51	76.19	73.40	70.67
<b>36</b> <sup>dd</sup>	101.16	73.91	82.33	74.96	75.88	69.74
<b>37</b> <sup>ee</sup>	82.78	71.12	72.09	65.84	76.37	62.80
<b>38</b> <sup>ff</sup>	82.64	70.15	72.09	65.51	76.07	62.56

*Table 4.* <sup>13</sup>C NMR Chemical Shifts for Compounds 1, 5–7, 10–14, 16, 18, 19, 22–24, and 26–38<sup>a</sup>

<sup>a</sup>For conditions of measurements, see Table 1.

 ${}^{b}\delta_{PhCH_{2}}$  71.44;  $\delta_{SCH_{2}}$  25.67;  $\delta_{SCH_{2}CH_{3}}$  14.92.

 $^{c}J_{C\text{-}1,\text{H}\text{-}1}\text{ 151.4 Hz}; \, \delta_{\text{CH}_2\text{CH}_3} \text{ 14.92}; \, \delta_{\text{CH}_2\text{CH}_3} \text{ 25.78}; \, \delta_{\text{COCH}_3} \text{ 20.65}, \text{ 20.59}, \text{ 20.51}, \text{ 20.47}.$ 

 ${}^{d}\delta_{SCH_2}$  24.94;  $\delta_{SCH_2CH_3}$  14.21.

 ${}^{e}\delta_{CH_{2}CH_{2}Si}$  65.81;  $\delta_{CH_{3}CH_{3}Si}$  17.80;  $\delta_{Si(CH_{3})_{3}}$  – 1.46;  $\delta_{COCH_{3}}$  20.89, 20.70 (3 C).

 ${}^{f}\delta_{SCH_2} 25.61; \delta_{SCH_2CH_3} 14.83.$ 

 ${}^{g}J_{C-1,H-1}$  171.2 Hz;  $\delta_{CH_{2}CH_{2}Si}$  66.16;  $\delta_{CH_{2}CH_{2}Si}$  17.99;  $\delta_{(CH_{3})_{3}Si}$  – 1.40.

<sup>h</sup>J<sub>C-1,H-1</sub> 172.1 Hz.

<sup>i</sup>δ<sub>SCH<sub>2</sub></sub> 25.32;  $δ_{SCH_2CH_3}$  14.75;  $δ_{C(CH_3)_3}$  39.05;  $\delta_{C(CH_3)_3}$  38.88, 38.82, 38.75, 38.68;  $\delta_{C(CH_3)_3}$  27.15, 27.09, 27.05, 27.02.

 ${}^{j}J_{C-1,H-1}$  170.3 Hz;  $\delta_{CH_2CH_2Si}$  65.89;  $\delta_{CH_2CH_2Si}$  17.89;  $\delta_{C(CH_3)_3}$  38.89, 38.86, 38.77, 38.72;  $\delta_{C(CH_3)_3}$  27.14, 27.12, 27.09, 27.06;  $\delta_{Si(CH_3)_3} - 1.38$ .

<sup>k</sup> $\delta_{CH_2CH_3}$  14.80;  $\delta_{CH_2CH_3}$  24.83;  $\delta_{CH_2Ph}$  75.07. 73.38, 72.00.

<sup>1</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{CH_2CH_3}$  15.25;  $\delta_{CH_2CH_3}$  25.27;  $\delta_{CH_2Ph}$  75.50. 73.88, 72.21.

 $^{m}\delta_{PhCH_{2}}$  75.10, 73.14; 71.37;  $\delta_{CH_{2}CH_{3}}$  14.96;  $\delta_{CH_{2}CH_{3}}$  25.45;  $\delta_{C(CH_{3})_{3}}$  39.05;  $\delta_{C(CH_{3})_{3}}$  27.11.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

465

complete and that one product was formed. A little ice, followed by saturated, aqueous NaHCO<sub>3</sub> was added, and the mixture was stirred for 2 h. After extraction with CH<sub>2</sub>Cl<sub>2</sub>, the organic phase was concentrated, and the residue was chromatographed, to give amorphous **1** (285 mg, 98%),  $[\alpha]_D + 65^\circ$  (*c* 1.45); CIMS: *m*/*z* 445 ([M+18]<sup>+</sup>).

Anal. Calcd for  $C_{22}H_{25}N_3O_4S$ : C, 61.81; H, 5.89; N 9.83; S, 7.50. Found: C, 61.98; H, 6.00; N 9.9.76; S, 7.60.

2-(Trimethylsilyl)ethyl 4-azido-2-*O*-benzoyl-3-*O*-benzyl-4,6-dideoxy- $\alpha$ -D-mannopyranoside (2). The conversion  $1 \rightarrow 2$  was a one product reaction. Chromatography gave, in virtually theoretical yield, compound 2, which was identical with the independently synthesized substance (NMR, mp), obtained previously<sup>[6]</sup> in 82% yield.

**Ethyl 1-thio-\alpha-D-mannopyranoside (6).** Conventional deacetylation (Zemplén) of **4** gave the title compound **6**, mp 126–127°C (from EtOH),  $[\alpha]_D + 20.2^\circ$  (*c* 0.85, H<sub>2</sub>O). Anal. Calcd for C<sub>8</sub>H<sub>16</sub>O<sub>5</sub>: C, 42.84; H,7.19. Found: C, 42.75; H, 6.94.

**2-(Trimethylsilyl)ethyl 2,3,4,6-tetra-O-benzoyl-\alpha-D-mannopyranoside (11).** One major and a minor product were formed. Chromatography gave first the amorphous **11**,  $[\alpha]_D - 42.4^\circ$  (c 1.5). CIMS: m/z 714 ( $[M+18]^+$ ).

<sup>r</sup>2-Carbon signal.

<sup>s</sup> $J_{C-1,H-1}$  154.7 Hz;  $\delta_{PhCH_2}$  75.08, 73.23; 71.00;  $\delta_{CH_2CH_2Si}$  67.05;  $\delta_{CH_2CH_2Si}$  17.86;  $\delta_{C(CH_3)_3}$  39.01;  $\delta_{C(CH_3)_3}$  27.22;  $\delta_{Si(CH_3)_3}$  – 1.35.

<sup>t</sup>δ<sub>COCH<sub>2</sub></sub> 20.59 (2 C), 20.54, 20.45.

 $^{u}J_{C-1,H-1}$  162.4 Hz; $\delta_{COCH_3}$  20.56 (2 C), 20.40,  $\delta_{C(CH_3)_3}$  39.15,  $\delta_{C(CH_3)_3}$  26.98.

<sup>v</sup>Measured in C<sub>6</sub>D<sub>6</sub>  $J_{C-1,H-1}$  177.8 Hz;  $\delta_{COCH_3}$  20.63, 20.52 (2 C), 20.30.

<sup>w</sup>δ<sub>COCH<sub>3</sub></sub> 20.85, 20.58 (2 C), 20.54;  $\delta_{C(CH_3)_3}$  39.06;  $\delta_{C(CH_3)_3}$  26.99.

 $^{x}J_{C-1,H-1}$  168.5 Hz; $\delta_{COCH_{3}}$  20.59,  $\delta_{CH_{2}CH_{3}}$  14.74,  $\delta_{CH_{2}CH_{3}}$  25.52.

 $^{y}J_{C-1,H-1}$  168.5 Hz;  $\delta_{COCH_{3}}$  20.60, 20.55, 20.49,  $\delta_{CH_{2}CH_{3}S}$  14.76,  $\delta_{CH_{2}CH_{3}S}$  25.40;  $\delta_{C(CH_{3})_{3}}$  39.02.

<sup>z</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{\text{COCH}_3}$  20.36, 20.23 (2C)  $\delta_{\text{CH}_2\text{CH}_2\text{Si}}$  65.85;  $\delta_{\text{CH}_2\text{CH}_2\text{Si}}$  17.84;  $\delta_{\text{Si}(\text{CH}_3)_3}$  – 1.38.

 ${}^{aa}_{COCH_3} 20.60, 20.55, 20.51; \delta_{CH_2CH_2Si} 65.72; \delta_{CH_2CH_2Si} 17.73; \delta_{C(CH_3)_3} 38.95; \delta_{Si(CH_3)_3} - 1.52.$ 

<sup>bb</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{CH_2CH_3}$  15.52,  $\delta_{CH_2CH_3}$  25.78,  $\delta_{PhCH_2}$  75.55, 73.82, 72.87; 72.49.

<sup>cc</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $J_{C-1,H-1}$  166.0 Hz;  $\delta_{PhCH_2}$  75.57, 73.90, 73.52, 72.62;  $\delta_{CH_2CH_2Si}$  65.34;  $\delta_{CH_2CH_2Si}$  18.45;  $\delta_{Si(CH_3)_3} = 0.93$ .

 $^{dd}J_{C-1,H-1} 153.5 \text{ Hz}; \ \, \delta_{PhCH_2} 75.12, 73.78, 73.43, 71.31; \ \, \delta_{CH_2CH_2Si} 67.19; \ \, \delta_{CH_2CH_2Si} 18.28, \\ \delta_{Si(CH_3)_3} - 1.39.$ 

 $^{ee}J_{C-1,H-1}$  153.9 Hz;  $\delta_{COCH_3}$  20.72, 20.66, 20.60;  $\delta_{CH_2,CH_3}$  14.89,  $\delta_{CH_2,CH_3}$  25.91.

<sup>ff</sup>*J*<sub>C-1,H-1</sub> 154.1 Hz;  $δ_{C(CH_3)_3}$  39.10;  $δ_{C(CH_3)_3}$  27.06;  $δ_{CH_2CH_3}$  14.90;  $δ_{CH_2CH_3}$  25.95;  $δ_{COCH_3}$  20.67, 20.64, 20.50.

Notes to Table 4 Continued:

<sup>&</sup>lt;sup>n</sup>Measured in C<sub>6</sub>D<sub>6</sub>.  $\delta_{PhCH_2}$  75.64, 73.68; 72.00;  $\delta_{CH_2CH_3}$  15.30;  $\delta_{CH_2CH_3}$  25.97;  $\delta_{C(CH_3)_3}$  39.39;  $\delta_{C(CH_3)_3}$  27.69.

<sup>&</sup>lt;sup>o</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  75.65, 73.91, 72.27;  $\delta_{CH_2CH_2Si}$  65.64;  $\delta_{CH_2CH_2Si}$  18.37;  $\delta_{COCH_3}$  20.99;  $\delta_{Si(CH_1)_2} = 0.99$ .

<sup>&</sup>lt;sup>p</sup>Measured in C<sub>6</sub>D<sub>6</sub>;  $\delta_{PhCH_2}$  75.68, 73.93, 72.08;  $\delta_{CH_2CH_2Si}$  65.72;  $\delta_{CH_2CH_2Si}$ , 18.35;  $\delta_{Si(CH_3)_3} - 0.96$ . <sup>q</sup> $\delta_{PhCH_2}$  75.11, 73.16; 71.24;  $\delta_{CH_2CH_2Si}$  65.11;  $\delta_{CH_2CH_2Si}$  17.84;  $\delta_{C(CH_3)_3}$  38.94;  $\delta_{C(CH_3)_3}$  27.12;  $\delta_{Si(CH_3)_3} - 1.41$ .

466

#### SAKSENA, ZHANG, AND KOVÁČ

Anal. Calcd for  $C_{39}H_{40}O_{10}Si$ : C, 67.72; H, 5.79. Found: C, 66.86; H, 5.86. Eluted next was 2,3,4,6-tetra-*O*-benzoyl- $\alpha$ , $\beta$ -D-mannopyranose (**12**), which was identified by comparison with an authentic sample,<sup>[21]</sup> (TLC, NMR); CIMS: *m/z* 614 ([M+18]<sup>+</sup>).

Ethyl 2,3,4,6-tetra-*O*-pivaloyl-1-thio-α-D-mannopyranoside (13). Pivaloylation of **6**, as described below for preparation of **18**, gave the title ethyl 1-thioglycoside **13** in 92% yield, mp 104–105°C (from EtOH),  $[\alpha]_D + 58°$  (*c* 0.9); CIMS: *m/z* 578 ([M+18]<sup>+</sup>). Anal. Calcd for C<sub>28</sub>H<sub>48</sub>O<sub>9</sub>S: C, 59.97; H, 8.63. Found: C, 60.12; H, 8.65.

**2-(Trimethylsilyl)ethyl 2,3,4,6-tetra-O-pivaloyl-\alpha-D-mannopyranoside (14).** One major product, faster moving than the starting material, and one minor, slower moving product was formed, as shown by TLC (30:1 toluene–acetone). Chromatography gave 14, mp 82–84°C (from EtOH containing a few drops of water,  $[\alpha]_D + 33.5^{\circ}$  (*c* 0.9); FAB MS: *m/z* 455 ([M+Na]<sup>+</sup>).

Anal. Calcd for C<sub>31</sub>H<sub>56</sub>O<sub>10</sub>Si: C, 60.36; H, 9.15. Found: C, 60.32; H, 9.14.

Ethyl 3,4,6-tri-*O*-benzyl-2-*O*-pivaloyl-1-thio-α-D-mannopyranoside (18). Pivaaloyl chloride (1.2 mL, 10 mmol) was added with stirring to a solution of ethyl 3,4,6tri-*O*-benzyl-1-thio-α-D-mannopyranoside<sup>[18]</sup> (16, 2.36 g, 5 mmol) in pyridine (1.2 mL, 15 mmol), and the mixture was stirred at room temperature with exclusion of atmospheric moisture overnight. TLC (15:1 hexane–acetone) showed that the reaction was complete and that one product was formed. Aqueous NaHCO<sub>3</sub> solution was added, to destroy excess of the reagent, and the mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and aqueous NaHCO<sub>3</sub> solution. The organic phase was dried, concentrated, and chromatography gave amorphous compound 18 (3.10 g, 87%),  $[\alpha]_D + 71^\circ$  (*c* 0.9); CIMS: *m/z* 596 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>34</sub>H<sub>42</sub>O<sub>6</sub>S: C, 70.56; H, 7.31. Found: C, 70.77; H,7.41.

2-(Trimethylsilyl)ethyl 2-O-acetyl-3,4,6-tri-O-benzyl- $\alpha$ -D-mannopyranoside (19). One major and three minor products were formed. Eluted first was the title compound 19,  $[\alpha]_D + 29^\circ$  (c 2.8), CIMS: m/z 591 ([M-1]).

Anal. Calcd for C<sub>33</sub>H<sub>44</sub>O<sub>7</sub>Si: C, 68.89; H, 7.48. Found: C, 68.98; H, 7.51.

Two of the three minor products formed were isolated in amounts sufficient to be identified by MS and NMR spectroscopy. Eluted next was the product of deacetylation of the major product **19**, 2-(trimethylsilyl)ethyl 3,4,6-tri-*O*-benzyl- $\alpha$ -D-mannopyranoside (**20**, ~ 20 mg). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  4.91 (d, 1 H,  $J_{1,2}$  1.7 Hz, H-1); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  98.68 ( $J_{C-1,H-1}$  168.6 Hz, C-1), 80.27 (C-3), 75.10, 73.40, 71.91 (3 CH<sub>2</sub>Ph), 73.37 (C-4), 70.93 (C-5), 68.96 (C-6), 68.51 (C-2), 64.88 (OCH<sub>2</sub>CH<sub>2</sub>Si), 17.85 (CH<sub>2</sub>Si), -1.37[(CH<sub>3</sub>)<sub>3</sub>]; FAB MS: m/z 573 ([M+Na]<sup>+</sup>), which was not further characterized. When this material was acetylated, the product was indistinguishable from compound **19** described above (TLC, NMR, MS).

Eluted last was the hemiacetal **21**, as indicated by its chromatographic mobility and m/z value 515 ( $[M+Na]^+$ ) in the FAB MS spectrum.

**2-(Trimethylsilyl)ethyl 2-O-benzoyl-3,4,6-tri-O-benzyl-α-D-mannopyranoside** (22). One major product, faster moving that the starting material, and one minor,

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

467

slower moving product was formed, as shown by TLC (40:1 toluene–acetone). Chromatography gave the amorphous **22**,  $[\alpha]_D - 2.3^\circ$  (*c* 3.4); CIMS: *m/z* 672 ( $[M+18]^+$ ). Anal. Calcd for C<sub>39</sub>H<sub>46</sub>O<sub>7</sub>Si: C, 71.53; H, 7.08. Found: C, 71.55; H, 7.17.

**2-(Trimethylsilyl)ethyl 3,4,6-tri-O-benzyl-2-O-pivaloyl-\alpha-D-mannopyranoside** (23). Three products were formed, the one showing the fastest chromatographic mobility largely predominating. Chromatography (25:1 toluene–EtOAc) gave first the title glycoside **23**,  $[\alpha]_{\rm D}$ +29° (*c* 0.9); CIMS: *m/z* 652 ([M+18]<sup>+</sup>).

Anal. Calcd for C37H50O7Si: C, 70.00; H, 7.94. Found: C, 70.06; H, 7.82.

Eluted next was a small amount of material whose NMR spectra showed it to be 2-(trimethylsilyl)ethyl 3,4,6-tri-O-benzyl-2-O-pivaloyl- $\beta$ -D-mannopyranoside (24).

**1,3,4,6-Tetra-***O***-acetyl-***2***-***O***-benzoyl-***β***-***D***-mannopyranose (26).** Benzoylation of **25**,<sup>[17]</sup> as described above for the preparation of **1** gave **26** in virtually theoretical yield, mp 103–104°C (from EtOH),  $[\alpha]_D - 65^\circ$  (*c* 0.8); CIMS: *m/z* 470 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>21</sub>H<sub>24</sub>O<sub>11</sub>: C, 55.75; H, 5.35 Found: C, 55.68; H, 5.35.

**1,3,4,6-Tetra-***O***-acetyl-***2***-***O***-pivaloyl-***β***-***D***-mannopyranose** (27). A reaction of the 2-hydroxy compound 25 with pivaloyl chloride, as described above for similar benzoylation gave, after chromatography, the title derivative 27 in virtually theoretical yield, mp 106–107°C (from EtOH),  $[\alpha]_D - 20.4^\circ$  (*c* 0.8); CIMS: *m/z* 455 ( $[M+18]^+$ ).

Anal. Calcd for C<sub>19</sub>H<sub>28</sub>O<sub>11</sub>: C, 52.77; H, 6.53. Found: C, 52.91; H, 6.47.

**1,3,4,6-Tetra-***O***-acetyl-***2***-***O***-benzoyl-***α***-***D***-mannopyranose** (**28**). A solution of **26** (1 g) in 10:4:0.1 Ac<sub>2</sub>O:AcOH:H<sub>2</sub>SO<sub>4</sub> (6 mL) was kept at room temperature overnight. TLC (3:1 hexane–acetone) showed that a faster moving product was formed. <sup>1</sup>H NMR showed that a ~93:7 equilibrium mixture of α- and β-anomers was present.<sup>b</sup> Solid NaOAc trihydrate was added with stirring, to neutralize the mineral acid, followed by aqueous NaHCO<sub>3</sub>, to destroy Ac<sub>2</sub>O and neutralize AcOH. The resulting mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and aqueous NaHCO<sub>3</sub> solution, the organic phase was dried, concentrated, and chromatography gave first the pure α-anomer **28** (0.83 g, 83%), mp 102–103°C (from EtOH),  $[\alpha]_D - 15°$  (*c* 0.8); FAB MS: *m/z* 475 ([M+Na]<sup>+</sup>).

Anal. Calcd for C<sub>21</sub>H<sub>24</sub>O<sub>11</sub>: C, 55.75; H, 5.35 Found: C, 55.72; H, 5.35.

Eluted next was a mixture of  $\alpha$ - and  $\beta$ -anomers. Such mixtures, obtained from several experiments, could be processed as described above, to obtain more of the desired **28**.

**1,3,4,6-Tetra-O-acetyl-2-O-pivaloyl-\alpha-D-mannopyranose (29).** Anomerization of **27**, as described above for the preparation of **26** gave, after chromatography, the amorphous **29** (90%),  $[\alpha]_D + 42^\circ$  (*c* 0.8); CIMS: *m/z* 455 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>19</sub>H<sub>28</sub>O<sub>11</sub>: C, 52.77; H, 6.53. Found: C, 52.79; H, 6.65.

<sup>&</sup>lt;sup>b</sup>That ratio had not changed when, in a separate experiment, the reaction time was extended to 48 h, as showed by <sup>1</sup>H NMR spectroscopy. The two anomers show almost identical mobility on TLC.

468

#### SAKSENA, ZHANG, AND KOVÁČ

Ethyl 3,4,6-tri-*O*-acetyl-2-*O*-benzoyl-1-thio-α-D-mannopyranoside (30). *A*. From 1,3,4,6-Tetra-*O*-acetyl-2-*O*-benzoyl-β-D-mannopyranose (26). Ethanethiol (0.45 mL, 3 mmol) followed by BF<sub>3</sub>-etherate (1.5 mL) was added at 0°C, with stirring and exclusion of moisture, to a solution of 26 (1.35 g, 3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (60 mL), and the mixture was kept in a refrigerator (5–7°C) until TLC (3:1 hexane–acetone) showed that almost all starting material was consumed (~3 days). One major and several minor products were formed. The mixture was slowly poured into aqueous NaHCO<sub>3</sub>, contained in a separatory funnel, the product was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and chromatography gave first the α-anomer 30 (0.785 g, 58%), mp 66–67°C (from EtOH),  $[\alpha]_D+127^\circ$  (*c* 1.7); CIMS: m/z 472 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>21</sub>H<sub>26</sub>O<sub>9</sub>S: C, 55.50; H, 5.77. Found: C, 55.62; H, 5.90.

Eluted next was a small amount of material whose NMR spectra showed it to be ethyl 3,4,6-tri-*O*-acetyl-2-*O*-benzoyl-1-thio- $\beta$ -D-mannopyranoside (**37**, 45 mg, 3%); CIMS: m/z 472 ([M+18]<sup>+</sup>).

*B*. From 1,3,4,6-tetra-*O*-acetyl-2-*O*-benzoyl- $\alpha$ -D-mannopyranose (**28**). The reaction of **28**, performed on the same scale as described above but with only 1.2 mL of BF<sub>3</sub>-etherate, was terminated after ~ 24 h. Chromatography gave **30** and **37** in 65 and 2% yield, respectively.

Ethyl 3,4,6-tri-*O*-acetyl-2-*O*-pivaloyl-1-thio-α-D-mannopyranoside (31). *A*. From 1,3,4,6-Tetra-*O*-acetyl-2-*O*-pivaloyl-β-D-mannopyranose (27). The reaction, carried out as described above for the preparation of 30 (*B*) gave, after a reaction time of 48 h and chromatography, the α-anomer 31 (680 mg, 52%), 93.5–94°C,  $[\alpha]_D$ +90° (*c* 0.8). CIMS: *m/z* 452 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>19</sub>H<sub>30</sub>O<sub>9</sub>S: C, 52.52; H, 6.96. Found: C, 52.53; H, 7.03.

Eluted next was the amorphous ethyl 3,4,6-tri-*O*-acetyl-2-*O*-pivaloyl-1-thio-β-D-mannopyranoside (**38**, 300 mg, 23%),  $[\alpha]_D - 76^\circ$  (c 1.6); CIMS: *m/z* 452 ([M+18]<sup>+</sup>). Found: C, 53.02; H, 6.77.

*B*. From 1,3,4,6-tetra-*O*-acetyl-2-*O*-pivaloyl- $\alpha$ -D-mannopyranose (**29**). Reaction as above (*A*) gave compounds **31** and **38** in 60 and 12%, respectively.

**2-(Trimethylsilyl)ethyl** 3,4,6-tri-*O*-acetyl-2-*O*-benzoyl- $\alpha$ -D-mannopyranoside (32). One major and one minor product were formed. Chromatography (5:1 he-xane-acetone) gave 32,  $[\alpha]_D$ +15.5° (*c* 0.5); CIMS: *m/z* 528 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>24</sub>H<sub>34</sub>O<sub>10</sub>Si: C, 56.45; H, 6.71. Found: C, 56.51; H, 6.61.

**2-(Trimethylsilyl)ethyl** 3,4,6-tri-*O*-acetyl-2-*O*-pivaloyl- $\alpha$ -D-mannopyranoside (33). One major and one minor product were formed. Chromatography (10:1 toluene–acetone) gave 33 [ $\alpha$ ]<sub>D</sub>+53° (*c* 1.1); CIMS: *m/z* 508 ([M+18]<sup>+</sup>).

Anal. Calcd for C<sub>22</sub>H<sub>38</sub>O<sub>10</sub>Si: C, 53.86; H, 7.81. Found: C, 53.77; H, 7.80.

2-(Trimethylsilyl)ethyl 2,3,4,6-tetra-*O*-benzyl- $\alpha$ - (35) and  $\beta$ -D-mannopyranoside (36). Two products, which showed similar chromatographic mobility were formed, as shown by TLC (10:1 hexane–EtOAc). The  $\alpha$ -anomer 35, eluted first, showed  $[\alpha]_D + 48^\circ$  (*c* 0.7). FABMS: *m*/*z* 663 ([M+Na]<sup>+</sup>).

Anal. Calcd for C<sub>39</sub>H<sub>48</sub>O<sub>6</sub>Si: C, 73.09; H,7.55. Found: C, 73.03; H,7.70.

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

#### 2-(TRIMETHYLSILYL)ETHYL α-D-MANNOPYRANOSIDES

469

The  $\beta$ -anomer **36**, which was eluted next, crystallized on standing. Recrystallization from EtOH gave material melting at 56–59°C,  $[\alpha]_D - 13.5^\circ$  (*c* 0.3), FABMS: *m/z* 663 ([M+Na]<sup>+</sup>). Found: C, 73.16; H, 7.70.

#### REFERENCES

- Jansson, K.; Ahlfors, S.; Frejd, T.; Kihlberg, J.; Magnusson, G.; Dahmen, J.; Noori, G.; Stenwall, K. 2-(Trimethylsilyl)ethyl glycosides. Synthesis, anomeric deblocking, and transformation into 1,2-trans 1-O-acyl sugars. J. Org. Chem. **1988**, *53*, 5629–5647.
- 2. Jansson, K.; Magnusson, G. 2-(Trimethylsilylethyl) glycosides. Treatment with electrophilic reagents to give trimethylsilyl and methoxymethyl glucopyranosides and glucopyranosyl chlorides. Tetrahedron **1990**, *46*, 59–64.
- Jansson, K.; Noori, G.; Magnusson, G. 2-(Trimethylsilyl)ethyl glycosides. Transformation into glycopyranosyl chlorides. J. Org. Chem. 1990, 55, 3181–3185.
- Kartha, K.P.R.; Jennings, H.J. A facile, one-step procedure for the conversion of 2-(trimethylsilyl)ethyl glycosides to their glycosyl chlorides. Tetrahedron Lett. 1990, 31, 2537–2540.
- 5. Pozsgay, V. A synthesis of 2-(trimethylsilyl)ethyl α-D-mannopyranoside. Tetrahedron Lett. **1993**, *34*, 7175–7178.
- Ogawa, Y.; Lei, P.-S.; Kováč, P. Synthesis of ligands related to the *Vibrio cholerae* O-specific antigen. Part 11. Synthesis of four glycosides of a disaccharide fragment representing the terminus of the O-polysaccharide of *Vibrio cholerae* O:1, serotype Inaba, bearing aglycons suitable for linking to proteins. Carbohydr. Res. **1996**, 288, 85–98.
- Ogawa, Y.; Lei, P.-S.; Kováč, P. Synthesis of ligands related to the *Vibrio cholerae* O-specific antigen. Part 12. Synthesis of eight glycosides of hexasaccharide fragments representing the terminus of the O-polysaccharide of *Vibrio cholerae* O:1, serotype Inaba and Ogawa, bearing aglycons suitable for linking to proteins. Carbohydr. Res. **1996**, *293*, 173–194.
- Kanie, O.; Ito, Y.; Ogawa, T. Orthogonal glycosylation strategy in oligosaccharide synthesis. J. Am. Chem. Soc. 1994, 116, 12073–12074.
- Konradson, P.; Udodong, U.E.; Fraser-Reid, B. Iodonium promoted reactions of disarmed thioglycosides. Tetrahedron Lett. 1990, 31, 4313–4316.
- Norberg, T. Glycosylation Properties and Reactivity of Thioglycosides, Sulfoxides and Other S-glycosides: Current Scope and Future Prospects. In *Modern Methods in Carbohydrate Synthesis*; Khan, S.H., O'Neill, R.A., Eds.; Harwood Academic Publishers: Amsterdam, 1996; 82–106.
- Sarbajna, S.; Roy, N. Synthesis of tri- and tetrasaccharide derivatives related to *Klebsiella* type 57. Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem. 1998, 37B (3), 252–256.
- 12. Fried, J.; Walz, D.E. Ethyl thioglycosides of D-mannose and D-galactose and a new synthesis of styracitol. J. Am. Chem. Soc. **1949**, *71*, 140–143.
- Kunz, H.; Harreus, A. Synthesis of glycosides using 2,3,4,6-tetra-O-pivaloyl-α-Dglucopyranosyl bromide. Liebigs Ann. Chem. 1982, 41–48.

470

#### SAKSENA, ZHANG, AND KOVÁČ

- Bock, K.; Pedersen, C. A study of 13CH coupling constants in hexopyranoses. J. Chem. Soc., Perkin Trans. 2 1974, 293–297.
- Das, S.K.; Roy, N. An improved method for the preparation of some ethyl 1thioglycosides. Carbohydr. Res. 1996, 296, 275–277.
- Helander, A.; Kenne, L.; Oscarson, S.; Peters, T.; Brisson, J.-R. Synthesis and conformational and NMR studies of α-D-mannopyranosyl- and α-D-mannopyranosyl-(1-2)-α-D-mannopyranosyl linked to L-serine and L-threonine. Carbohydr. Res. 1992, 230, 299–318.
- Bovin, N.V.; Zurabyan, S.E.; Khorlin, A.Y. On the nucleophilic displacement at C-2 of hexopyranoses. Izv. Akad. Nauk, Ser. Khim. 1981, 1638–1641.
- Zhang, Y.-M.; Mallet, J.-M.; Sinaÿ, P. Glycosylation using a one-electron-transfer, homogeneous reagent. Application to an efficient synthesis of the trimannosyl core of *N*-glycosylproteins. Carbohydr. Res. **1992**, *236*, 73–88.
- Barresi, F.; Hindsgaul, O. The synthesis of β-mannopyranosides by intramolecular aglycon delivery: Scope and limitation of the existing methodology. Can. J. Chem. 1994, 72, 1447–1465.
- Peters, T.; Bundle, D.R. Synthetic antigenic determinants of the *Brucella* A polysaccharide: A disaccharide thioglycoside for block synthesis of pentasaccharide and lower homologues of a 1,2-linked 4,6-dideoxy-4-formamido-α-D-mannose. Can. J. Chem. **1989**, 67, 491–496.
- Zhang, J.; Kováč, P. An alternative method for regioselective, anomeric deacylation. J. Carbohydr. Chem. 1999, 18, 461–469.

Received March 14, 2002 Accepted June 27, 2002