

Accepted Manuscript

Multicomponent Diversity Oriented Synthesis of Multivalent Glycomimetics Containing Hexafluorovaline

M.C. Bellucci, A. Sganappa, M. Sani, A. Volonterio



PII: S0040-4020(15)01147-3

DOI: [10.1016/j.tet.2015.07.070](https://doi.org/10.1016/j.tet.2015.07.070)

Reference: TET 27015

To appear in: *Tetrahedron*

Received Date: 10 June 2015

Revised Date: 14 July 2015

Accepted Date: 27 July 2015

Please cite this article as: Bellucci MC, Sganappa A, Sani M, Volonterio A, Multicomponent Diversity Oriented Synthesis of Multivalent Glycomimetics Containing Hexafluorovaline, *Tetrahedron* (2015), doi: 10.1016/j.tet.2015.07.070.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Multicomponent Diversity Oriented Synthesis of Multivalent Glycomimetics Containing Hexafluorovaline

M.C. Bellucci,^a A. Sganappa,^b M. Sani,^c and A. Volonterio^{b,c,*}

^a *Department of Food, Environmental and Nutritional Sciences, Università degli Studi di Milano, via
Celoria 2, 20133 Milano, Italy.*

^b *Department of Chemistry, Materials, and Chemical Engineer “G. Natta”, Politecnico di Milano, via
Mancinelli 7, 20131 Milano, Italy*

^c *C.N.R. Istituto del Riconoscimento Molecolare, via Mancinelli 7, 20131 Milano, Italy.*

alessandro.volonterio@polimi.it

Abstract

The chemoselective introduction of fluorinated moieties into biologically relevant scaffolds is now an established strategy to modulate and to study the properties of molecular leads. In this article we propose, for the first time, the diversity oriented synthesis of multivalent glycomimetics incorporating hexafluorovaline through a straightforward multicomponent sequential process which occurs with high yield and in mild conditions. The same process has been successfully applied to the chemistry of

aminoglycosides producing a neomycin-hexafluorovaline-galactose conjugate and providing a general, efficient strategy to functionalized aminoglycosides with sugar-hexafluorovaline tags.

Keywords: Diversity Oriented Synthesis, Multicomponent Synthesis, Multivalent Glycomimetic, Fluorine, Aminoglycoside.


1. Introduction

The introduction of a fluorine atom or perfluorinated groups into chemical scaffolds is now a standard strategy for modulating the chemical and biophysical properties of molecular leads and for studying the biological aspects associated with the fluorine substitution/introduction, such as metabolism, excretion properties, and fluorine-containing ligand binding interactions.¹ This is due to the unique properties of the fluorine atom, its strongest inductive effect compared to other elements, its small size, and low polarizability. For instance, the highly fluorinated amino acids hexafluorovaline (hfVal) and hexafluoroleucine (hfLeu), being often well tolerated, have been used to modulate such properties of peptides and proteins like their hydrophobicity, acidity/basicity, folding and stability.² Moreover, ¹⁹F is the most stable and abundant fluorine isotope, thus fluorinated peptides and proteins can be easily tracked with NMR and/or MRI spectroscopy, even in living cells.³ Thus, since fluorine containing organic molecules are almost absent in nature,⁴ there is great interest in developing new synthetic strategies to introduce selectively fluorinated moieties into chemical scaffolds.

Multivalent glycomimetics are a class of very intriguing compounds.⁵ Indeed, there is a growing interest in glycosidase inhibition, and in elucidating fundamental biological pathways through the synthesis of glycomimetics. Multivalency can increase dramatically the binding potency of the corresponding monovalent ligands as already demonstrated.⁶ However, despite such interest, compared to monovalent glycomimetics, there are relatively few example in the literature of multivalent glycosidase inhibitors, and no fluorinated multivalent glycosidase inhibitors, probably

because they are not synthetically accessible in an easy way.⁶ One strategy to overcome such a deficiency could be the development of straightforward methodologies for the diversity oriented synthesis (DOS) of these compounds.⁷ Indeed, DOS, often combined with multicomponent reactions (MCRs), is an indispensable tool for modern pharmaceutical and drug discovery research programs because they offer the possibility to access molecular complexity with a minimum of effort, in addition to atom economy, operational simplicity, and bond-forming efficiency.⁸ This is well evidenced by the growing application of such strategies for the synthesis of libraries of different collections of molecules. However, to the best of our knowledge, there are no examples of DOS of multivalent, selectively fluorinated glycomimetics.⁹

Very recently, in the frame of a research program dealing with the discovery of novel MCRs for the synthesis of heterocycles¹⁰ and glycoconjugates,¹¹ we have reported for the first time the MC combinatorial synthesis of diversity oriented multifunctional glycomimetics **1-4**, where the sugar moieties were tethered through artificial linkers such as aspartic acid, a hydantoin ring, or urea frameworks (Figure 1).¹² Herein we wish to report an extension of such process, using commercially available 4,4,4-trifluoro-3-trifluoromethyl(Tfm)-crotonic acid in order to prepare multifunctional glycomimetics where the Asp moiety of **1-4** is substituted by hfVal.



previous work $R^1 = H$, $R^2 = COOEt$ or $COOBn$

this work $R^1 = R^2 = CF_3$


Figure 1. Multivalent glycomimetics containing hfVal

We propose that the introduction of hfVal in the multivalent glycomimetic scaffolds can impart interesting new properties to this intriguing class of compounds such as, for instance, higher metabolic

stability and lipophilicity and would facilitate the investigation of target-ligand interactions through ^{19}F NMR spectroscopy.

2. Results and Discussion

4,4,4-Trifluoro-3-Tfm-crotonic acid is a commercially available fluorinated building block that, due to the high electronegativity of the trifluoromethyl groups, has been used for the synthesis of hfVal amino acid derivatives through *anti*-Michael addition of *N*-nucleophiles.¹³ Recently, our group has exploited the reactivity of 4,4,4-trifluoro-3-Tfm-crotonic acid **8** for the synthesis of glycoconjugates containing hfVal **10** and **11** through the MC sequential domino process depicted in Scheme 1.^{10,11} The reaction of **8** with *in situ* generated *N*-glycosyl, *N'*-*tert*-butyl carbodiimides **7** produces intermediate **A** which undergoes intramolecular *anti*-aza-Michael reaction affording a second highly reactive intermediate **B**. The latter step is totally regioselective when the steric hindrance of the two *N*-substituents in the carbodiimide framework is very different, namely a primary versus a tertiary alkyl group. In the absence of a *N*-nucleophile, **9** intermediate **B** rearranges through an O→N acyl migration process affording hydantoin scaffold **10** containing a hfVal moiety (path A, Scheme 1).¹⁴ Instead, in the presence of *N*-nucleophile **9**, such as amines, amino esters, or peptides, intermediate **B** undergoes nucleophilic attack leading to the formation of urea-hfVal-glycopeptides **11** (path B, Scheme 1).




Scheme 1. Mechanism of the domino MC process

Since this process is high yielding, totally regioselective and operationally very simple (MC domino process which does not require high temperatures, dry solvents, difficult purification steps), we wished to exploit it for the synthesis of multivalent glycomimetics containing hfVal by









Disappointingly, even in the presence of highly polar DMF the yields were raised only to 51% (Scheme 2).

Once proved that the reaction works efficiently even with *O*-protected *N*-glycosyl amines as nucleophiles in the standard conditions, we applied the process for the four-component sequential synthesis of a library of divalent glycoconjugates containing hfVal **1** starting from different *N*-*tert*-butyl, *N'*-glycosyl carbodiimides **7**, which are formed *in situ* by Staudinger reaction between sugar azides **5** and *tert*-butyl isocyanate **6**, and *N*-glycosyl nucleophiles **9** (Table 1).

Table 1. MC combinatorial synthesis of multivalent glycomimetics **1**.



Entry	Sugar-N ₃	Carbodiimide	Sugar-NH ₂	Product	Yield (%) ^a
1					81 ^b
2					53 ^{b,d}
3					78 ^c
4					68 ^b

5					83 ^b
6					76 ^b


^aIsolated yields. ^bAn almost 3:1 diastereoisomeric mixture. ^cAn almost 1.5:1 diastereoisomeric mixture. ^dA 30% of the corresponding hydantoin was recovered.

Indeed, the reaction between glycosyl azides **5a-e** and commercially available *tert*-butyl isocyanate **6** lead to the formation of the corresponding carbodiimides **7a-e**, respectively, upon treatment with Ph_3P in CH_3CN at rt (Staudinger reaction). The resulting carbodiimides **7a-e** can also be isolated by quick flash chromatography and characterized. However, compounds **7a-e** can be used *in situ* in order to achieve a four component sequential process. In this case, once the carbodiimide is formed (TLC monitoring), the temperature is cooled to 0°C and 2,4,6-trimethylpyridine (TMP) followed by nucleophile **9** and, finally, acid **8** are added. The process was very general, working efficiently either starting from sugar azides (where the functional group is attached to primary carbon 6 in hexoses or carbon 5 in pentoses) leading to the formation of multivalent glycomimetics having enzymatically stable “ CH_2NH_2 ” backbone mimicking glycine (entries 1-4, Table 1),¹⁶ or attached to the anomeric carbon through a suitable linker (entries 5 and 6, Table 1). In all cases the reaction was completely regioselective leading to the formation of regioisomers **1a-f** as a mixture of diastereoisomers, which arose from the nucleophilic attack of the less sterically congested primary *N*-glycosyl moiety in the intramolecular *anti*-aza-Michael step (Scheme 1, path B).¹⁷ The yields were generally high except when glycosylamine **9c** was used as nucleophile (entry 2, Table 1). This result was due to the poor nucleophilicity of the hemiacetalic amino group and scarce solubility of **9c** as already stated in previous works.^{11d} In this way we were able to obtain a small library of six divalent glycomimetics where the sugar moieties are connected through a urea-hfVal linker. The reaction is very general, and any kind of


























appropriately functionalized carbohydrates could be used, such as ribose, galactose, glucose, glucosamine, and mannose.

In order to obtain other multivalent glycomimetics containing hfVal, where the sugar moieties are connected with different scaffolds (DOS), we tested the same reaction starting with carbodiimides having two *N*-sugar substituents. In this way, considering the synthetic pathway depicted in Scheme 1, we would be able to obtain three new multivalent glycomimetics, *i.e.* divalent glycomimetics **2**, **3** where the carbohydrates are connected through a hydantoin scaffold or a urea linker, respectively, and trivalent glycomimetic **4** where the sugar moieties are tethered through a urea-hfVal scaffold. *N,N'*-Disugar carbodiimides **16** could be prepared starting from the corresponding sugar azides **5** and sugar isothiocyanates **15** (Table 2).

Table 2. MC combinatorial synthesis of multivalent glycomimetics **2-4**.



Entry	Sugar- N_3	Sugar-NCS	Carbodiimide	Sugar- NH_2	Product	Yield (%) ^a
1				/		78 ^b
2				/		82 ^c

3						75 ^c
4						72 ^b
5						76 ^b
6						62 ^b
7						64 ^c


^aIsolated yields. ^bAn almost 1.5:1 diastereoisomeric mixture. ^cAn almost 3:1 diastereoisomeric mixture.

Accordingly, Staudinger reaction between ribose azide **5a** and isothiocyanate **15a** lead to the clean formation of carbodiimide **16a** along with Ph₃PS (TLC monitoring). By adding *in situ* TMP followed by acid **8**, we obtained the formation of divalent glycomimetic **2a**, as an almost 1.5:1 mixture of diastereoisomers, where the sugar moieties are connected through a hydantoin ring (entry 1, Table 2). Under the same conditions we obtained the formation of diglycohydantoin **2b** in very high yields starting with galactose azide **5b** and isothiocyanate **15b** (entry 2, Table 2). Interestingly, the reaction carried out in the presence of α -amino esters nucleophiles such as H-Ala-OBn **17a** and H-Leu-OBn **17b** furnished divalent glycopeptides **3a,b**, respectively (entry 3

and 4, Table 2). The latter reaction is particularly promising because it permits the unprecedented multicomponent synthesis of a class of very intriguing scaffolds, *i.e.* divalent glycopeptides incorporating the unnatural amino acid hfVal. Moreover, the process works very efficiently also starting from dipeptide nucleophiles such as H-Phe-Leu-OBn **17c** leading to the one pot formation of divalent glycotriptides containing hfVal **3c** in high yields (entry 5, Table 2). Finally, in order to demonstrate that this process could also be exploited for the preparation of more complicated trivalent glycomimetics, we tested the reaction starting with *in situ* generated symmetric *N,N'*-diglyco carbodiimide **16a,b** and sugar nucleophile **9a,d**, respectively. We were delighted that in this case the reaction also works effectively giving rise to the formation of trivalent glycomimetics **4a,b** in good yields (entries 6 and 7, Table 2). It is worth noting that, in order to increase complexity, the synthesis of glycomimetics **2-4** could be also achieved starting with asymmetric *N,N'*-diglyco carbodiimides. However, even if high yielding, due to the similarity on the steric hindrance of the *N*-substituents, the process lead to the formation of mixtures of regioisomers that are difficult to separate (data not shown).

Beside the synthesis of multivalent glycomimetics **1-4**, this process could be exploited as a general procedure for the introduction of sugar-hfVal tags into biologically relevant molecules in a simple and convenient way. With this in mind, we sought to functionalize aminoglycoside antibiotics. Aminoglycosides are a class of polyaminosugars which selectively bind RNA.¹⁸ The selective functionalization of such scaffolds has been exploited for the synthesis of conjugates with better characteristics, such as, for instance, activity, selectivity, and ability to fight against aminoglycosides resistance.¹⁹ In this sense, the introduction of fluorinated probes in the aminoglycoside scaffolds could facilitate the investigation of RNA-ligand interaction through ¹⁹F NMR spectroscopy.²⁰ However, besides the interest in the development of new practical ways to selectively functionalize aminoglycosides, there are only a few examples that use a multicomponent one-pot procedure, and, to the best of our knowledge, none of them are used for tethering aminoglycosides with fluorinated tags.²¹ Thus, the successful application of our process to aminoglycoside chemistry would provide a general

way to selectively functionalize such molecules with other scaffolds (in this case carbohydrates) along with a fluorinated tag (hfVal) with a practical one pot multicomponent procedure. Accordingly, we decided to use neomycin derivative **18** as nucleophile in our multicomponent process. To our delight, reacting carbodiimide **6b** in the presence of acid **8** and neomycin derivative **18** under the same conditions described above, *i.e.* TMP, CH₃CN at 0 °C, we obtained the clean regiospecific formation of conjugate **19** in very good yield, as a mixture of diastereoisomers (Scheme 3).²²



Scheme 3. Synthesis of neomycin-hfVal-galactose conjugate **19**

The application of this procedure for the combinatorial synthesis of different aminoglycoside conjugates bearing a hfVal probe will be studied more in detail and reported in a forthcoming paper.

3. Conclusions

In conclusion, we have demonstrated that the MC sequential process recently developed by us can also be for the DOS of multivalent glycomimetics containing hfVal amino acid. Indeed, the reaction between *in situ* generated *N*-glycosyl, *N'*-*tert*-butyl carbodiimides or *N*, *N'*-diglycosyl carbodiimides with 4,4,4-trifluoro-3-Tfm-crotonic in the presence or absence of *N*-nucleophiles, such as α -aminoesters, peptides

or glycosyl amines, gave rise to the clean formation of di- or trivalent glycomimetics containing hfVal where the glycosyl moieties are tethered through different scaffolds, such as urea, hydantoin ring, and urea-peptide linkers. The MC sequential domino process is completely regioselective, high yielding, occurs under mild conditions, and is very general, working efficiently with different sugars and linkers. This process could also be exploited as a favourable tool to conjugate biologically important compounds with a sugar-hfVal tag. For instance, the reaction performed using an appropriately functionalized neomycin derivative such as *N*-nucleophile allowed us to obtain the synthesis of a neomycin-hfVal conjugate in a very efficient way. The combinatorial synthesis of different aminoglycoside-hfVal-sugar conjugates and the determination of their antibacterial activity and RNA interaction through ^{19}F NMR spectroscopy is ongoing in our laboratories and will be reported in a forthcoming paper.

4. Experimental Section

4.1 General Methods: Commercially available reagent-grade solvents were employed without purification. Primary glycosylazides **5** were prepared following reported procedures.¹¹ Glycosylisothiocyanate **15** were prepared by aza-Wittig reaction with CS_2 .²³ Glycosylamines **9** were prepared by catalytic hydrogenation of the corresponding azides. Neomycin derivative **18** was obtained as reported in ref. 19a. ^1H NMR spectra were run on spectrometers operating at 400 or 500 MHz. Chemical shifts are expressed in ppm (δ), using tetramethylsilane (TMS) as internal standard for ^1H and ^{13}C nuclei (δ_{H} and $\delta_{\text{C}} = 0.00$). ESIMS was performed with an Esquire 3000 plus ion-trap mass spectrometer equipped with an ESI source. The IR spectra were obtained by a Varian 640 high-performance FTIR spectrometer. Elemental analysis was obtained on FlashEA 1112 NC Analyzers. TLC was run on silica gel 60 F254 Merck. Flash chromatography (FC) was performed with silica gel 60 (60–200 mm, Merck).

4.2 Synthesis of divalent glycomimetics 1: general procedure. To a solution of glycosylazide **5** (1 equiv.) in CH_3CN (0.1 M) *tert*-butyl isocyanate (1.05 equiv.) **6** followed by Ph_3P (1.05 equiv.)

were added at rt. The solution was stirred until complete formation of the corresponding carbodiimide **7** was achieved (TLC monitoring). The temperature was lowered to 0 °C and TMP (1 equiv.), a solution of glycosylamine **9** (1 equiv.) in a minimum amount of CH₃CN followed by a solution of 4,4,4-trifluoro-3-Tfm-crotonic acid **8** (1 equiv.) in a minimum amount of CH₃CN were added. The temperature was slowly left to reach rt and the reaction, when finished (TLC monitoring, ca. 3 h), was quenched with a 1M solution of HCl. The mixture was extracted with AcOEt, the organic phases collected, dried over Na₂SO₄, filtered, the solvent removed under reduced pressure and the crude purified by flash chromatography.

1a: Major diastereoisomer: R_f = 0.29 (hexane/ AcOEt 80:20); FTIR (neat) ν 1779, 1756, 1726 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), δ : 8.44 (br s, 1H), 5.49-5.46 (m, 2H), 5.11 (s, 1H), 4.63 (d, J = 5.6 Hz, 1H), 4.59 (dd, J = 8.0 and 2.4 Hz, 2H), 4.53 (d, J = 5.6 Hz, 1H), 4.38-4.36 (m, 1H), 4.29 (dd, J = 5.2 and 2.4 Hz, 1H), 3.98-3.96 (m, 1H), 3.89-3.86 (m, 1H), 3.47 (s, 3H), 3.33-3.31 (m, 1H), 2.90-2.88 (m, 1H), 1.49 (s, 3H), 1.49 (s, 3H), 1.47 (s, 3H), 1.41 (s, 3H), 1.35 (s, 3H), 1.32 (s, 3H), 1.30 (s, 12H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 168.3, 154.9, 123.1 (q, J = 281.7 Hz, CF₃), 112.6, 111.5, 109.4, 108.5, 96.2, 87.0, 84.4, 71.4, 70.8, 70.5, 70.4, 56.0, 51.4, 45.9 (septet, J = 25.8 Hz), 40.4, 29.0, 26.3, 25.9, 25.7, 24.9, 24.7; ESI (m/z) 774.7 [M^+ +Na, (100)]; Anal. calcd. for C₃₁H₄₇F₆N₃O₁₁: C 49.53, H 6.30, N 5.59; found: C 49.53, H 6.32, N 5.60. **Minor diastereoisomer:** R_f = 0.22 (hexane/ AcOEt 80:20); FTIR (neat) ν 1781, 1754, 1722 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), δ : 7.12 (br s, 1H), 5.77 (s, 1H), 5.49 (d, J = 5.2 Hz, 2H), 5.11 (s, 1H), 4.98 (s, 1H), 4.60-4.56 (m, 2H), 4.41-4.39 (m, 1H), 4.32-4.28 (m, 2H), 4.15 (dd, J = 8.0 and 2.0 Hz, 1H), 3.97-3.94 (m, 1H), 3.47-3.44 (m, 3H), 3.41 (s, 3H), 3.24 (dd, J = 16.4 and 5.2 Hz, 1H), 1.49 (s, 3H), 1.49 (s, 3H), 1.47 (s, 3H), 1.45 (s, 3H), 1.35 (s, 9H), 1.34 (s, 3H), 1.32 (s, 3H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 168.0, 157.0, 123.3 (q, J = 281.5 Hz, CF₃), 112.9, 110.3, 109.4, 108.7, 96.3, 85.8, 84.5, 82.1, 71.3, 70.9, 70.5, 65.4, 55.5, 51.5, 46.9 (s, J = 26.1 Hz), 40.2, 29.0, 26.5, 26.0, 25.9, 25.1, 24.9, 24.4; ESI (m/z) 774.6 [M^+ +Na, (100)]; Anal. calcd. for C₃₁H₄₇F₆N₃O₁₁: C 49.53, H 6.30, N 5.59; found: C 49.55, H 6.31, N 5.58.

4.3 Synthesis of divalent glycomimetics 2: general procedure. To a solution of glycosylazide **5** (1 equiv.) in CH₃CN (0.1 M) a solution of glycosylisothiocyanate **15** (1.05 equiv.) in a minimum amount of CH₃CN followed by Ph₃P (1.05 equiv.) were added at rt. The solution was stirred until complete formation of the corresponding carbodiimide **16** was achieved (TLC monitoring). A solution of 4,4,4-trifluoro-3-Tfm-crotonic acid **8** (1 equiv.) in a minimum amount of CH₃CN was added at rt and the resulting solution stirred until the reaction was finished (TLC monitoring, *ca.* 3 h). The reaction was quenched with a 1M solution of HCl. The mixture was extracted with AcOEt, the organic phases collected, dried over Na₂SO₄, filtered, the solvent removed under reduced pressure and the crude purified by flash chromatography.

2a: Mixture of two diastereoisomers: R_f = 0.24 (hexane/ AcOEt 80:20); FTIR (neat) ν 1777, 1754, 1728 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), *major diastereoisomer* δ : 4.96 (s, 2H), 4.75 (s, 1H), 4.63-4.61 (m, 4H), 4.34-4.32 (m, 2H), 4.09 (dd, J = 12.0 and 4.8 Hz, 1H), 3.95-3.92 (m, 1H), 3.71-3.68 (m, 2H), 3.36 (s, 3H), 3.35 (s, 3H), 3.12 (dd, J = 12.0 and 6.4 Hz, 1H), 1.44 (s, 3H), 1.43 (s, 3H), 1.31 (s, 3H), 1.29 (s, 3H); *minor diastereoisomer* δ : 4.95 (s, 2H), 4.75 (s, 1H), 4.63-4.61 (m, 4H), 4.34-4.32 (m, 1H), 4.31-4.29 (m, 1H), 3.95-3.92 (m, 2H), 3.71-3.68 (m, 2H), 3.40-3.37 (m, 1H), 3.37 (s, 3H), 3.32 (s, 3H), 1.48 (s, 3H), 1.47 (s, 3H), 1.32 (s, 3H), 1.27 (s, 3H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 169.0, 156.5, 156.0, 113.1, 112.5, 110.5, 110.2, 110.0, 109.8, 85.2, 85.0, 84.6, 83.8, 83.4, 83.3, 82.3, 82.2, 82.1, 81.9, 56.0, 55.5, 55.4, 55.3, 55.1, 48.8 (septet, J = 28.2 Hz), 45.9, 44.7, 42.9, 42.7, 29.6, 26.6, 26.5, 26.4, 26.3, 25.1, 25.0, 24.9, 24.8; ESI (m/z) 645.2 [M^+ +Na, (100)]; Anal. calcd. for C₂₄H₃₂F₆N₂O₁₀: C 46.31, H 5.18, N 4.50; found: C 46.33, H 5.20, N 4.53.

4.4 Synthesis of divalent glycomimetics 2: general procedure. To a solution of glycosylazide **5** (1 equiv.) in CH₃CN (0.1 M) a solution of glycosylisothiocyanate **15** (1.05 equiv.) in a minimum amount of CH₃CN followed by Ph₃P (1.05 equiv.) were added at rt. The solution was stirred until

complete formation of the corresponding carbodiimide **16** was achieved (TLC monitoring). The temperature was lowered to 0 °C and TMP (2 equiv.), solid α -aminoester hydrochloride **17** (1 equiv.) followed by a solution of 4,4,4-trifluoro-3-Tfm-crotonic acid **8** (1 equiv.) in a minimum amount of CH₃CN were added. The temperature was slowly left to reach rt and the reaction, when finished (TLC monitoring, *ca.* 3 h), was quenched with a 1M solution of HCl. The mixture was extracted with AcOEt, the organic phases collected, dried over Na₂SO₄, filtered, the solvent removed under reduced pressure and the crude purified by flash chromatography.

3a: Major diastereoisomer: R_f = 0.34 (hexane/ AcOEt 70:30); FTIR (neat) ν 1769, 1753, 1723 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), δ : 8.11 (br s, 1H), 7.38-7.35 (m, 5H), 6.57 (s, 1H), 5.48 (d, J = 4.3 Hz, 1H), 5.44 (d, J = 4.2 Hz, 1H), 5.18 (d, J = 12.1 Hz, 1H), 5.11 (d, J = 12.1 Hz, 1H), 4.78 (br s, 1H), 4.65-4.56 (m, 3H), 4.31-4.28 (m, 2H), 4.25-4.22 (m, 2H), 4.11 (q, J = 7.0 Hz, 1H), 3.40-3.38 (m, 4H), 1.47 (s, 6H), 1.45 (s, 3H), 1.43 (s, 3H), 1.36 (s, 6H), 1.33 (s, 3H), 1.30-1.26 (m, 6H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 172.5, 167.5, 160.0, 136.0, 128.9, 128.6, 128.4, 123.7 (q, J = 282.2 Hz), 123.2 (q, J = 282.2 Hz), 110.0, 109.6, 109.4, 109.0, 96.6, 71.8, 71.3, 71.2, 70.9, 70.7, 67.2, 66.4, 65.4, 48.8, 47.4 (septet, J = 26.5 Hz), 41.6, 26.4, 26.3, 26.2, 26.0, 25.3, 25.2, 24.9, 24.6, 17.9; ESI (m/z) 936.3 [M^+ +Na, (100)]; Anal. calcd. for C₄₀H₅₃F₆N₃O₁₄: C 52.57, H 5.85, N 4.60; found: C 52.58, H 5.82, N 4.62. **Minor diastereoisomer:** R_f = 0.30 (hexane/ AcOEt 70:30); FTIR (neat) ν 1770, 1761, 1721 cm⁻¹; ¹H NMR (400 MHz, CDCl₃), δ : 7.65 (br s, 1H), 7.38-7.35 (m, 5H), 6.34 (br s, 1H), 5.49 (d, J = 5.08 Hz, 1H), 5.47 (d, J = 5.03 Hz, 1H), 5.18 (d, J = 12.1 Hz, 1H), 5.11 (d, J = 12.1 Hz, 1H), 4.56-4.54 (m, 2H), 5.45 (septet, J = 7.0 Hz, 1H), 4.30-4.26 (m, 2H), 4.22-4.19 (m, 2H), 3.97-3.94 (m, 1H), 3.85-3.83 (m, 1H), 3.55-3.54 (m, 1H), 3.23-3.21 (m, 2H), 1.44 (s, 6H), 1.42 (s, 3H), 1.36 (s, 6H), 1.33 (s, 3H), 1.13 (s, 6H), 1.29 (s, 3H), 1.20 (s, 3H); ¹³C NMR (100.6 MHz, CDCl₃) δ : 171.9, 168.0, 135.9, 128.9, 128.7, 128.5, 123.5 (q, J = 281.0 Hz), 123.3 (q, J = 282.2 Hz), 109.8, 109.6, 109.5, 108.9, 96.7, 96.6, 96.5, 72.2, 71.9, 71.3, 70.94, 70.91, 69.2, 67.3, 67.2, 49.4, 46.6 (septet, J = 26.7 Hz), 41.5, 26.4, 26.3, 26.1, 25.3, 24.8,

24.5; ESI (m/z) 936.2 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{40}H_{53}F_6N_3O_{14}$: C 52.57, H 5.85, N 4.60; found: C 52.59, H 5.86, N 4.58.

4.5 Synthesis of trivalent glycomimetics 4: general procedure. To a solution of glycosylazide **5** (1 equiv.) in CH_3CN (0.1 M) a solution of glycosylisothiocyanate **15** (1.05 equiv.) in a minimum amount of CH_3CN followed by Ph_3P (1.05 equiv.) were added at rt. The solution was stirred until complete formation of the corresponding carbodiimide **16** was achieved (TLC monitoring). The temperature was lowered to 0 °C and TMP (2 equiv.), a solution of glycosylamine **9** (1 equiv.) followed by a solution of 4,4,4-trifluoro-3-Tfm-crotonic acid **8** (1 equiv.) in a minimum amount of CH_3CN were added. The temperature was slowly left to reach rt and the reaction, when finished (TLC monitoring, *ca.* 3 h), was quenched with a 1M solution of HCl. The mixture was extracted with AcOEt, the organic phases collected, dried over Na_2SO_4 , filtered, the solvent removed under reduced pressure and the crude purified by flash chromatography.

4a: Mixture of two diastereoisomers: R_f = 0.21 (hexane/ AcOEt 20:80); FTIR (neat) ν 1778, 1755, 1721 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$), *major diastereoisomer* δ : 6.01 (br s, 1H), 5.46 (d, J = 5.1 Hz, 1H), 5.14 (s, 1H), 4.97 (s, 1H), 4.62-4.54 (m, 6H), 4.18-4.15 (m, 3H), 4.14-4.11 (m, 2H), 3.91-3.90 (m, 1H), 3.84 (d, J = 9.2 Hz, 1H), 3.78-3.76 (m, 1H), 3.53-3.50 (m, 2H), 3.42 (s, 3H), 3.37 (s, 3H), 3.25-3.22 (m, 2H), 3.08-3.05 (m, 1H), 1.48 (s, 3H), 1.45 (s, 3H), 1.39 (s, 3H), 1.34 (s, 3H), 1.31 (s, 3H), 1.30 (s, 3H), 1.29 (s, 3H), 1.28 (s, 3H); *minor diastereoisomer* δ : 6.38 (br s, 1H), 5.49 (d, J = 5.2 Hz, 1H), 5.14 (s, 1H), 4.96 (s, 1H), 4.62-4.54 (m, 6H), 4.18-4.15 (m, 3H), 4.14-4.11 (m, 2H), 3.91-3.90 (m, 1H), 3.84 (d, J = 9.2 Hz, 1H), 3.78-3.76 (m, 1H), 3.53-3.50 (m, 2H), 3.48 (s, 3H), 3.41 (s, 3H), 3.25-3.22 (m, 2H), 3.08-3.05 (m, 1H), 1.47 (s, 3H), 1.44 (s, 3H), 1.39 (s, 3H), 1.34 (s, 3H), 1.31 (s, 3H), 1.30 (s, 3H), 1.29 (s, 3H), 1.28 (s, 3H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 168.1, 167.6, 158.2, 156.0, 123.2 (q, J = 280.7 Hz), 123.0 (q, J = 281.4), 113.1, 112.8, 112.4, 112.3, 111.4, 110.3, 110.2, 110.0, 109.4, 109.3,

108.6, 108.5, 96.3, 96.2, 86.5, 86.2, 85.6, 85.4, 85.3, 84.5, 82.0, 81.9, 81.7, 71.5, 71.4, 70.9, 70.5, 66.8, 65.6, 60.3, 57.9, 56.8, 55.8, 55.2, 55.1, 53.8, 51.4, 47.0, 46.7, 46.5, 46.3, 46.0, 45.8, 44.2, 43.7, 40.6, 40.3, 26.5, 26.4, 26.35, 26.31, 25.9, 25.8, 25.7, 25.0, 24.9, 24.8, 24.7, 24.6, 24.4, 24.3; ESI (m/z) 904.7 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{36}H_{53}F_6N_3O_{15}$: C 49.03, H 6.06, N 4.77; found: C 49.00, H 6.08, N 4.76.

Acknowledgments. Politecnico di Milano and ICRM-CNR are gratefully acknowledged for economic support.

Supporting Information Available: characterization of compounds **1b-f**, **2b**, **3b,c**, **4b**, **13** and **19**. Copies of 1H , ^{13}C NMR, and ESI-MS spectra of all new compounds.

References

- 1 a) Wang, J.; Sanchez-Rosello, M.; Acena, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432; b) Isanbor, C.; O'Hagan, D.; *J. Fluorine Chem.* **2006**, *127*, 303. See also the thematic issue: *Curr. Top. Med. Chem.* **2014**, *14*, 855.
- 2 a) Salwiczek, M.; Nyakamura, E. K.; Gerling, U. I. M.; Ye, S.; Kokscho, B. *Chem. Soc. Rev.* **2012**, *41*, 2135; b) Marsh, E. N. G. *Acc. Chem. Res.* **2014**, *47*, 2878.
- 3 a) Cobb, S. L.; Murphy, C. D. *J. Fluorine Chem.* **2009**, *130*, 132. b) Ulrich, A. S. *Prog. Nucl. Magn. Reson. Spectrosc.* **2006**, *46*, 1; c) Ahrens, E. T.; Zhong, J. *NMR Biomed.* **2013**, *26*, 860; d) Tirota, I.; Dichiarante, V.; Pigliacelli, C.; Cavallo, G.; Terraneo, G.; Baldelli Bombelli, F.; Metrangolo, P.; Resnati, G. *Chem. Rev.* **2015**, *115*, 1106.
- 4 O'Hagan, D.; Harper, B. *J. Fluorine Chem.* **1999**, *100*, 127.

- 5 a) Kiessling, L. L.; Phol, N. L. *Chem. Biol.* **1996**, *3*, 71; b) Fasting, C.; Schalley, C. A.; Weber, M.; Seitz, O.; Hecht, S.; Kokscho, B.; Dornedde, J.; Graf, C.; Knapp, E.-W.; Haag, R. *Angew. Chem., Int. Ed.* **2012**, *51*, 10472; c) Pieters, R. J. **2009**, *7*, 2013; c) Bernardi, A.; Jimenez-Barbero, J.; Casnati, A.; De Castro, C.; Darbre, T.; Fieschi, F.; Finne, J.; Funken, H.; Jaeger, K.-E.; Lahmann, M.; Lindhorst, T. K.; Marradi, M.; Messner, P.; Molinaro, A.; Murphy, P. V.; Nativi, C.; Oscarson, S.; Penade's, S.; Peri, F.; Pieters, R. J.; Renaudet, O.; Reymond, J.-L.; Richichi, B.; Rojo, J.; Sansone, F.; Schaffer, C.; Turnbull, W. B.; Velasco-Torrijos, T.; Vidal, S.; Vincent, S.; Wennekes, T.; Zuilhofxy, H.; Imberty, A. *Chem. Soc. Rev.* **2013**, *42*, 4709.
- 6 Compain, P.; Bodlenner, A.; *ChemBioChem* **2014**, *15*, 1239.
- 7 a) Schreiber, S. L. *Science*, **2000**, *287*, 1964; b) Spring, D. R. *Org. Biomol. Chem.* **2003**, *1*, 3867; c) Burke, M. D.; Schreiber, S. L. *Angew. Chem. Int. Ed.* **2004**, *43*, 46; d) Tan, D. S. *Nat. Chem. Biol.* **2005**, *1*, 74; e) Galloway, W. R. J.; Isidro-Llobet, A.; Spring, D. R. *Nat. Commun.* **2008**, *80*, 1.
- 8 a) Dömling, A.; Ugi, I. *Angew. Chem. Int. Ed.* **2000**, *39*, 3168; b) Dömling, A. *Chem. Rev.* **2006**, *106*, 17; c) Touré, B. B.; Hall, D. G. *Chem. Rev.* **2009**, *109*, 4439; d) Ganem, B. *Acc. Chem. Res.* **2009**, *42*, 463; e) Biggs-Houck, J. E.; Younai, A.; Shaw, J. T. *Curr. Opin. Chem. Biol.* **2010**, *14*, 371; f) Ruijter, E.; Scheffelaar, R.; Orru, R. V. A. *Angew Chem. Int. Ed.* **2011**, *50*, 6234.
- 9 Concerning the DOS of non-fluorinated multivalent glycomimetics through MC processes, see: a) Lockhoff, O. *Angew. Chem. Int. Ed.* **1998**, *37*, 3436; b) Westermann, B.; Dörner, S. *Chem. Commun.* **2005**, 2116.
- 10 a) Volonterio, A.; Ramirez de Arellano, C.; Zanda, M. *J. Org. Chem.* **2005**, *70*, 2161; b) Olimpieri, F.; Fustero, S.; Volonterio, A.; Zanda, M. *Synthesis* **2010**, *4*, 651; c) Bellucci, M. C.; Marcelli, T.; Scaglioni, L.; Volonterio, A. *RSC Adv.* **2011**, *1*, 1250; d) Olimpieri, F.; Bellucci, M. C.; Marcelli, T.; Volonterio, A. *Org. Biomol. Chem.* **2012**, *10*, 9538

- 11 a) Bellucci, M. C.; Volonterio, A. *Adv. Synth. Catal.* **2010**, 252, 2791; b) Bellucci, M. C.; Ghilardi, A.; Volonterio, A. *Org. Biomol. Chem.* **2011**, 9, 8379; c) Bellucci, M. C.; Terraneo, G.; Volonterio, A. *Org. Biomol. Chem.* **2013**, 11, 2421; d) Bellucci, M. C.; Volonterio, A. *Eur. J. Org. Chem.* **2014**, 2386.
- 12 Bellucci, M. C.; Sani, M.; Sganappa, A.; Volonterio, A. *ACS Comb. Chem.* **2014**, 16, 711.
- 13 a) Hsieh, K.-h.; Needleman, P.; Marshall, G. R. *J. Med. Chem.* **1987**, 30, 1097; b) Eberle, M. K.; Keese, R.; Stoekli-Evans, H. *Helv. Chim. Acta* **1998**, 81, 182; c) Eberle, M. K.; Keese, R. *Helv. Chim. Acta.* **2010**, 93, 1583.
- 14 For a DFT study on the mechanism of this process, see: Marcelli, T.; Olimpieri, F.; Volonterio, A. *Org. Biomol. Chem.* **2011**, 9, 5156.
- 15 The chiral aminosugar **9a** steps in the reaction once the new stereogenic center at the hfVal moiety is already formed.
- 16 Samantaray, S.; Marathe, U.; Dasgupta, S.; Nandicoori, V. K.; Roy, R. P.; *J. Am. Chem. Soc.* **2008**, 130, 2132.
- 17 In these cases we did not obtained an equimolar ratio of diastereoisomers because the intramolecular aza-Michael nucleophilic attack arose from the chiral *N*-sugar moiety of the carbodiimides. However, due to the distance between the reactive electronpoor C=C form the stereogenic centers of the sugars, the diastereoselections are low, form *ca.* 3:1 in the case of galactose to *ca.* 1.5:1 in the case of the other sugars. Since the mechanism and the stereochemical behaviour of this process is related to that described in a previous paper (see ref. 11a), the stereochemistry of the new formed stereocenter of the major diastereoisomer was assessed to be (*R*) on the basis of the structure of the corresponding peptide-sugar conjugate incorporating hexafluorovaline which was determined by X-ray diffraction
- 18 a) Tor, Y.; Hermann, T.; Westhof, E. *Chem. Biol.* **1998**, 5, R277; b) Tor, Y. *ChemBioChem* **2003**, 4, 998; c) Houghton, J. L.; Green, K. D.; Chen, W.; Garneau-Tsodikova, S.

- ChemBioChem.* **2010**, *11*, 880; d) McCoy, L. S.; Xie, Y.; Tor, Y. *Wiley Interdiscip. Rev. RNA* **2011**, *2*, 209.
- 19 a) Luedtke, N. W.; Liu, Q.; Tor, Y. *Biochemistry* **2003**, *42*, 11391; b) Fridman, M.; Belakhov, V.; Yaron, S.; Baasov, T.; *Org. Lett.* **2003**, *5*, 3575; c) Blount, K. F.; Tor, Y. *ChemBioChem.* **2006**, *7*, 1612; d) Jeon, J.; Lee, K. H.; Rao, J. *Chem. Commun.* **2012**, *48*, 10034; e) Kumar, S.; Xue, L.; Arya, D. P. *J. Am. Chem. Soc.* **2010**, *133*, 7361.
- 20 Lombès, T.; Mounné, R.; Larue, V.; Prost, E.; Catala, M.; Lecourt, T.; Dardel, F.; Micouin, L.; Tisné, C. *Angew. Chem. Int. Ed.* **2012**, *51*, 9530.
- 21 a) Park, W. K. C.; Auer, M.; Jaksche, H.; Wong, C.-H. *J. Am. Chem. Soc.* **1996**, *118*, 10150; b) Nunns, C. L.; Spence, L. A.; Slater, M. J.; Berrisford, D. J. *Tetrahedron Lett.* **1999**, *40*, 9341.
- 22 Neomycin derivative **18** is very soluble in CH₃CN at 0°C, thus the protection of the hydroxyls was not necessary to obtain high yield.
- 23 Garcia-Fernandez, J. M.; Ortiz Mellet, C.; Fuentes, J. *J. Org. Chem.* 1993, **58**, 5192.

Supporting Information**Multicomponent Diversity Oriented Synthesis of
Multivalent Glycomimetics Containing
Hexafluorovaline**

Maria Cristina Bellucci, Aurora Sganappa, Monica Sani, and Alessandro Volonterio,*

Department of Food, Environmental and Nutritional Sciences, Università degli Studi di Milano, via

Celoria 2, 20133 Milano, Italy; C.N.R. Istituto di Chimica del Riconoscimento Molecolare, via

Mancinelli 7, 20131 Milano, Italy; Department of Chemistry, Materials, and Chemical Engineer

“G. Natta”, Politecnico di Milano, via Mancinelli 7, 20131 Milano, Italy

alessandro.volonterio@polimi.it

Table of Contents:

Pages S2-S15 Characterization of all new compounds.

Pages S15-S91 ^1H NMR, ^{13}C NMR, and MS spectra of all new compounds.

13: Mixture of two diastereoisomers: $R_f = 0.38$ (hexane/ AcOEt 60:40); ^1H NMR (400 MHz, CDCl_3), major diastereoisomer δ : 8.25 (br s, 1H), 5.48 (t, $J = 4.8$ Hz, 1H), 4.67 (m, 2H), 4.57 (d, $J = 2.0$ Hz, 1H), 4.36 (m, 1H), 4.27 (m, 1H), 4.13 (m, 1H), 3.97 (m, 1H), 3.82 (m, 2H), 3.57 (m, 2H), 3.35 (m, 1H), 1.44 (m, 6H), 1.28 (m, 18H); minor diastereoisomer δ : 8.25 (br s, 1H), 5.48 (t, $J = 4.8$ Hz, 1H), 4.67 (m, 2H), 4.55 (d, $J = 2.4$ Hz, 1H), 4.36 (m, 1H), 4.27 (m, 1H), 4.13 (m, 1H), 3.97 (m, 1H), 3.82 (m, 2H), 3.57 (m, 2H), 3.23 (m, 1H), 1.44 (m, 6H), 1.28 (m, 18H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 169.3, 158.1, 123.0 (q, $J = 281.2$ Hz, CF_3), 122.8 (q, $J = 284.1$ Hz, CF_3), 109.3, 108.4, 108.3, 96.3, 96.2, 71.2, 70.8, 70.6, 65.9, 65.8, 56.2, 55.9, 50.2, 50.0, 47.0 (s, $J = 25.9$ Hz), 40.0, 39.8, 25.9, 25.8, 25.7, 24.8, 24.3, 23.0, 21.2, 20.7, 20.5; ESI (m/z) 616.3 [$\text{M}^+ + \text{Na}$, (100)]; Anal. calcd. for $\text{C}_{24}\text{H}_{37}\text{F}_6\text{N}_3\text{O}_7$: C 48.56, H 6.28, N 7.08; found: C 48.58, H 6.30, N 7.09.

1b: Mixture of two diastereoisomers: $R_f = 0.27$ (hexane/ AcOEt 60:40); ^1H NMR (400 MHz, CDCl_3), major diastereoisomer δ : 8.44 (br s, 1H), 5.47 (s, 1H), 5.26-5.19 (m, 2H), 5.07-5.492 (m, 2H), 4.60-4.53 (m, 4H), 4.30 (m, 1H), 4.22 (m, 1H), 4.05-4.00 (m, 2H), 3.78 (m, 1H), 3.45 (m, 1H), 3.40 (s, 3H), 3.20 (dd, $J = 16.0$ and 3.6 Hz, 1H), 2.03-1.94 (four s, 12H), 1.47 (s, 3H), 1.31 (s, 3H), 1.30 (s, 9H); minor diastereoisomer δ : 8.42 (br s, 1H), 5.47 (s, 1H), 5.26-5.19 (m, 2H), 5.07-5.492 (m, 2H), 4.60-4.53 (m, 4H), 4.30 (m, 1H), 4.22 (m, 1H), 4.05-4.00 (m, 2H), 3.78 (m, 1H), 3.45 (m, 1H), 3.38 (s, 3H), 3.00 (m, 1H), 2.03-1.94 (four s, 12H), 1.53 (s, 3H), 1.31 (s, 3H), 1.28 (s, 9H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 170.45, 170.41, 169.9, 169.8, 169.7, 169.6, 169.4, 168.3, 122.9 (q, $J = 279.7$ Hz), 122.5 (q, $J = 278.5$ Hz), 113.4, 112.8, 110.7, 110.3, 87.0, 86.6, 85.2, 84.1, 84.0, 83.7, 82.0, 78.2, 73.8, 73.3, 73.2, 73.1, 70.5, 70.4, 68.3, 68.2, 61.8, 55.8, 55.3, 54.3, 51.6, 51.5, 47.0 (septet, $J = 26.1$ Hz), 45.8 (septet, $J = 28.2$ Hz), 29.4, 28.9, 28.8, 26.5, 26.3, 24.9, 24.7, 20.5, 20.4, 20.3, 20.1; ESI (m/z) 862.6 [$\text{M}^+ + \text{Na}$, (100)]; Anal. calcd. for $\text{C}_{33}\text{H}_{47}\text{F}_6\text{N}_3\text{O}_{15}$: C 47.20, H 5.64, N 5.00; found: C 47.23, H 5.62, N 4.99.

1c: Mixture of two diastereoisomers: $R_f = 0.24$ (hexane/ AcOEt 80:20); ^1H NMR (400 MHz, CDCl_3), major diastereoisomer δ : 5.53 (d, $J = 4.8$ Hz, 1H), 4.96 (s, 1H), 4.60 (m, 4H), 4.32 (m, 1H), 4.24 (m, 2H), 3.98 (m, 1H), 3.68 (m, 2H), 3.50 (m, 1H), 3.37 (s, 3H), 3.36 (m, 1H), 3.10 (m, 2H), 1.46-1.45 (three s, 9H), 1.35-1.33 (three s, 9H), 1.32 (s, 9H); minor diastereoisomer δ : 5.52 (d, $J = 4.6$ Hz, 1H), 4.95 (s, 1H), 4.60 (m, 4H), 4.32 (m, 1H), 4.24 (m, 2H), 4.01 (m, 1H), 3.71 (m, 2H), 3.50 (m, 1H), 3.37 (s, 3H), 3.38 (m, 1H), 3.10 (m, 2H), 1.46-1.45 (three s, 9H), 1.35-1.33 (three s, 9H), 1.29 (s, 9H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 168.6, 167.7, 162.6, 158.3, 156.2, 123.0 (q, $J = 283.7$ Hz), 112.35, 112.30, 109.8, 109.7, 109.6, 109.5, 109.4, 109.2, 96.3, 96.2, 96.1, 85.5, 85.4, 85.3, 85.2, 82.2, 71.6, 71.5, 70.8, 70.7, 70.6, 70.5, 70.4, 70.3, 68.6, 55.1, 51.25, 51.22, 46.4 (septet, $J = 26.1$ Hz), 42.7, 36.4, 29.4, 28.9, 28.8, 28.3, 26.4, 26.3, 26.0, 25.9, 25.8, 25.7, 25.6, 25.0, 24.8, 24.7, 24.3, 24.1; ESI (m/z) 774.6 [$\text{M}^+ + \text{Na}$, (100)]; Anal. calcd. for $\text{C}_{31}\text{H}_{47}\text{F}_6\text{N}_3\text{O}_{11}$: C 49.53, H 6.30, N 5.59; found: C 49.55, H 6.31, N 5.60.

1d: Mixture of two diastereoisomers: $R_f = 0.58$ (hexane/ AcOEt 50:50); ^1H NMR (400 MHz, CDCl_3), major diastereoisomer δ : 7.27 (m, 6H), 6.27 (br s, 1H), 5.63 (d, $J = 10.4$ Hz, 1H), 5.32 (m, 1H), 4.94 (m, 2H), 4.86 (m, 1H), 4.75 (s, 1H), 4.56-4.28 (m, 6H), 4.19 (m, 1H), 3.52 (m, 2H), 3.47 (s, 3H), 3.21 (m, 2H), 1.84 (s, 3H), 1.46 (s, 3H), 1.35-1.12 (three s, 30H); minor diastereoisomer δ : 7.27 (m, 6H), 5.98 (br s, 1H), 5.60 (d, $J = 10.4$ Hz, 1H), 5.32 (m, 1H), 4.94 (m, 2H), 4.86 (m, 1H), 4.78 (s, 1H), 4.56-4.28 (m, 6H), 4.08 (m, 1H), 3.52 (m, 2H), 3.32 (s, 3H), 3.21 (m, 2H), 1.83 (s, 3H), 1.42 (s, 3H), 1.35-1.12 (three s, 30H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 178.6, 178.5, 177.0, 176.9, 169.7, 167.6, 167.3, 162.5, 158.7, 135.9, 128.8, 128.7, 128.6, 128.5, 128.2, 128.0, 127.9,

123.0 (q, $J = 284.8$ Hz), 122.7 (q, $J = 285.7$ Hz), 112.4, 109.9, 109.8, 96.4, 95.9, 95.8, 85.7, 85.5, 85.4, 85.3, 82.1, 82.0, 70.8, 70.4, 70.3, 70.2, 70.0, 69.9, 69.8, 69.7, 69.6, 69.4, 68.9, 68.5, 55.3, 55.1, 51.8, 51.7, 51.1, 50.9, 47.7 (septet, $J = 25.1$ Hz), 43.5, 42.7, 42.5, 38.9, 38.8, 36.4, 31.4, 29.6, 29.3, 29.1, 29.0, 28.6, 28.4, 28.3, 27.0, 26.9, 26.5, 26.4, 26.3, 24.8, 23.0, 22.9, 22.8; ESI (m/z) 993.8 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{44}H_{64}F_6N_4O_{13}$: C 54.43, H 6.64, N 5.77; found: C 54.45, H 6.65, N 5.80.

1e: Mixture of two diastereoisomers: $R_f = 0.37$ (hexane/ AcOEt 50:50); 1H NMR (400 MHz, $CDCl_3$), major diastereoisomer δ : 7.32 (br s, 1H), 5.28 (m, 2H), 5.22 (m, 1H), 4.97 (d, $J = 2.8$ Hz, 1H), 4.80 (d, $J = 2.0$ Hz, 1H), 4.36 (m, 1H), 4.27 (m, 1H), 4.13 (m, 1H), 3.97 (m, 1H), 3.82 (m, 2H), 3.57 (m, 2H), 3.35 (m, 1H), 1.44 (m, 6H), 1.28 (m, 18H); minor diastereoisomer δ : 8.25 (br s, 1H), 5.48 (t, $J = 4.8$ Hz, 1H), 4.67 (m, 2H), 4.55 (d, $J = 2.4$ Hz, 1H), 4.36 (m, 1H), 4.27 (m, 1H), 4.13 (m, 1H), 3.97 (m, 1H), 3.82 (m, 2H), 3.57 (m, 2H), 3.23 (m, 1H), 1.44 (m, 6H), 1.28 (m, 18H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 169.3, 158.1, 123.0 (q, $J = 281.2$ Hz, CF_3), 122.8 (q, $J = 284.1$ Hz, CF_3), 109.3, 108.4, 108.3, 96.3, 96.2, 71.2, 70.8, 70.6, 65.9, 65.8, 56.2, 55.9, 50.2, 50.0, 47.0 (s, $J = 25.9$ Hz), 40.0, 39.8, 25.9, 25.8, 25.7, 24.8, 24.3, 23.0, 21.2, 20.7, 20.5; ESI (m/z) 616.3 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{36}H_{53}F_6N_3O_{16}$: C 48.16, H 5.95, N 4.68; found: C 48.17, H 5.97, N 4.67.

1f: Major diastereoisomer: $R_f = 0.45$ (hexane/ AcOEt 50:50); 1H NMR (400 MHz, $CDCl_3$), δ : 7.80 (s, 1H), 6.25 (br s, 1H), 5.83 (d, $J = 9.6$ Hz, 1H), 5.42 (t, $J = 9.6$ Hz, 1H), 5.37 (t, $J = 9.6$ Hz, 1H), 5.26 (t, $J = 9.6$ Hz, 1H), 4.93 (s, 1H), 4.49 (m, 2H), 4.42 (br s, 1H), 4.37 (dd, $J = 12.2$ and 4.4 Hz, 1H), 4.25 (m, 1H), 4.21 (m, 2H), 4.00 (m, 2H), 3.95 (m, 1H), 3.34 (s, 3H), 3.22 (m, 1H), 2.11 (s, 3H), 2.08 (s, 3H), 2.05 (s, 3H), 1.89 (s, 3H), 1.60 (s, 3H), 1.45 (s, 3H), 1.33 (s, 9H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 170.3, 169.9, 169.2, 168.7, 167.9, 155.9, 144.8, 122.8 (q, $J = 282.1$ Hz), 121.6, 112.4, 109.9, 86.1, 85.9, 85.2, 85.1, 81.8, 75.4, 72.5, 70.9, 67.8, 61.2, 60.3, 55.2, 51.5, 46.6 (septet, $J = 27.2$ Hz), 42.5, 29.6, 28.9, 28.3, 26.4, 24.8, 20.9, 20.4, 20.0, 14.2; ESI (m/z) 943.7 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{36}H_{50}F_6N_6O_{15}$: C 46.96, H 5.47, N 9.13; found: C 46.98, H 5.48, N 9.11.

Minor diastereoisomer: $R_f = 0.38$ (hexane/ AcOEt 50:50); 1H NMR (400 MHz, $CDCl_3$), δ : 7.71 (s, 1H), 5.97 (br s, 1H), 5.81 (d, $J = 9.2$ Hz, 1H), 5.43 (t, $J = 9.2$ Hz, 1H), 5.35 (t, $J = 9.2$ Hz, 1H), 5.25 (t, $J = 9.2$ Hz, 1H), 4.97 (s, 1H), 4.57 (d, $J = 6.0$ Hz, 1H), 4.53 (d, $J = 6.0$ Hz, 1H), 4.49 (br s, 1H), 4.39 (d, $J = 13.2$ Hz, 1H), 4.32 (dd, $J = 12.8$ and 5.2 Hz, 1H), 4.21 (m, 1H), 4.18 (d, $J = 13.2$ Hz, 1H), 4.00 (m, 2H), 3.67 (m, 1H), 3.41 (s, 3H), 3.23 (m, 1H), 2.11 (s, 3H), 2.08 (s, 3H), 2.05 (s, 3H), 1.86 (s, 3H), 1.68 (s, 3H), 1.47 (s, 3H), 1.30 (s, 9H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 170.4, 169.9, 169.3, 168.5, 167.9, 155.9, 144.5, 123.0 (q, $J = 282.7$ Hz), 121.2, 112.4, 109.1, 86.0, 85.6, 85.5, 82.0, 75.4, 75.3, 72.5, 70.4, 67.6, 61.4, 55.5, 51.5, 46.7 (septet, $J = 26.2$ Hz), 42.3, 29.7, 28.9, 26.3, 24.7, 20.6, 20.5, 20.0; ESI (m/z) 943.7 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{36}H_{50}F_6N_6O_{15}$: C 46.96, H 5.47, N 9.13; found: C 46.96, H 5.49, N 9.14.

2b: Mixture of two diastereoisomers: $R_f = 0.24$ (hexane/ AcOEt 80:20); 1H NMR (400 MHz, $CDCl_3$), major diastereoisomer δ : 5.41 (d, $J = 5.2$ Hz, 1H), 5.38 (d, $J = 4.8$ Hz, 1H), 4.62 (s, 1H), 4.56 (m, 2H), 4.24 (m, 2H), 4.14 (m, 4H), 3.82 (m, 2H), 3.46 (m, 1H), 3.21 (dd, $J = 14.8$ and 5.2 Hz, 1H), 1.44-1.37 (four s, 12H), 1.30-1.24 (four s, 12H); minor diastereoisomer δ : 5.46 (d, $J = 5.2$ Hz, 1H), 5.38 (d, $J = 4.8$ Hz, 1H), 4.91 (s, 1H), 4.56 (m, 2H), 4.24 (m, 2H), 4.14 (m, 4H), 3.82 (m, 2H), 3.45 (m, 2H), 1.44-1.37 (four s, 12H), 1.30-1.24 (four s, 12H); ^{13}C NMR (100.6 MHz, $CDCl_3$)

δ : 169.7, 169.6, 169.5, 157.2, 156.8, 122.8 (q, $J = 281.5$ Hz), 122.3 (q, $J = 280.8$ Hz), 109.9, 109.8, 109.7, 109.5, 109.0, 108.7, 108.7, 108.5, 96.3, 96.23, 96.20, 96.1, 71.5, 71.4, 71.3, 71.1, 70.9, 70.8, 70.7, 70.5, 70.44, 70.40, 70.3, 66.9, 64.6, 64.3, 63.6, 57.2, 57.1, 55.4, 49.2 (septet, $J = 28.5$ Hz), 48.9 (septet, $J = 27.9$ Hz), 42.2, 42.1, 42.0, 40.2, 39.9, 29.6, 26.0, 25.9, 25.8, 25.76, 25.70, 25.05, 25.03, 24.7, 24.6, 24.55, 24.51, 24.3 ; ESI (m/z) 757.1 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{30}H_{40}F_6N_2O_{12}$: C 49.05, H 5.49, N 3.81; found: C 49.02, H 5.50, N 3.79.




3b: *Mixture of two diastereoisomers:* $R_f = 0.35$ (hexane/ AcOEt 60:40); 1H NMR (400 MHz, $CDCl_3$), *major diastereoisomer* δ : 8.21 (br s, 1H), 7.34 (m, 5H), 6.60 (br s, 1H), 5.16 (d, $J = 12.0$ Hz, 1H), 5.12 (d, $J = 12.0$ Hz, 1H), 5.00 (s, 1H), 4.97 (s, 1H), 4.62-4.57 (m, 7H), 4.28 (t, $J = 6.0$ Hz, 1H), 4.00 (m, 1H), 3.48 (m, 1H), 3.44 (s, 3H), 3.40 (s, 3H), 3.33 (m, 1H), 1.63 (m, 3H), 1.47 (s, 3H), 1.46 (s, 3H), 1.30 (s, 3H), 1.29 (s, 3H), 0.91 (m, 6H); *minor diastereoisomer* δ : 7.52 (br s, 1H), 7.34 (m, 5H), 6.16 (t, $J = 5.6$ Hz, 1H), 5.17 (d, $J = 12.0$ Hz, 1H), 5.11 (d, $J = 12.0$ Hz, 1H), 5.04 (s, 1H), 4.97 (s, 1H), 4.62-4.57 (m, 7H), 4.34 (dd, $J = 8.4$ and 6.0 Hz, 1H), 4.00 (m, 1H), 3.48 (m, 1H), 3.41 (s, 3H), 3.40 (s, 3H), 3.06 (m, 1H), 1.63 (m, 3H), 1.47 (s, 3H), 1.46 (s, 3H), 1.33 (s, 3H), 1.31 (s, 3H), 0.91 (m, 6H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 171.9, 171.6, 167.8, 167.3, 158.3, 156.2, 135.6, 135.5, 128.5, 128.4, 128.3, 128.28, 128.21, 128.1, 128.0, 122.7 (q, $J = 289.2$ Hz), 122.4 (q, $J = 282.6$ Hz), 113.1, 112.9, 112.3, 112.2, 111.4, 110.4, 110.3, 110.2, 87.0, 86.3, 86.2, 85.6, 85.4, 84.6, 84.4, 82.3, 82.0, 66.9, 66.8, 56.0, 55.8, 55.3, 55.2, 51.5, 51.4, 46.6 (septet, $J = 26.8$ Hz), 46.0 (septet, $J = 26.5$ Hz), 44.0, 43.8, 41.2, 40.8, 26.5, 26.4, 26.3, 24.9, 24.8, 24.7, 24.6, 22.7, 21.7, 21.6; ESI (m/z) 866.1 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{37}H_{51}F_6N_3O_{12}$: C 52.67, H 6.09, N 4.98; found: C 52.68, H 6.10, N 4.96.



3c: *Major diastereoisomer:* $R_f = 0.34$ (hexane/ AcOEt 40:60); 1H NMR (500 MHz, $CDCl_3$), δ : 7.58 (br s, 1H), 7.37-7.22 (m, 10H), 6.12 (br s, 1H), 5.18 (d, $J = 12.5$ Hz, 1H), 5.14 (d, $J = 12.5$ Hz, 1H), 4.97 (s, 1H), 4.86 (m, 1H), 4.64-4.47 (m, 8H), 4.28 (m, 1H), 3.75-3.68 (m, 2H), 3.44 (s, 3H), 3.41 (m, 2H), 2.99 (m, 2H), 2.96 (m, 3H), 1.71 (m, 1H), 1.57 (m, 2H), 1.49 (s, 3H), 1.45 (s, 3H), 1.31 (s, 3H), 1.26 (s, 3H), 0.92 (d, $J = 6.5$ Hz, 3H), 0.90 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 171.8, 170.6, 167.9, 137.0, 135.8, 128.7, 128.5, 128.4, 128.1, 128.0, 126.6, 123.6 (q, $J = 280.7$ Hz), 123.2 (q, $J = 281.7$ Hz), 113.1, 112.4, 110.9, 110.8, 86.3, 85.8, 84.1, 81.9, 81.3, 66.7, 58.8, 55.8, 55.5, 54.2, 53.4, 51.4, 46.1 (septet, $J = 26.2$ Hz), 43.4, 40.8, 37.0, 29.6, 26.4, 26.2, 24.8, 24.4, 22.6, 22.1; ESI (m/z) 1013.4 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{46}H_{60}F_6N_4O_{13}$: C 55.75, H 6.10, N 5.65; found: C 55.76, H 6.09, N 5.67.


Minor diastereoisomer: $R_f = 0.28$ (hexane/ AcOEt 40:60); 1H NMR (400 MHz, $CDCl_3$), δ : 7.30-7.17 (m, 12H), 6.52 (br s, 1H), 6.02 (br s, 1H), 5.08 (s, 2H), 4.93 (s, 1H), 4.91 (s, 1H), 4.58-4.49 (m, 6H), 4.34 (q, $J = 7.2$ Hz, 1H), 4.23 (t, $J = 6.8$ Hz, 1H), 4.17 (m, 1H), 3.42 (m, 1H), 3.34 (s, 3H), 3.33 (s, 3H), 3.20-3.14 (m, 4H), 1.65-1.52 (m, 5H), 1.43 (s, 3H), 1.42 (s, 3H), 1.27 (s, 3H), 1.26 (s, 3H), 0.86 (d, $J = 6.8$ Hz, 3H), 0.86 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ : 172.3, 169.8, 167.3, 158.1, 137.3, 135.6, 129.1, 128.5, 128.4, 128.2, 128.0, 126.7, 123.2 (q, $J = 284.7$ Hz), 113.4, 112.4, 110.5, 110.1, 86.2, 85.4, 84.5, 81.9, 81.8, 66.7, 56.1, 56.0, 55.1, 51.1, 46.7 (q, $J = 27.2$ Hz), 44.0, 41.1, 36.3, 29.6, 26.5, 26.4, 24.9, 24.8, 24.7, 24.6, 22.7, 21.8 ; ESI (m/z) 1013.4 [$M^+ + Na$, (100)]; Anal. calcd. for $C_{46}H_{60}F_6N_4O_{13}$: C 55.75, H 6.10, N 5.65; found: C 55.75, H 6.11, N 5.66.

4b: Mixture of two diastereoisomers: $R_f = 0.18$ (hexane/ AcOEt 20:80); ^1H NMR (400 MHz, CDCl_3), major diastereoisomer δ : 5.58 (d, $J = 5.2$ Hz, 1H), 5.54 (d, $J = 4.8$ Hz, 1H), 5.30 (s, 1H), 5.28 (s, 1H), 5.23 (m, 1H), 4.80 (d, $J = 2.0$ Hz, 1H), 4.59 (m, 2H), 4.32 (m, 1H), 4.29 (m, 2H), 4.26 (m, 1H), 4.11 (m, 2H), 3.98 (m, 2H), 3.84 (m, 2H), 3.74 (m, 1H), 3.49 (m, 3H), 3.21 (m, 3H), 2.16 (s, 3H), 2.11 (s, 3H), 2.41 (s, 3H), 1.99 (s, 3H), 1.84 (m, 2H), 1.46 (s, 3H), 1.45 (s, 9H), 1.35 (s, 3H), 1.34 (s, 6H), 1.27 (s, 3H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 170.6, 169.9, 169.7, 168.2, 122.9 (q, $J = 280.7$ Hz), 109.5, 109.3, 108.5, 97.5, 96.3, 96.1, 71.5, 70.9, 70.7, 70.5, 70.4, 69.6, 69.1, 68.5, 66.7, 66.2, 65.9, 62.5, 46.3, 46.1, 45.8, 37.2, 29.3, 28.9, 26.0, 25.9, 25.8, 25.0, 24.96, 24.89, 24.3, 24.1, 20.8, 20.7, 20.6, 20.5; ESI (m/z) 1123.1 [$\text{M}^+ + \text{Na}$, (100)]; Anal. calcd. for $\text{C}_{47}\text{H}_{67}\text{F}_6\text{N}_3\text{O}_{22}$: C 49.52, H 5.92, N 3.69; found: C 49.50, H 5.94, N 3.70.

19: Mixture of two diastereoisomers: $R_f = 0.23$ ($\text{CHCl}_3/\text{CH}_3\text{OH}$ 95:5); ^1H NMR (400 MHz, CD_3OD), major diastereoisomer δ : 6.36 (br s, 1H), 6.34 (br s, 1H), 5.59 (s, 1H), 5.26 (br s, 1H), 5.16 (d, $J = 2.8$ Hz, 1H), 4.92 (s, 1H), 4.39 (m, 1H), 4.31 (m, 1H), 4.19 (m, 3H), 4.09 (m, 2H), 3.89 (m, 3H), 3.76 (m, 2H), 3.69 (m, 1H), 3.47-3.17 (m, 15H), 2.89 (m, 2H), 2.68 (m, 2H), 1.97 (m, 1H), 1.47-1.28 (eleven s, 102H); minor diastereoisomer δ : 6.36 (br s, 1H), 6.34 (br s, 1H), 5.58 (s, 1H), 5.26 (br s, 1H), 5.14 (d, $J = 2.6$ Hz, 1H), 4.92 (s, 1H), 4.39 (m, 1H), 4.31 (m, 1H), 4.19 (m, 3H), 4.09 (m, 2H), 3.89 (m, 3H), 3.76 (m, 2H), 3.69 (m, 1H), 3.47-3.17 (m, 15H), 2.89 (m, 2H), 2.68 (m, 2H), 1.97 (m, 1H), 1.47-1.28 (eleven s, 102H); ^{13}C NMR (100.6 MHz, CDCl_3) δ : 168.3, 168.0, 157.5, 157.1, 156.8, 156.3, 122.7 (q, $J = 282.3$ Hz), 122.3 (q, $J = 282.7$ Hz), 109.3, 109.2, 109.0, 99.3, 98.0, 96.5, 96.4, 96.2, 85.7, 81.2, 80.1, 79.3, 79.0, 78.0, 74.3, 74.1, 74.0, 73.2, 72.1, 72.0, 71.6, 71.2, 70.8, 70.7, 70.5, 70.4, 70.2, 67.5, 66.1, 55.6, 52.9, 51.0, 50.9, 50.8, 50.4, 45.9 (septet, $J = 27.2$ Hz), 41.4, 40.3, 39.6, 34.4, 34.1, 30.9, 30.8, 28.2, 28.0, 27.7, 27.5, 27.4, 27.3, 25.6, 25.2, 25.1, 24.9, 24.8, 24.0, 23.9, 23.2; ESI (m/z) 1846.0 [$\text{M}^+ + \text{Na}$, (71)]; Anal. calcd. for $\text{C}_{77}\text{H}_{129}\text{F}_6\text{N}_9\text{O}_{31}\text{S}$: C 50.73, H 7.13, N 6.92; found: C 50.70, H 7.15, N 6.89.








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av Id78.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 10/17/13 15:08:50
Method Copy of _01esquirew.MS

Operator
Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av Id78b.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 10/29/13 16:02:38
Method Copy of _01esquirew.MS

Operator
Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av 2510.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 10/29/13 15:59:05
Method Copy of _01esquirew.MS

Operator
Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av 2522.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volontero


Acquisition Date 11/15/13 15:39:40
Method Copy of _01esquirew.MS

Operator
Instrument

Walter Panzeri
esquire3000plus










-L.G.S. - Laboratorio Grandi Strumenti - Display Report


Analysis Name av_2511.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio

Acquisition Date 10/29/13 15:50:35
Method Copy(3) of _01esquirew.MS
Operator Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av 2587.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 03/24/14 16:17:51
Method Copy of _01esquirew.MS

Operator
Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report


Analysis Name av 2581a.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio

Acquisition Date 03/12/14 16:21:59
Method Copy of _01esquirew.MS


Operator
Instrument


Walter Panzeri
esquire3000plus








minor diast.









- L.G.S. - Laboratorio Grandi Strumenti - Display Report

Analysis Info

Analysis Name av cb8.d

Sample Name

Comment 1 mg/mL dil 1:100 MeOH

Richiedente: Volonterio

Acquisition Date 02/02/12 15:16:26

Method Copy of _01walt
07.09.11.MS


Operator


Instrument


Walter Panzeri

esquire3000plus


Acquisition Parameter


Ion Source Type ESI
Scan Begin 50 m/z
Capillary Exit 147.3 VoltMass Range Mode Std/Normal
Scan End 950 m/z
Skim 1 40.0 VoltIon Polarity Positive
Averages 5 Spectra
Trap Drive 64.3Alternating Ion Polarity off
Accumulation Time 75 μ s
Auto MS/MS off







major diast.











minor diast










-L.G.S. - Laboratorio Grandi Strumenti - Display Report


Analysis Name av 2623.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio

Acquisition Date 02/18/15 13:38:15
Method Copy of _01tmix_posneg
Im.MS


Operator
Instrument


Walter Panzeri
esquire3000plus





major diast







-L.G.S. - Laboratorio Grandi Strumenti - Display Report

Analysis Name av 2625b.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 03/06/15 15:19:38
Method Copy of _01tmix_posneg
Im.MS


Operator
Instrument
Walter Panzeri
esquire3000plus





minor diast.






-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av 2625c.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 03/06/15 15:25:43
Method Copy of _01tmix_posneg
Im.MS

Operator
Instrument

Walter Panzeri
esquire3000plus








-L.G.S. - Laboratorio Grandi Strumenti - Display Report



Analysis Name av 2528.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Volonterio


Acquisition Date 11/14/13 10:16:08
Method Copy of _01esquirew.MS


Operator
Instrument

Administrator
esquire3000plus











19





-L.G.S. - Laboratorio Grandi Strumenti - Display Report

Analysis Name av 2626 bis.d
Sample Name
Comment 1 mg/ml dil 1:100 MeOH
Richiedente: Fronza

Acquisition Date 03/13/15 10:28:03
Method Copy(3) of _01tmix_posneg
hm.MS
Operator Instrument
Walter Panzeri
esquire3000plus

