

# Communication

# Ligand-Enabled, Copper-Catalyzed Regio- and Stereoselective Synthesis of Trialkylsubstituted Alkenylboronates from Unactivated Internal Alkynes

Taisuke Itoh, Yohei Shimizu, and Motomu Kanai

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.6b04646 • Publication Date (Web): 07 Jun 2016

Downloaded from http://pubs.acs.org on June 7, 2016

# **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Ligand-Enabled, Copper-Catalyzed Regio- and Stereoselective Synthesis of Trialkylsubstituted Alkenylboronates from Unactivated Internal Alkynes

Taisuke Itoh,<sup>†</sup> Yohei Shimizu,<sup>†</sup> and Motomu Kanai<sup>\*,†,§</sup>

<sup>†</sup>Graduate School of Pharmaceutical Sciences, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan <sup>§</sup>ERATO, Japan Science Technology Agency, Kanai Life Science Catalysis Project, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Supporting Information Placeholder

**ABSTRACT:** We report the first copper-catalyzed regio- and stereoselective borylalkylation of dialkylsubstituted internal alkynes with bis(pinacolato)diboron and alkyl halides. A catalytically-generated borylcopper species containing a novel  $\pi$ accepting NHC ligand chemoselectively reacted with unactivated internal alkynes over alkyl halides. The intermediate alkenylcopper species subsequently reacted with alkyl halides, affording the desired products. The copper catalyst differentiated steric demands between the two aliphatic substituents on the C $\equiv$ C triple bond of the alkyne substrates to exhibit high regioselectivity from a wide range of alkyne/alkyl halide combinations. This method is useful for the straightforward synthesis of trialkylsubstituted alkenylboronates, i.e., versatile precursors for tetrasubstituted alkenes containing three or four different alkylsubstituents, which are difficult to synthesize by other methods.

Alkenylboronates are synthetically versatile building blocks. Despite many reports of the synthesis of alkenylboronates, regioand stereoselective synthesis of tetrasubstituted alkenylboronates, which are useful precursors for tetrasubstituted alkenes, is particularly challenging.<sup>1-8</sup> Suginome and coworkers reported seminal examples of the regio- and stereoselective synthesis of tetrasubstituted alkenylboronates through nickel- or palladium-catalyzed intramolecular boryl transfer to a  $C \equiv C$  triple bond using substrates containing a reactive boryl moiety (Scheme 1a).<sup>1</sup> Soon after, palladium- and nickel-catalyzed intermolecular carboboration of alkynes was reported (Scheme 1b).<sup>2,3,4</sup> More recently, copper-catalyzed borylalkylation of alkynes through three-component coupling with bis(pinacolato)diboron [(Bpin)<sub>2</sub>] and alkyl halides has emerged as a powerful method to furnish tetrasubstituted alkenylboronates (Scheme 1c).<sup>5,6</sup> Three main events proceeded in a well-ordered manner in this one-pot, three-component reaction: (1) generation of a borylcopper species from a copper alkoxide catalyst and (Bpin)<sub>2</sub>, (2) regioselective borylcupration of alkynes to afford alkenylcopper intermediate 1, and (3) alkylation of 1 with alkyl halides. To date, however, the substrate scopes of those metal-catalyzed reactions (Scheme 1b and Scheme 1c, previous works) have been predominantly limited to aromatic-substituted alkynes.

For regio- and stereoselective synthesis of tetrasubstituted alkenylboronates containing three alkyl groups, Nishihara developed a zirconium-mediated stepwise C–C bond formation to alkylsubstituted alkynylboronates (Scheme 1d,  $R^1 = alkyl$ ).<sup>8a,c</sup> This method requires multiple steps, however, including regioselective zirconacycle formation, selective protonolysis of an sp<sup>3</sup> carbon– metal bond, alkenyl iodide formation, and a Negishi coupling reaction.<sup>8,9</sup> Here we report the first straightforward synthesis of tetrasubstituted alkenylboronates containing three different alkyl groups by copper-catalyzed regio- and stereoselective borylalkylation of unactivated, dialkylsubstituted internal alkynes (Scheme 1c, this work).

# Scheme 1. Regio- and Stereoselective Synthesis of Tetrasubstituted Alkenylboronates

(a) Carboboration of alkynes via intramolecular boryl transfer

$$R^{1} CI \xrightarrow{B} O \xrightarrow{R^{2}SnBu_{3} \text{ or } R^{2}ZrCp_{2}CI} R^{2} \xrightarrow{R^{1}} N^{1}Pr_{2}$$

(b) Intermolecular carboboration of aryl alkynes

$$\begin{array}{c} \text{Ar} \\ R^{1} \\ R^{2} \text{Bpin} \end{array} \xrightarrow{R^{2} \text{Bpin}} \left[ \begin{array}{c} R^{2} - M \\ Ar \\ R^{1} \end{array} \right] \xrightarrow{R^{2}} \text{Ar} \xrightarrow{R^{2}} \text{Bpin} \\ R^{1} \end{array} \right]$$

(c) Copper-catalyzed three-component coupling



(d) Previous synthesis of trialkylsubstituted alkenylboronates

$$R^{1} \xrightarrow{\text{Cp}_{2}\text{Zr}} \xrightarrow{PrOH;} Pd \text{ cat.} \xrightarrow{R^{1}} Bpin$$
Bpin
$$R^{2} \xrightarrow{Pd \text{ cat.}} \xrightarrow{R^{2}\text{Znl}} Et \xrightarrow{R^{2}} Bpin$$

Two main hurdles must be overcome to achieve a coppercatalyzed three-component coupling reaction with dialkylsubstituted alkynes, (Bpin)<sub>2</sub>, and alkyl halides; chemo- and regioselectivity. The borylcopper species must chemoselectively react with alkynes in the presence of alkyl halides, but the differentiation between the two electrophiles is more difficult in the reaction using dialkylsubstituted unactivated alkynes than when using reactive aromatic-substituted alkynes. Moreover, regioselective borylcupration of electronically unbiased dialkylsubstituted internal alkynes is very difficult to achieve.<sup>10</sup> We envisioned that these two hurdles could be overcome by identifying a proper ligand for the copper catalyst, based on the previous development of liganddependent, copper-catalyzed regiodivergent borylalkylation,<sup>11</sup> borylamination,<sup>12</sup> and boryloxygenation<sup>13</sup> of alkenes.

 Table 1. Optimization of Copper-Catalyzed Regioselective

 Borylalkylation of 2a<sup>a</sup>

PnO	+ BnBr	Cu(I) cat. (10 m ligand (12 mol (Bpin) <sub>2</sub> (1.3 eq KO <i>t</i> Bu (1.5 eq	ol%) %) uiv) BnO uiv)	Bn Bpin 4ag
ыю	22.	DMF, rt		Spiin Spiin Bn
	2a 3g (3 equiv	)	BnO	5ag
entry	Cu(I) cat.	ligand	yield (%) <sup>b</sup>	r.r.
1	CuCl	dppe	3	2.5 : 1
2	CuCl	dppbz	4	1.5 : 1
3	CuCl	Xantphos	1	>20 : 1
4	IMesCuCl	-	15	7.9:1
5	IPrCuCl	-	10	2.6 : 1
6	<sup>NQ</sup> IMesCuCl	-	74 (72)	13 : 1
7	MeIMesCuCI	_	12	5.5 : 1
8 <sup>c</sup>	<sup>NQ</sup> IMesCuCl	_	100 (90)	10 : 1
A	r-N-Ar	Mes <sup>-N</sup> Y <sup>N-N</sup>	Nes O=	
	Cu Cl	Cu CI	Mes <sup></sup>	N N N N Nes
IMes IPrCu	CuCl: Ar = Mes JCl: Ar = 2,6- <i>i</i> Pr <sub>2</sub> -C <sub>6</sub> H	<sup>Me</sup> lMesCuC I <sub>3</sub>	l	Ċu ĊI
	(M	es = 2,4,6-Me <sub>3</sub> -C <sub>6</sub>	<sub>3</sub> H <sub>2</sub> ) <sup>NQ</sup>	IMesCuCl
<i>ac</i>	1			· ) G (1)

<sup>*a*</sup>General reaction conditions: **2a** (0.10 mmol), **3g** (3 equiv), Cu(I) catalyst (10 mol%), ligand (12 mol%), (Bpin)<sub>2</sub> (1.3 equiv), and KOtBu (1.5 equiv) in DMF (0.5 mL) at rt for 4 h. Yield and regioisomeric ratio (r.r. = **4ag** : **5ag**) were determined by <sup>1</sup>H NMR analysis of a crude mixture using 1,1,2,2-tetrachloroethane as an internal standard. <sup>*b*</sup>Combined yield of **4ag** and **5ag**. Combined isolated yield is shown in parenthesis (entries 6 and 8). <sup>*c*</sup>Butyl iodide (**3a**) was used instead of **3g**. Combined yield of **4aa** and **5aa**, and regioisomeric ratio (r.r. = **4aa** : **5aa**) are shown.

Thus, we began our study by focusing on the ligand effects in the reaction with dialkylsubstituted alkyne 2a, (Bpin)<sub>2</sub>, and benzyl bromide (3g) (Table 1). Phosphine ligands were ineffective for the reaction (entries 1-3). In contrast, an N-heterocyclic carbene (NHC)-copper catalyst, IMesCuCl, exhibited potential reactivity and regioselectivity (15% yield, r.r. = 7.9 : 1, entry 4). Use of a bulkier NHC-copper catalyst, IPrCuCl, decreased the regioselectivity (r.r. = 2.6 : 1, entry 5). This tendency to decrease regioselectivity is likely due to destabilization of the reaction pathway to generate intermediate 1 ( $R^1 = BnO(CH_2)_2$ ,  $R^2 = Me$ ), where the sterically more hindered carbon atom is attached to the copper atom when the ligand is sterically demanding.<sup>14</sup> The low yield in entries 1-5 was due to the lack of chemoselectivity: significant amounts (50-62% based on (Bpin)<sub>2</sub>) of BnBpin were generated as a side product through the reaction of borylcopper with benzyl bromide (3g), rather than with alkyne 2a. We hypothesized that a  $\pi$ -accepting NHC–copper catalyst would exhibit increased affinity toward the  $\pi$ -electrons of alkynes,<sup>15</sup> leading to chemoselective

promotion of the desired borylcupration of **2a**, rather than undesired borylation of **3g**. As expected, use of a  $\pi$ -accepting naphthoquinone-conjugated NHC–copper catalyst, <sup>NQ</sup>IMesCuCl,<sup>16</sup> dramatically improved both the yield and regioselectivity (74% yield, r.r. = 13 : 1, entry 6) while markedly suppressing the undesired BnBpin generation (18%). In contrast, use of a simple methylsubstituted NHC–copper catalyst, <sup>Me</sup>IMesCuCl, as a control provided almost the same yield of borylalkylation products (12%, entry 7) as IMesCuCl (entry 4). When a less reactive alkylating reagent, BuI (**3a**), was used instead of BnBr (**3g**), product **4aa** was obtained in quantitative yield with high regioselectivity under the optimized conditions using the <sup>NQ</sup>IMesCuCl catalyst (r.r. = 10 : 1, entry 8).

Table	2.	Scope	of	<b>Copper-Catalyzed</b>	Regioselective
Boryla	lkyla	tion of l	Dialk	ylsubstituted Intern	al Alkynes <sup>a</sup>



1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60 <sup>a</sup>General reaction conditions: **2** (0.20 mmol), **3** (3 equiv), <sup>NQ</sup>IMesCuCl (10 mol%), (Bpin)<sub>2</sub> (1.3 equiv), and KOtBu (1.5 equiv) in DMF (1.0 mL) at rt for 4 h. Combined isolated yield of **4** and **5** and regioisomeric ratio (**4** : **5**) are shown. Regioisomeric ratio was determined by <sup>1</sup>H NMR analysis of a crude mixture. <sup>b</sup>Regioisomeric ratio was determined after isolation. <sup>c</sup>Corresponding alkyl bromides were used instead of alkyl iodides.

With the optimized conditions in hand, we next examined the substrate scope of this three-component coupling reaction (Table 2). A non-functionalized simple alkyne, 2-octyne (2b), afforded a synthetically useful yield and regioselectivity (4ba, 74% yield, r.r. = 6.9: 1). A larger difference in the steric demand between two alkyl substituents on the C=C triple bond (2c) led to increased regioselectivity (4ca, 17 : 1). The reaction proceeded without any difficulties with 3-hexyne (2d) bearing two symmetric ethyl substituents (4da). Silyl ether (2e), ester (2f and 2g), and amide (2h) functionalities were tolerated, and the products were obtained in high yield and regioselectivity (4ea, 4fa, 4ga, and 4ha). In the cases of propargyl ethers 2i and 2j and propargyl amine 2l, the alkylation proceeded predominantly at the proximate carbons to the hetero atoms, independent of the steric bulkiness of the substituents on the C=C triple bonds (4ia, 4ja, and 4la). In homopropargyl ether 2k, however, the directing effect of the heteroatom was not operative (4ka, 1.7 : 1). For alkyl-aromatic-disubstituted internal alkyne 2m, the alkylation proceeded predominantly at the benzylic position (4ma). Terminal alkyne 2n afforded a trisubstituted alkenyl boronate 4na in high yield, albeit with moderate regioselectivity (3.2:1).

The scope of alkyl halides was also broad. Unactivated alkyl iodide 3b afforded 4ab in high yield and regioselectivity. The C=C double bond (3c) and ester (3d) functionalities in the alkylation reagents were compatible with the reaction conditions. The reaction using (iodomethyl)cyclopropane (3e) solely produced the desired borylalkylation product 4ae and no ring-opening product, indicating that the alkylation step from alkenylcopper species 1 proceeds through a non-radical process.<sup>17</sup> Using highly reactive methyl iodide (3f) as an alkylating reagent, 4af was obtained in high yield and regioselectivity when 3f was added dropwise over 1 h. For introduction of other activated alkyl groups, such as benzyl and ethoxycarbonylmethyl groups, use of the corresponding alkyl bromides produced a higher yield (4ag and 4ah). Sterically congested isobutyl iodide (3i) gave 4ai in moderate yield and regioselectivity. A secondary alkyl iodide, isopropyl iodide (3j), was considerably unreactive, and the desired product was not obtained.

To demonstrate that the products are versatile precursors to tetrasubstituted alkenes, we conducted Suzuki-Miyaura coupling of **4ia** with iodobenzene and 1-bromododecane (Scheme 2). Tetrasubstituted alkenes **6** and **7**, containing three and four different alkyl substituents, respectively, were stereoselectively obtained in high yield. Therefore, the two-step sequence, i.e., copper-catalyzed regio- and stereoselective borylalkylation, followed by Suzuki-Miyaura coupling, is a useful method for the straightforward synthesis of tetrasubstituted alkenes containing three or four different alkylsubstituents, which are difficult to synthesize by other methods.<sup>8a,18</sup>

#### Scheme 2. Synthesis of Tetrasubstituted Alkenes Containing Three or Four Alkyl Substituents



In conclusion, we developed the first straightforward catalytic method for regio- and stereoselective synthesis of trialkylsubstituted alkenylboronates, versatile precursors for tetrasubstituted alkenes containing three or four different alkyl groups. This group of alkenes comprises useful building blocks for organic synthesis, but their efficient and concise synthesis has remained challenging. The  $\pi$ -accepting ability of the <sup>NQ</sup>IMes ligand on the copper(I) catalyst was found to be essential for controlling the high regio- and chemoselectivity in the three-component, one-pot borylalkylation process of dialkylsubstituted internal alkynes with (Bpin)<sub>2</sub> and alkyl halides. Studies are in progress to elucidate the origin of the high selectivity exhibited by the  $\pi$ -accepting NHC ligand in copper(I) chemistry.

### ASSOCIATED CONTENT

#### **Supporting Information**

The Supporting Information is available free of charge via the Internet at http://pubs.acs.org.

Experimental details and characterization data (PDF)

#### **AUTHOR INFORMATION**

#### **Corresponding Author**

kanai@mol.f.u-tokyo.ac.jp

#### Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENT

This work was supported in part by ERATO from JST (MK) and Grant-in-Aid for Scientific Research (C) from JSPS (YS). TI thanks JSPS for the fellowship.

## REFERENCES

(1) (a) Suginome, M.; Yamamoto, A.; Murakami, M. J. Am. Chem. Soc. 2003, 125, 6358. (b) Yamamoto, A.; Suginome, M. J. Am. Chem. Soc. 2005, 127, 706. (c) Daini, M.; Yamamoto, A.; Suginome, M. J. Am. Chem. Soc. 2008, 130, 2918.

(2) (a) Suginome, M.; Yamamoto, A.; Murakami, M. Angew. Chem. Int. Ed. **2005**, 44, 2380. (b) Suginome, M.; Shirakura, M.; Yamamoto, A. J. Am. Chem. Soc. **2006**, 128, 14438. (c) Daini, M.; Suginome, M. Chem. Commun. **2008**, 5224. (d) Mannathan, S.; Jeganmohan, M.; Cheng, C.-H. Angew. Chem. Int. Ed. **2009**, 48, 2192.

(3) Iron-catalyzed alkylboration of symmetrical alkynes was reported: Nakagawa, N.; Hatakeyama, T.; Nakamura, M. *Chem. Eur. J.* **2015**, *21*, 4257.

(4) For a review of base metal-catalyzed borylation of alkynes, see: Yoshida, H. *ACS Catal.* **2016**, *6*, 1799.

(5) For copper-catalyzed borylalkylation of arylalkynes, see: (a) Akfaro, R.; Parra, A.; Alemán, J.; Luis, J.; Ruano, G.; Tortosa, M. J. Am. Chem. Soc. **2012**, *134*, 15165. (b) Yoshida, H.; Kageyuki, I.; Takaki, K.

*Org. Lett.* **2013**, *15*, 952. (c) Bidal, Y. D.; Lazreg, F.; Cazin, C. S. J. *ACS Catal.* **2014**, *4*, 1564. (d) Tai, C.-C.; Yu, M.-S.; Chen, Y.-L.; Chuang, W.-H.; Lin, T.-H.; Yap, G. P. A.; Ong, T.-G. *Chem. Commun.* **2014**, *50*, 4344. (e) Bin, H.-Y.; Wei, X.; Zi, J.; Zuo, Y.-J.; Wang, T.-C.; Zhong, C.-M. *ACS Catal* **2015**, *5*, 6670. For a pioneering work of copper-mediated borylalkylation of activated alkynes, see: (f) Okuno, Y.; Yamashita, M.; Nozaki, K. *Angew. Chem. Int. Ed.* **2011**, *50*, 920.

(6) For copper-catalyzed carboboration of arylalkynes, see: (a) Zhang, L.; Cheng, J.; Carry, B.; Hou, Z. J. Am. Chem. Soc. **2012**, *134*, 14314. (b) Zhou, Y.; You, W.; Smith, K. B.; Brown, M. K. Angew. Chem. Int. Ed. **2014**, *53*, 3475. (c) Lesieur, M.; Bidal, Y. D.; Lazreg, F.; Nahra, F.; Cazin, C. S. J. ChemCatChem **2015**, *7*, 2108.

(7) For other examples for catalyzed regio- and stereoselective synthesis of tetrasubstituted alkenylboronates, see: (1) Itami, K.; Kamei, T.; Yoshida, J.-I. *J. Am. Chem. Soc.* **2003**, *125*, 14670. (b) Shimizu, M.; Nakamaki, C.; Shimono, K.; Schelper, M.; Kurahashi, T.; Hiyama, T. *J. Am. Chem. Soc.* **2005**, *127*, 12506.

(8) (a) Nishihara, Y.; Okada, Y.; Jiao, J. Suetsugu, M.; Lan, M.-T.; Kinoshita, M.; Iwasaki, M.; Takagi, K. *Angew. Chem. Int. Ed.* **2011**, *50*, 8660. For the related stereoselective synthesis of tetrasubstituted alkenylboronates containing aromatic substituents, see: (b) Nishihara, Y.; Miyasaka, M.; Okamoto, M.; Takahashi, H.; Inoue, E.; Tanemura, K.; Takagi, K. *J. Am. Chem. Soc.* **2007**, *129*, 12634. (c) Nishihara, Y.; Suetsugu, M.; Saito, D.; Kinoshita, M.; Iwasaki, M. Org. Lett. **2013**, *15*, 2418.

(9) For another non-catalytic method, see: Endo, K.; Hirokami, M.; Shibata, T. J. Org. Chem. 2010, 75, 3469.

(10) For selected examples of regioselective addition to electronically unbiased internal alkynes, see: (a) Malik, H. A.; Sormunen, G. J.; Montgomery, J. J. Am. Chem. Soc. 2010, 132, 6304. (b) Liu, P.; Montgomery, J.; Houk, K. N. J. Am. Chem. Soc. 2011, 133, 6956. (c) Miller, K. M.; Jamison, T. F. J. Am. Chem. Soc. 2004, 126, 15342. (d) Liu, P. McCarren, P.; Cheong, P. H.-Y.; Jamison, T. F.; Houk, K. N. J. Am. Chem. Soc. 2010, 132, 2050. (e) Barchuk, A.; Ngai, M.-Y.; Krische, M. J. J. Am. Chem. Soc. 2007, 129, 8432. (f) Reichard, H. A.; McLaughlin, M.; Chen, M. Z.; Micalizio, G. C. Eur. J. Org. Chem. 2010, 391. (g) Chen, Q.-A.; Cruz, F. A.; Dong, V. M. J. Am. Chem. Soc. 2015, 137, 3157. (h) Liang, T.; Nguyen, K. D.; Zhang, W.; Krische, M. J. J. Am. Chem. Soc. 2015, 137, 3161. (i) Moure, A. L.; Arrayás, R. G.; Cárdenas, D. J.; Alonso, I.; Carretero, J. C. J. Am. Chem. Soc. 2012, 134, 7219.

(11) Su, W.; Gong, T.-J.; Lu, X.; Xu, M.-Y.; Yu, C.-G.; Xu, Z.-Y.; Yu, H.-Z.; Xiao, B.; Fu, Y. Angew. Chem. Int. Ed. 2015, 54, 12957.

(12) Sakae, R.; Hirano, K.; Miura, M. J. Am. Chem. Soc. 2015, 137, 6460.

(13) Itoh, T.; Matsueda, T.; Shimizu, Y.; Kanai, M. Chem. Eur. J. 2015, 21, 15955.

(14) Consistent with this explanation, regioselectivity decreased according to the decreased steric demand of boronates. When less bulky bis[(-)-pinanediolato]diboron and bis(hexylene glycolato)diboron were used instead of (Bpin)<sub>2</sub>, regioselectivity decreased to 5.9 : 1 and 1 : 1.5, respectively, using a <sup>NQ</sup>IMesCuCl catalyst (cf. 10 : 1 with (Bpin)<sub>2</sub>, see Table 1, entry 8).

(15)  $\pi$ -Accepting NHC- and cyclic (alkyl)(amino)carbene (CAAC)gold catalysts promoting addition reactions to alkenes and alkynes were reported. See: (a) López-Gómez, M. J.; Martin, D.; Bertrand, G. *Chem. Commun.* **2013**, 49, 4483. (b) Hu, X.; Martin, D.; Malaimi, M.; Bertrand, G. J. Am. Chem. Soc. **2014**, 136, 13594. (c) Lavallo, V.; Frey, G. D.; Donnadieu, B.; Soleilhavoup, M.; Bertrand, G. Angew. Chem. Int. Ed. **2008**, 47, 5224. (d) Kinjo, R.; Donnadieu, B.; Bertrand, G. Angew. Chem. Int. Ed. **2011**, 50, 5560. (e) Alcarazo, M.; Stork, T.; Anoop, A.; Thiel, W.; Fürstner, A. Angew. Chem. Int. Ed. **2010**, 49, 2542.

(16) The use of <sup>NQ</sup>IMesCuCl in organic synthesis is unprecedented. A <sup>NQ</sup>IMes-nickel complex was used in Kumada coupling reaction. See: (a) Sanderson, M. D.; Kamplain, J. W.; Bielawski, C. W. J. Am. Chem. Soc. **2006**, *128*, 16514. (b) Tennyson, A. G.; Lynch, V. M.; Bielawski, C. W. J. Am. Chem. Soc. **2010**, *132*, 9420.

(17) In contrast, the iron-catalyzed alkylboration provided a ringopening product. See ref 3.

(18) For a review of stereoselective synthesis of tetrasubstituted alkenes, see: Flynn, A. B.; Ogilvie, W. W. *Chem. Rev.* **2007**, *107*, 4698.

