This article was downloaded by: [University of Tennessee, Knoxville] On: 19 December 2014, At: 10:07 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

FORMATION OF 2'-DEOXY-2-NITROADENOSINES BY REACTION OF 2'-DEOXYADENOSINES WITH COPPER(II) NITRATE/ACETIC ANHYDRIDE

Toyo Kaiya^a, Haruko Tanaka^a & Kohfuku Kohda^b

^a Graduate School of Pharmaceutical Sciences , Nagoya City University , Tanabedori, Mizuhoku, Nagoya, 467-8603, Japan

^b Graduate School of Pharmaceutical Sciences, Nagoya City University, Tanabedori, Mizuhoku, Nagoya, 467-8603, Japan

Published online: 31 Aug 2006.

To cite this article: Toyo Kaiya , Haruko Tanaka & Kohfuku Kohda (2002) FORMATION OF 2'-DEOXY-2-NITROADENOSINES BY REACTION OF 2'-DEOXYADENOSINES WITH COPPER(II) NITRATE/ACETIC ANHYDRIDE, Nucleosides, Nucleotides and Nucleic Acids, 21:6-7, 427-433

To link to this article: http://dx.doi.org/10.1081/NCN-120014815

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS Vol. 21, Nos. 6 & 7, pp. 427–433, 2002

FORMATION OF 2'-DEOXY-2-NITROADENOSINES BY REACTION OF 2'-DEOXYADENOSINES WITH COPPER(II) NITRATE/ACETIC ANHYDRIDE

Toyo Kaiya, Haruko Tanaka, and Kohfuku Kohda*

Graduate School of Pharmaceutical Sciences, Nagoya City University, Tanabedori, Mizuho-ku, Nagoya, 467-8603, Japan

ABSTRACT

Nitration of 9-substituted [ethyl, $(Ac)_2$ -2'-deoxyribosyl, $(Ac)_3$ -ribosyl] N^6 -acetyladenine derivatives with Cu(NO₃)₂·3H₂O/Ac₂O was examined. Nitration proceeded at the 2-position, although the yield was low. Removal of the acetyl groups gave 2'-deoxy-2-nitroadenosine derivatives.

INTRODUCTION

Nitric oxide (NO) is involved in important physiological functions such as vasodilation^[1] and neurotransmission.^[2] However, it can have adverse effects: NO forms harmful peroxynitrite (ONOO⁻) by reaction with superoxide, and it has been reported that peroxynitrite reacts with the guanine moiety of DNA to form 8-nitroguanine and 8-oxoguanine,^[3] the latter of which is considered to be a DNA modification responsible for mutation/ cancer induction.^[4] The adenine moiety of DNA also forms 8-oxoadenine and 2-hydroxyadenine by treatment with oxidizing agents.^[5–6] Nitrated adenine derivatives, however, had not been described until a recent report by

427

DOI: 10.1081/NCN-120014815 Copyright © 2002 by Marcel Dekker, Inc. 1525-7770 (Print); 1532-2335 (Online) www.dekker.com

^{*}Corresponding author. Fax: +81-52-834-9309; E-mail: kohda@phar.nagoya-cu.ac.jp

Koomen's group who synthesized 2-nitroadenosine via 6-chloro-2-nitropurineriboside derivatives that were prepared by the nitration of 6-chloropurine derivative with tetrabutylammonium nitrate/trifluoroacetic anhydride (TBAN/TFAA).^[7] These investigators also obtained 2-nitroadenosine by nitration of $N^6, N^6, O^{2'}, O^{3'}, O^{5'}$ -pentaacetyladenosine with TBAN/TFAA and subsequent deacetylation, however, the yield was low due to a side reaction in the course of the deacetylation.^[7] Except for their study, direct nitration of adenine derivatives has been unreported, although many studies have been published on direct nitration of uracil derivatives using a variety of nitrating agents; $NO_2^+BF_4^-$, ^[8] NO_2OCOCF_3 , ^[9-10] $NO_2^-O_3$, ^[11] and $Cu(NO_3)_2 \cdot 3H_2O/$ Ac₂O.^[12] Sodum et al. reported quite recently on the reaction products of 2'deoxyadenosine with peroxynitrite. They showed that the main product was 2'-deoxy-8-oxoadenosine, however, the minor one was only suggested to be some nitrated derivative.^[13] In this study, we examined the preparation of nitrated 2'-deoxyadenosine derivatives. Since the glycosidic bond of 2'deoxyadenosine is less stable than that of adenosine against acid, milder conditions that cause low yield of nitration product are required for the preparation of 2'-deoxy-2-nitroadenosine. Reaction of $N^6, O^{3'}, O^{5'}$ -triacetyl-2'deoxyadenosine with Cu(NO₃)₂·3H₂O/Ac₂O afforded 2-nitro derivatives and subsequent removal of the acetyl groups gave 2'-deoxy-2-nitroadenosine.

RESULTS AND DISCUSSION

Nitration of 9-substituted N^6 -acetyladenine derivative (1) (Sch. 1) was carried out with Cu(NO₃)₂·3H₂O/Ac₂O. The reaction gave a small yield of



Scheme 1. Syntheses of 2'-deoxy-2-nitroadenosine derivatives.



Figure 1. HMBC correlations of compound 2a.

products (2 and 3) with unreacted starting material. The products were separated by preparative thin layer chromatography (PLC) (silica gel). The yield of 2 was slightly larger than that of 3 and the total yield of 2 and 3 was $7 \sim 33\%$. The mass spectrum of **2a** showed that it has a mono nitrated structure. The NMR spectrum of 2a suggested that the position of the nitro group is C2 or C8. The correct position of the nitro group was finally determined to be C2 by means of HMBC correlations (Fig. 1). From these results, the structure of 2a was determined to be N⁶-acetyl-9-ethyl-2-nitroadenine. Similarly, the structures of 2b,c were determined to be 2-nitroadenine derivatives. Deacetylation of 2a-c with NH₄OH/MeOH gave the corresponding 9-ethyl-2-nitroadenine (4a), 2'-deoxy-2-nitroadenosine (4b) and 2-nitroadenosine (4c), respectively. Compound 4 has a characteristic UV absorption spectrum with a λ max at a long wavelength due to the effect of the nitro group (Fig. 2). In the course of deacetylation, 2-methoxyadenine derivatives were obtained as a by-product by replacement of the nitro group of **4** with the methoxy group as reported.^[7] The yield of 2-methoxyadenine derivatives was less than 25%. Structures of 3a-c were determined to be N^6 -deacetylated **2a**-c from spectral data. Deacetylation of **3b**,c with NH₄OH/MeOH also gave 4b,c. When adenine derivatives without the N° -acetyl group were employed as substrate, no reaction proceeded within the Cu(NO₃)₂·3H₂O/Ac₂O system. This indicates that radical nitration mechanisms may be involved.^[7] It has been reported that 2'-deoxy-8-nitroguanosine is very unstable and decomposes to 8-nitroguanine within a few minutes.^[3,13] The stability of **4b** was then examined using HPLC. Compound **4b** was fairly stable and no decomposition was observed under treatment with aqueous solution (pH 6.8) at 60°C for 4 h. Acid treatment (0.1 N HCl at 60° C for 40 min) of **4b** gave 2-nitroadenine (5). When N⁶-acetyladenine was allowed to react with $Cu(NO_3)_2 \cdot 3H_2O/Ac_2O$, the reaction did not give N⁶acetyl-2-nitroadenine but gave other products (unstable ring-nitrogen-mononitrated N^6 -acetyladenine and stable N^6 -acetyl-2(8)-nitro-8(2)-oxoadenine) and their structures were deduced from NMR and mass spectroscopic data. When 9-ethylguanine was allowed to react with $Cu(NO_3)_2 \cdot 3H_2O/Ac_2O$, the reaction did not give a nitro derivative but gave instead N-ethylparabanic acid.



Figure 2. UV absorption spectrum of compound 4b at pH 6.8.

EXPERIMENTAL SECTION

Materials and Methods

¹H and ¹³C NMR spectra were recorded on a JEOL EX 270, GSX 400, or ALPHA 500 spectrometer, and chemical shifts were expressed in ppm using Me₄Si as the internal standard. EI and FAB mass spectra were obtained with a JEOL JMS-DX300 spectrometer. ESI mass spectra were obtained with a Micromass Quattro II spectrometer. HPLC analyses were carried out using a Shimadzu LC-10AD apparatus equipped with an SPD-M6A, a photodiode array UV detector. The Merck LiChrosphere 100 RP-18e, 4×250 mm column used was eluted at a flow rate of 0.6 mL/min with a 0.15 M phosphate buffer (pH 7.0)–MeOH system. Melting points were measured with a Yanagimoto micro melting point apparatus and are uncorrected. Silica gel 60 PF₂₅₄ (Merck) was used for preparative thin-layer chromatography (PLC).

Nitration of N^6 -Acetyladenine Derivatives (1) with Cu(NO₃)₂·3H₂O/Ac₂O

As a general practice, the reaction of **1c** which showed the highest yield was described. $Cu(NO_3)_2 \cdot 3H_2O$ (800 mg, 3.32 mmol) was dissolved in 6 mL of

2'-DEOXY-2-NITROADENOSINES

Ac₂O. After 15 min, **1c** (200 mg, 0.46 mmol) was added and the mixture was left at room temperature for 22 h. MeOH (80 mL) was then added and the mixture was evaporated to dryness. Water was added to the residue and the products were extracted with AcOEt. Separation and purification of the products were carried out by PLC (silica gel, AcOEt). The yields of **2c** and **3c** were 41 mg (19%) and 29 mg (14%), respectively, and **1c** was recovered in a 20 mg (10%) yield. In the case of **1b**, separation of the products was carried out by PLC (silica gel, AcOEt:MeOH = 40:1), and the yields of **2b** and **3b** were 3.4% and 3.3%, respectively, and the recovery of starting material was 11%. A similar procedure was employed for the reaction of **1a** and products **2a** and **3a** were separated by PLC (silica gel, AcOEt) in total yield 19%.

*N*⁶-Acetyl-9-ethyl-2-nitroadenine (2a). Colorless needles (MeOH-AcOEt), mp 194–196°C; ¹H NMR (500 MHz, DMSO-*d*₆): δ 1.49 (t, 3H, J = 7.3 Hz, CH₂CH₃), 2.32 (s, 3H, COCH₃), 4.34 (q, 2H, CH₂CH₃), 8.80 (s, 1H, 8-H), 11.28 (bs, 1H, NH); ¹³C NMR (DMSO-*d*₆) δ 14.8 (CH₂CH₃), 24.4 (COCH₃), 39.1 (CH₂CH₃), 125.2 (5-C), 147.5 (8-C), 149.6 (6-C), 151.8 (4-C), 153.6 (2-C), 168.9 (COCH₃); EI-MS: m/z 250 (M⁺), 208 (M⁺ – Ac + H), 162 (M⁺ – Ac + H – NO₂); HRMS calcd for C₉H₁₀N₆O₃: 250.0814, found 250.0814.

 $N^{6}, O^{3'}, O^{5'}$ -Triacetyl-2'-deoxy-2-nitroadenosine (2b). Colorless solid; ¹H NMR (CDCl₃): δ 2.10, 2.17 (each s, each 3H, 3'-, 5'-OCOCH₃), 2.77 (m, 1H, 2'-H), 2.78 (s, 3H, N-COCH₃), 2.89 (m, 1H, 2"-H), 4.42 (m, 3H, 4'-H, 5'-H, 5"-H), 5.45 (m, 1H, 3'-H), 6.55 (dd, 1H, J = 6.1, 7.6 Hz, 1'-H), 8.45 (s, 1H, 8-H), 8.94 (bs, 1H, NH); EI-MS: m/z 422 (M⁺); HRMS calcd for C₁₆H₁₈N₆O₈: 422.1185, found 422.1183.

 $N^{6}, O^{2'}, O^{3'}, O^{5'}$ -Tetraacetyl-2-nitroadenosine (2c). Colorless solid; ¹H NMR (CDCl₃): δ 2.10, 2.14, 2.20 (each s, each 3H, 2'-, 3'-, 5'-OCOCH₃), 2.79 (s, 3H, N-COCH₃), 4.47 (m, 2H, 4'-H, 5'-H), 4.53 (m, 1H, 5"-H), 5.62 (dd, 1H, J = 4.3, 5.5 Hz, 3'-H), 5.78 (m, 1H, 2'-H), 6.30 (d, J = 5.8 Hz, 1'-H), 8.41 (s, 1H, 8-H), 8.90 (bs, 1H, NH): EI-MS: m/z 480 (M⁺); HRMS calcd for C₁₈H₂₀N₆O₁₀: 480.1241, found 480.1241.

9-Ethyl-2-nitroadenine (3a = 4a). Yellow powder (H₂O), mp 280°C (decomp); ¹H NMR (DMSO- d_6): δ 1.44 (t, 3H, J = 7.3 Hz, CH₂CH₃), 4.23 (q, 2H, CH₂CH₃), 8.16 (bs, 2H, NH₂), 8.44 (s, 1H, 8-H); EI-MS: m/z 208 (M⁺); HRMS calcd for C₇H₈N₆O₂: 208.0708, found 208.0704; UV λ max (nm) (pH 6.8) 245, 268 (sh), 346.

 $O^{3'}$, $O^{5'}$ -Diacetyl-2'-deoxy-2-nitroadenosine (3b). Colorless solid; ¹H NMR (CDCl₃): δ 2.10, 2.16 (each s, each 3H, 3'-, 5'-OCOCH₃), 2.74 (m, 1H, 2'-H), 2.88 (m, 1H, 2"-H), 4.39 (m, 2H, 5'-H, 5"-H), 4.42 (m, 1H, H-4'), 5.44 (m, 1H, 3'-H), 6.48 (bs, 2H, NH₂), 6.49 (dd, 1H, J = 5.9, 7.6 Hz, 1'-H), 8.24 (s, 1H, 8-H); EI-MS: m/z 380 (M⁺); HRMS calcd for C₁₄H₁₆N₆O₇: 380.1082, found 380.1082.

 $O^{2'}, O^{3'}, O^{5'}$ -Triacetyl-2-nitroadenosine (3c). Colorless solid; ¹H NMR (CDCl₃): δ 2.10, 2.13, 2.18 (each s, each 3H, 2'-, 3'-, 5'-OCOCH₃), 4.47 (m, 2H, 4'-H, 5'-H), 4.50 (m, 1H, 5"-H), 5.64 (dd, 1H, J = 4.6, 5.5 Hz, 3'-H), 5.77 (m, 1H, 2'-H), 6.24 (d, J = 5.2 Hz, 1'-H), 6.27 (bs, 2H, NH₂), 8.17 (s, 1H, 8-H); EI-MS: m/z 438 (M⁺); HRMS calcd for C₁₆H₁₈N₆O₉: 438.1134, found 438.1137.

Preparation of 2-Nitroadenine Derivatives (4)

Compound 2a, 2b, 2c, 3b or 3c was treated with conc. $NH_4OH/MeOH$ (1:3) at 50°C for 30 min and the product was separated and purified by PLC. The yield of 4 was about 70%.

2'-Deoxy-2-nitroadenosine (4b). Pale yellow powder; ¹H NMR (D₂O): δ 2.60 (ddd, 1H, J = 3.7, 6.4, 14.0 Hz, 2'-H), 2.87 (m, 1H, 2"-H), 3.83 (dd, 1H, J = 4.6, 12.5 Hz, 5'-H), 3.88 (dd, 1H, J = 3.4, 12.5 Hz, 5"-H), 4.19 (m, 1H, 4'-H), 4.69 (dd, J = 3.4, 6.1 Hz, 3'-H), 6.50 (t, J = 6.7 Hz, 1'-H), 8.49 (s, 1H, 8-H); ESI-MS: positive ion mode, m/z 296.95 (M + H)⁺, negative ion mode, m/z 294.98 (M – H)⁻; UV: λ max (nm) (pH 6.8) 236, 262 (sh), 334.

2-Nitroadenosine (4c).^[7] Pale yellow powder; ¹H NMR (CD₃OD): δ 3.79, 3.90 (each m, each 1H, 5'-H, 5"-H), 4.15 (m, 1H, 4'-H), 4.36 (dd, 1H, J = 3.7, 4.9 Hz, 3'-H), 4.69 (m, 1H, 2'-H), 6.03 (d, 1H, J = 5.5 Hz, 1'-H), 8.53 (s, 1H, 8-H); ESI-MS: positive ion mode, m/z 312.91 (M + H)⁺, negative ion mode, m/z 310.98 (M – H)⁻; UV: λ max (nm) (pH 6.8) 236, 261 (sh), 334.

2-Methoxyadenosine.^[7] Yield <25%; ¹H NMR (DMSO-*d*₆): δ 3.53. 3.64 (each m, each 1H, 5'-H, 5"-H), 3.82 (s, 3H, CH₃), 3.93 (m, 1H, 4'-H), 4.15 (dd, 1H, *J* = 3.4, 4.9 Hz, 3'-H), 4.62 (m, 1H, 2'-H), 5.78 (d, *J* = 6.1 Hz, 1'-H), 7.31 (bs, 2H, NH₂), 8.14 (s, 1H, 8-H); FAB-MS *m*/*z* 298 (M + H)⁺.

Preparation of 2-Nitroadenine (5)

Acid hydrolysis of **4b** and **4c** was carried out in 0.1N HCl at 60° C for 40 min and in 1N HCl at 90°C for 1h, respectively. Yellow precipitates of **5** which appeared after cooling the reaction mixture in almost quantitative yield were collected.

2-Nitroadenine (5). mp >300°C; ¹H NMR (DMSO- d_6): δ 8.05 (bs, 2H, NH₂), 8.40 (s, 1H, 8-H); EI-MS m/z 180 (M⁺), 134 (M⁺ – NO₂); HRMS

2'-DEOXY-2-NITROADENOSINES

calcd for $C_5H_4N_6O_2$: 180.0396, found 180.0397. UV: λ max (nm) (pH 6.8) 235, 262 (sh), 280 (sh), 334.

Reaction of N^6 -Acetyladenine with Cu(NO₃)₂·3H₂O/Ac₂O

 N^6 -Acetyladenine was allowed to react with Cu(NO₃)₂·3H₂O/Ac₂O as described for the nitration of N^6 -acetyladenine derivatives (1). Products were separated by PLC (CHCl₃:MeOH = 9:1).

Ring-nitrogen-mono-nitrated adenine. Yield 3.1%; ¹H NMR (DMSOd₆): δ 7.73 (bs, 2H, NH₂), 8.33 and 8.97 (each s, each 1H, 2-H and 8-H); EI-MS: m/z 180 (M⁺); HRMS calcd for C₅H₄N₆O₂: 180.0395, found 180.0401. The position (1, 3, 7 or 9) of the N-nitro group could not be deduced.

*N*⁶-Acetyl-2(8)-nitro-8(2)-oxoadenine. Yield 2.1%; ¹H NMR (DMSOd₆): δ 2.15 (s, 3H, CH₃), 11.22 (bs, 1H, NH); ¹³C NMR (DMSO-d₆): δ 23.0 (CH₃), 115.0, 136.5, 152.0, 152.6, 152.8, 169.5; EI-MS: m/z 238 (M⁺), 196 (M⁺ - C₂H₂O), 150 (M⁺ - C₂H₂O - NO₂); HRMS calcd for C₇H₆N₆O₄: 238.0450, found 238.0455.

ACKNOWLEDGMENT

We thank Dr. R. Taguchi for ESI mass analyses.

REFERENCES

- 1. Loscalzo, J.; Welch, G. Prog. Cardiovasc. Dis. 1995, 38, 87.
- 2. Rand, M.J.; Li, C.G. Annu. Rev. Physiol. 1995, 57, 659.
- 3. Yermilov, V.; Rubio, J.; Ohsima, H. FEBS Lett. 1995, 376, 207.
- Cheng, K.C.; Cahill, D.S.; Kasai, H.; Nishimura, S.; Loeb, L.A. J. Biol. Chem. 1992, 267, 166.
- Murata-Kamiya, N.; Kamiya, H.; Muraoka, M.; Kaji, H.; Kasai, H. J. Radiat. Res. 1997, 38, 121.
- 6. Kamiya, H.; Kasai, H. J. Biol. Chem. 1995, 270, 19446.
- 7. Deghati, P.Y.; Wanner, M.J.; Koomen, G.-J. Tetrahedron Lett. 2000, 41, 1291.
- 8. Huang, G.-F. (Kohda, K.); Torrence, P.F.J. Org. Chem. 1977, 42, 3821.
- 9. Ariza, X.; Bou, V.; Vilarrasa, J. J. Am. Chem. Soc. 1995, 117, 3665.
- 10. Ariza, X.; Farras, J.; Serra, C.; Vilarrasa, J. J. Org. Chem. 1997, 62, 1547.
- 11. Suzuki, H.; Nonoyama, N. J. Chem. Research (S). 1996, 244.
- 12. Giziewicz, J.; Wnuk, S.T.; Robins, M.J. J. Org. Chem. 1999, 64, 2149.
- 13. Sodum, R.S.; Fiala, E.S. Chem. Res. Toxicol. 2001, 14, 438.

Received March 11, 2002 Accepted May 28, 2002