# organic compounds

Acta Crystallographica Section C Crystal Structure Communications ISSN 0108-2701

# C—H··· $\pi$ and $\pi$ - $\pi$ interactions in the supramolecular structure of 3-methyl-1,4-diphenyl-1*H*-pyrazolo[3,4-*b*]pyridine

## John N. Low,<sup>a</sup>\* Justo Cobo,<sup>b</sup> Manuel Nogueras,<sup>b</sup> Adolfo Sánchez,<sup>b</sup> Harlen Torres<sup>c</sup> and Braulio Insuasty<sup>c</sup>

<sup>a</sup>Department of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, <sup>b</sup>Departamento de Química Inorgánica y Orgánica, Universidad de Jaén, 23071 Jaén, Spain, and <sup>c</sup>Grupo de Investigación de Compuestos Heterocíclicos, Departamento de Química, Universidad del Valle, AA25360 Cali, Colombia Correspondence e-mail: jnlow111@hotmail.com

Received 4 April 2002 Accepted 9 April 2002 Online 30 April 2002

The supramolecular structure of the title compound,  $C_{19}H_{15}N_3$ , is defined by  $\pi$ - $\pi$ -stacking and C-H··· $\pi$  interactions. There are no conventional hydrogen bonds in the structure.

#### Comment

Previously, we have reported different syntheses of biologically active compounds with a pyrazolo[3,4-b]pyridine skeleton. 5-Aminopyrazoles were used as starting materials with  $\alpha,\beta$ -unsaturated ketones, as shown in the Scheme (Quiroga, Insuasty, Cruz et al., 1998; Quiroga, Cruz et al., 2001), with benzaldehydes and Meldrum's acid (Quiroga, Hormaza et al., 1998; Quiroga, Insuasty, Hormaza et al., 1998) or with dimedone (Quiroga, Mejía et al., 2001) as reactants. We now report the crystal structure of fully aromatized 3-methyl-1,4-diphenyl-1*H*-pyrazolo[3,4-b]pyridine, (I), which was prepared to test the synthetic application of 5-(3-aryl-2-propenylidene)-2,2-dimethyl-1,3-dioxane-4,6-diones [the structures of some examples have been reported recently; see, for example, Low et al. (2002)]. Reaction of the phenyl derivative with 5-amino-3-methyl-1-phenylpyrazole yielded (I). In this reaction, the Meldrum's acid fragment was completely eliminated, resulting in a fully aromatized product.

The pyrazole and pyridine rings in (I) are planar within experimental error and have a dihedral angle between them of  $5.13 (10)^{\circ}$ . The only other compound with a similar pyrazolopyridine structure reported in the Cambridge Structural Database (CSD; Version 5.22 of October 2001; Allen & Kennard, 1993) is 6-amino-5-cyano-3-methyl-1,4-diphenylpyrazolo[3,4-*b*]pyridine (Quiroga *et al.*, 1999), (II), which crystallizes in space group  $P2_1/n$ . In (II), the dihedral angle between the pyrazole and pyridine rings is 3.6 (3)°. N.B. The atomic numbering differs for (I) and (II); all values are quoted with respect to the numbering of (I).



The torsion angles for the phenyl groups in (I) are given in Table 1. As can be seen from the examples given below, the conformations of the phenyl rings in compounds (I) and (II) are very similar, for example, in the case of ring 4 (atoms C41–C46) [values for (II) are given in parenthesis],  $C5-C4-C41-C42 = 112.8 (2)^{\circ}$  [118.2 (5)°], but differ by nearly 23° in the case of phenyl ring 3 (atoms C11–C16), where C7a–N1–C11–C12 = 135.2 (2)° [158.0 (5)°].

In (I), the  $\pi$ - $\pi$  stacking involves the pyridine ring of the pyrazolo-pyridine group. The ring in the molecule at (x, y, z) stacks above the ring at (-x, 2 - y, 1 - z) across the centre of inversion at  $(0, 1, \frac{1}{2})$ , with a distance of 3.563 (2) Å between the ring centroids, a perpendicular distance between the rings of 3.443 (2) Å and a centroid offset of 0.913 (2) Å (Fig. 2a). Although compound (II) does contain a conventional N-



#### Figure 1

View of (I) with the atomic numbering scheme. Displacement ellipsoids are plotted at the 30% probability level.

H···N hydrogen bond, there is similar  $\pi - \pi$  stacking between the pyridine rings related by the centre of inversion at (0,1,1), with a  $Cg \cdot \cdot Cg$  distance of 3.701 (3) Å, a perpendicular distance of 3.552 Å and an offset between the centres of 1.04 Å. Fig. 2(*b*) shows the stacking in (II), labelled in this case with the numbering system used for (I). Figs. 2(*a*) and 2(*b*) show the differences in the stacking in the two compounds. These differences result from the different polarizations of the pyridine rings in the two compounds, which result from the presence of the amino and cyano substituents in compound (II).

There are two C-H··· $\pi$  interactions involving each of phenyl rings 3 (atoms C11-C16, centroid *Cg*3) and 4 (atoms C41-C46, centroid *Cg*4). The C43-H43···*Cg*3<sup>ii</sup> interaction (Table 2) links the molecules into centrosymmetrically related dimers *via* the centre of inversion at  $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$  [symmetry code:



View of the  $\pi$ - $\pi$  stacking between the molecules in (a) (I) and (b) (II).





View of the linked dimer chains and sheet formed by the  $\pi$ - $\pi$  stacking in (I).

(ii) 1 - x, 1 - y, 1 - z]. These dimers are then linked by a C13-H13...Cg4<sup>i</sup> interaction and the equivalent interaction *via* the centre at  $(\frac{1}{2}, \frac{1}{2}, 0)$ , so forming chains of dimers running parallel to [001] [symmetry code: (i) x, y, -1 + z]. These C-H... $\pi$  and  $\pi$ - $\pi$  interactions link the molecules into two-dimensional sheets lying parallel to (110) (Fig. 3). In (II), the equivalent ring to ring 3 is involved in a C-H... $\pi$  interaction. In this case, the atom equivalent to H42 is involved in an interaction with the ring in the molecule at  $(\frac{1}{2} - x, \frac{1}{2} + y, \frac{3}{2} - z)$ , with an H...Cg distance of 2.85 Å, a C...Cg distance of 3.735 (5) Å and an angle at H of 160°.

### **Experimental**

An ethanol solution containing equimolar quantities of 5-(3-phenyl-2-propenylidene)-2,2-dimethyl-[1,3]dioxane-4,6-dione and 5-amino-3-methyl-1-phenylpyrazole in ethanol was heated under reflux for

2 h. The solvent was then removed under vacuum and the resulting crude solid was purified by silica-gel column chromatography using hexane/ethyl acetate (1:1  $\nu/\nu$ ) as eluant, yielding (I) (55%, m.p. 408 K). Analysis, found: C 79.9, H 5.2, N 14.7%; C<sub>19</sub> H<sub>15</sub>N<sub>3</sub> requires: C 80.0, H 5.3, N 14.7%. Crystals suitable for single-crystal X-ray diffraction analysis were obtained from a chloroform solution.

Z = 2

 $D_x = 1.319 \text{ Mg m}^{-3}$ 

Cell parameters from 3135

 $0.20 \times 0.10 \times 0.10$  mm

Mo K $\alpha$  radiation

reflections

 $\mu=0.08~\mathrm{mm}^{-1}$ 

T = 120(1) K

Block, yellow

 $\theta = 3.1 - 27.4^{\circ}$ 

#### Crystal data

 $\begin{array}{l} C_{19}H_{15}N_3 \\ M_r = 285.34 \\ Triclinic, P\overline{1} \\ a = 8.1392 \ (4) \ \mathring{A} \\ b = 8.4351 \ (4) \ \mathring{A} \\ c = 11.3852 \ (6) \ \mathring{A} \\ a = 77.951 \ (3)^\circ \\ \beta = 81.192 \ (2)^\circ \\ \gamma = 70.681 \ (2)^\circ \\ V = 718.39 \ (6) \ \mathring{A}^3 \end{array}$ 

#### Data collection

Nonius KappaCCD diffractometer<br/> $\varphi$  scans, and  $\omega$  scans with  $\kappa$  offsets1682 reflections with  $I > 2\sigma(I)$ <br/> $R_{int} = 0.092$ Absorption correction: multi-scan<br/>(DENZO-SMN; Otwinowski &<br/>Minor, 1997) $\theta_{max} = 27.4^{\circ}$ <br/> $h = -10 \rightarrow 10$ <br/> $k = -10 \rightarrow 10$ <br/> $l = -14 \rightarrow 14$ 9243 measured reflections<br/>3135 independent reflectionsIntensity decay: negligible

#### Refinement

Refinement on $F^2$	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.053$	$w = 1/[\sigma^2(F_o^2) + (0.0455P)^2]$
$wR(F^2) = 0.123$	where $P = (F_o^2 + 2F_c^2)/3$
S = 0.94	$(\Delta/\sigma)_{\rm max} < 0.001$
3135 reflections	$\Delta \rho_{\rm max} = 0.23 \ {\rm e} \ {\rm \AA}^{-3}$
200 parameters	$\Delta \rho_{\rm min} = -0.35 \ {\rm e} \ {\rm \AA}^{-3}$

#### Table 1

Selected torsion angles ( $^{\circ}$ ).

C7a-N1-C11-C12	135.2 (2)	C5-C4-C41-C46	-66.5(3)
N2-N1-C11-C12	-39.0(3)	C3a-C4-C41-C46	111.6 (2)
C7a-N1-C11-C16	-45.1(3)	C5-C4-C41-C42	112.8 (2)
N2-N1-C11-C16	140.78 (18)	C3a-C4-C41-C42	-69.1(3)

#### Table 2

Hydrogen-bonding geometry (Å, °).

Cg3 and Cg4 are the centroids of the C11–C16 and C41–C46 rings, respectively.

$D-\mathrm{H}\cdots A$	$D-\mathrm{H}$	$H \cdot \cdot \cdot A$	$D \cdots A$	$D - \mathbf{H} \cdot \cdot \cdot A$
$\begin{array}{c} \text{C13}-\text{H13}\cdots\text{Cg4}^{\text{i}}\\ \text{C43}-\text{H43}\cdots\text{Cg3}^{\text{ii}} \end{array}$	0.95	2.74	3.543 (3)	142
	0.95	2.79	3.579 (3)	142

Symmetry codes: (i) x, y, z - 1; (ii) 1 - x, 1 - y, 1 - z.

Compound (I) crystallized in the triclinic system; space group  $P\overline{1}$  was assumed and confirmed by the analysis. H atoms were treated as riding, with C–H distances of 0.95 (aromatic) and 0.98 Å (methyl).

Data collection: *KappaCCD Server Software* (Nonius, 1997); cell refinement: *DENZO–SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO–SMN*; program(s) used to solve structure: *SHELXS*97 (Sheldrick, 1997); program(s) used to refine structure: *SHELXL*97 (Sheldrick, 1997); molecular graphics: *ORTEPII* (Johnson, 1976) and *PLATON* (Spek, 2002); software used to prepare material for publication: *SHELXL*97 and WordPerfect macro *PRPKAPPA* (Ferguson, 1999).

X-ray data were collected at the EPSRC, X-ray Crystallographic Service, University of Southampton. The authors thank the staff for all their help and advice. JNL thanks NCR Self-Service, Dundee, for grants which have provided computing facilities for this work. MN, AS and JC thank the 'Ministerio de Educación, Cultura y Deportes (Programa de Cooperación con Iberoamerica, AECI)' of Spain for financial support of part of this work

Supplementary data for this paper are available from the IUCr electronic archives (Reference: GD1203). Services for accessing these data are described at the back of the journal.

#### References

Allen, F. H. & Kennard, O. (1993). Chem. Des. Autom. News, 8, 1, 31-37.

- Ferguson, G. (1999). PRPKAPPA. University of Guelph, Canada.
- Johnson, C. K. (1976). ORTEPII. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Low, J. N., Cobo, J., Nogueras, M., Sánchez, A., Insuasty, B. & Torres, H. (2002). Acta Cryst. C58, 039-041.
- Nonius (1997). *KappaCCD Server Software*. Windows 3.11 Version. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). Methods in Enzymology, Vol. 276, Macromolecular Crystallography, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Quiroga, J., Alvarado, M., Insuasty, B., Moreno, R., Ravina, E., Estevez, I. & de Almedia, R. H. (1999). J. Heterocycl. Chem. 36, 1311–1316.
- Quiroga, J., Cruz, S., Insuasty, B., Abonía, R., Cobo, J., Sánchez, A., Nogueras, M. & Low, J. N. (2001). J. Heterocycl. Chem. 38, 53–60.
- Quiroga, J., Hormaza, A., Insuasty, B. & Márquez, M. (1998). J. Heterocycl. Chem. 35, 409–412.
- Quiroga, J., Insuasty, B., Cruz, S., Hernández, P., Bolaños, A., Moreno, R., Hormaza, A. & de Almedia, R. H. (1998). J. Heterocycl. Chem. 35, 333–338.
- Quiroga, J., Insuasty, B., Hormaza, A., Cabildo, P., Claramunt, R. M. & Elguero, J. (1998). J. Heterocycl. Commun. 5, 115–122.
- Quiroga, J., Mejía, D., Insuasty, B., Abonía, R., Nogueras, M., Sánchez, A. & Cobo, J. (2001). *Tetrahedron*, 57, 6947–6953.
- Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.
- Spek, A. L. (2002). PLATON. March 2002 Version. University of Utrecht, The Netherlands.