

Copper-Catalyzed Intermolecular Trifluoromethylazidation and Trifluoromethylthiocyanation of Allenes: Efficient Access to CF₃-Containing Allyl Azides and Thiocyanates


Na Zhu,[†] Fei Wang,[‡] Pinhong Chen,[‡] Jinxing Ye,^{*,†} and Guosheng Liu^{*,‡}

[†]Engineering Research Centre of Pharmaceutical Process Chemistry, Ministry of Education, School of Pharmacy, East China University of Science and Technology, 130 Meilong Road, Shanghai, China 200237

[‡]State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai, China 200032

S Supporting Information

ABSTRACT: A mild and efficient method for copper-catalyzed trifluoromethylazidation and trifluoromethylthiocyanation of allenes was explored. A series of CF₃-containing allyl azides and thiocyanates were obtained with high yields and good stereoselectivities, which can be used for further transformation to some valuable compounds.



Allyl amines and sulfides are versatile building blocks for the synthesis of organic molecules of higher complexity.¹ These structural motifs are frequently found in natural products and pharmaceuticals and commonly exhibit biological activities.² Generally, the introduction of a trifluoromethyl group into lead compounds was considered to modify their activity and biocompatibility.³ Thus, exploration of new methods for introducing the CF₃ group into organic compounds has been of broad interest.⁴ Some CF₃-substituted allyl amines and sulfides have been recognized to be biologically active; some of which are listed in Figure 1.⁵ However, traditional strategies for the




Figure 1. Representative bioactive compounds with CF₃-allylic amines and sulfides.

synthesis of these skeletons suffered from multistep transformations and low efficiency.⁵ Thus, new methods for constructing CF₃-containing allyl amines and sulfides are in high demand. Herein, we report a novel copper-catalyzed intermolecular trifluoromethylazidation and -thiocyanation of allenes in which azide or thiocyanate and CF₃ group are incorporated simultaneously; a series of CF₃-containing allyl azides and thiocyanates are efficiently synthesized.


In the past decade, copper-catalyzed trifluoromethylation of alkenes has been well documented, and an array of new chemical bonds were incorporated, such as C–X, C–O, C–N, C–S, and C–C bonds, depending on the nature of the functional group used to trap the key intermediate.^{6,7} The related trifluoromethylation of alkynes has also been achieved to synthesize various CF₃-containing vinyl products.⁸ In contrast, the trifluoromethylation of allenes is relatively rare.⁹ In 2013, Ma and Yu disclosed a copper-catalyzed intramolecular trifluoromethyloxylation of 2,3-allenoic acids to provide CF₃-substituted butenolides.^{9b} Later, an intermolecular trifluoromethyloxylation reaction was explored by Liu and co-workers, while the substrate scope was limited to heteroatom substituted allenes.^{9c} Recently, our group developed a series of copper-catalyzed intermolecular difunctionalization of alkenes¹⁰ and alkynes,¹¹ including trifluoromethylated azidation, arylation, cyanation, and thiocyanation, in which a mutual activation model between ether-type Togni's CF₃⁺ reagent and TMSNu or ArB(OH)₂ was presented to be vital for the success of these reactions (Scheme 1a). With our continuous efforts on this catalytic system, we speculated that if allenes were subjected to our reaction systems, a battery of functional groups, such as azide and thiocyanate, would simultaneously be easily introduced into the products with CF₃ group. However, the reaction of allenes presented more complex reactivity than that of alkenes because of more reaction sites of allene moiety, which could deliver more possible trifluoromethylation products (Scheme 1b). The interesting but questionable selectivity motivated us to investigate this chemistry.

To test our hypothesis, the initial investigation was focused on the reaction of substrate **1a**, which was treated with the


Received: June 8, 2015

Scheme 1. Copper-Catalyzed Intermolecular Trifluoromethylations

a) Reaction of alkenes and alkynes (previous work)



b) Reaction of allenes (this work)



previous reaction conditions of alkenes.^{10a} We were delighted to find the desired CF₃-substituted allyl azide product **2a** was detected in 17% yield with excellent regioselectivity but poor stereoselectivity (Table 1, entry 1). Further optimization of the


Table 1. Optimization of the Reaction Conditions^a

entry	solvent	copper	yield (%) ^b	Z/E ^b
1	DMAc	Cu(CH ₃ CN) ₄ PF ₆	17	1/1
2	MeOH	Cu(CH ₃ CN) ₄ PF ₆	65	1/13
3	CH ₃ CN	Cu(CH ₃ CN) ₄ PF ₆	48	1/23
4	THF	Cu(CH ₃ CN) ₄ PF ₆	30	1/9
5	DCM	Cu(CH ₃ CN) ₄ PF ₆	38	1/22
6	MeOH	Cu(OTf) ₂	35	1/11
7	MeOH	Cu(OAc) ₂	18	1/17
8	MeOH	CuI	16	1/14
9	MeOH	CuCl ₂	50	1/9
10	MeOH	CuOAc	35	1/16
11 ^c	MeOH	Cu(CH ₃ CN) ₄ PF ₆	47	1/17
12 ^d	MeOH	Cu(CH ₃ CN) ₄ PF ₆	44	1/15
13 ^e	MeOH	Cu(CH ₃ CN) ₄ PF ₆	48	1/10
14 ^f	MeOH	Cu(CH ₃ CN) ₄ PF ₆	42	1/8

^aAll the reactions were run at 0.1 mmol scale. ^b¹⁹F-NMR yield with CF₃-DMA as an internal standard; E/Z ratio was detected from the crude product. ^cEster-type Togni's CF₃⁺ reagent was used. ^dAt 0 °C. ^eAt 40 °C. ^fAt 60 °C.

solvent revealed that MeOH provided the best result, giving the corresponding product **2a** in 65% yield with excellent regio- and stereoselectivity (entries 2–5). Later, various Cu(I) and Cu(II) salts were also examined, and Cu(CH₃CN)₄PF₆ was determined to be the best. When ester-type Togni's CF₃⁺ reagent was used, the reaction also provided product **2a** but with a slightly lower yield (entry 11). In addition, under the lower or elevated temperature, the reactions suffered decreased yields (entries 12–14). It should be noted that the stereoselectivity of the reactions varied with the change of reaction conditions.


With the optimized reaction conditions in hand, the substrate scope of the reaction was examined, and the results are summarized in Scheme 2. Various 1,1-disubstituted aryl allenes **1** were suitable for the reaction to yield the corresponding trifluoromethylated allylazides in moderate to good yields with

Scheme 2. Substrate Scope for Trifluoromethylazidation of Allenes^{a,b}

^aAll the reactions were run at 0.2 mmol scale. ^bIsolated yield. ^cEster-type Togni's CF₃⁺ reagent.

good *E*-isomer selectivity. Among them, a variety of functional groups, such as Br, F, alkyl, methoxyl, were compatible with the reaction conditions. Allene substrates with linear *n*-pentyl, *n*-butyl, and methoxypropyl groups generated the desired products in good yields with good regio- and stereoselectivities. However, substrates **1d** and **1j** with steric bulky groups provided products **2d** and **2j** with poor stereoselectivities around a 2:1 ratio. For the aryl group, both electron-rich and electron-poor are suitable for the transformation, furnishing the corresponding products in good yields and stereoselectivities. In contrast, substrate **1g** with an *ortho*-substituent delivered product **2g** in a slightly lower stereoselectivity. These observations revealed that the stereoselectivity was sensitive to the steric hindrance of both alkyl and aryl groups. Unfortunately, quite limited substrate scope was observed: the reaction of monosubstituted, 1,3-disubstituted allenes provided the corresponding products in low yields with poor regioselectivities. In addition, 1,1-dialkyl substituted allenes also exhibited unsatisfactory reactivity toward this reaction.

Inspired by the present results, we turned our attention to extending such a system to the trifluoromethylthiocyanation reaction. To our great delight, when substrate **1h** was used as a template by treatment with TMSNCS instead of TMSN₃ under the above system, the desired product **3c** was detected in 60% yield. Further optimization demonstrated that the best yield was obtained in DMSO at 40 °C (see the Supporting Information). Next, the scope of the trifluoromethylthiocyanation reaction was evaluated under the modified conditions, and the results are listed in Scheme 3. Generally, similar to the previous transformation, this reaction furnished the desired products in good to excellent yields, but with poor stereoselectivities. Again, the reactions provided *E*-isomer as the main products. In contrast, when MeOH was used as the solvent, the reactions provided a better stereoselectivity, but


Scheme 3. Substrate Scope for the Trifluoromethylthiocyanation Reaction^{a,b}

^aAll the reactions were run at 0.2 mmol scale. ^bIsolated yield. ^cMeOH as solvent, at rt.

with slightly lower yields. Finally, the reactions exhibited good functional group compatibility, such as Br, F, and alkyl, under the reaction conditions.

To demonstrate the practicability of the present methods, further transformation of **2a** and **3a** were investigated (Scheme 4). We found that **2a** could be readily converted to the


Scheme 4. Further Transformations of Products



corresponding allylamine **4a** in 89% yield in the presence of PPh₃ and H₂O. In addition, treatment of **2a** and phenylacetylene with Cu catalyst delivered the triazole **5a** in 95% yield.^{10a} Furthermore, allylthiocyanate **3a** could react with TMSCF₃ through nucleophilic trifluoromethylation to generate related SCF₃-substituted product **6a** in 65% yield. Very interestingly, the reaction of **3a** and NaN₃ resulted in the corresponding product **7a** in 74% yield.^{10d} Notably, these CF₃-decorated *tetra*-substituted olefins were synthesized efficiently from simple allenes, which was difficult to achieve through previously reported methods.

Based on our previous reports on the mutual activation model,¹⁰ the proposed mechanism is provided in Scheme 5: the activated CF₃⁺ reagent by TMSNu could react with Cu(I) catalyst to release Cu(II), Nu[−], and CF₃ radical species, which was then trapped by allenes, giving a CF₃-allyl radical species with high regioselectivity.¹² For the further process of CF₃-allyl radical with Cu(II)/Nu[−], two scenarios could be used to address final C–N and C–S bonds formation: (1) The N₃ or SCN radical species might be generated from the oxidation of Nu[−] with Cu(II) then coupled with CF₃-allyl radical species to


Scheme 5. Plausible Mechanism



give final products (path b). (2) CF₃-allyl radical species combined with Cu(II)/Nu[−] to give a CF₃-allyl-Cu(III) species, which underwent reductive elimination to furnish the final products (path a). Both processes seem to be reasonable; however, at present, it is difficult to draw a conclusion.

We assumed that the addition of CF₃ radical to allene originally generated the corresponding two allyl radical intermediates (**int.I** and **int.II**) with poor selectivity. Then, species **int.II** would isomerize to **int.I**, which is relatively thermodynamically stable, because of the steric hindrance between the aryl and CF₃ group. For the trifluoromethylazidation, good stereoselectivity was observed when R is a less bulky group, while poor selectivity was provided when a sterically hindered alkyl group was incorporated (such as **2d** and **2j**). We reasoned that the poor selectivity may result from the steric effect between R and CF₃ groups, which pushed the equilibrium from **int.I** to **int.II** (Scheme 5). Compared to the azidation, the related thiocyanation reaction was obtained with poor stereoselectivities (around 3:1). Although a detailed mechanistic understanding cannot currently be given to address these differences, we speculated that trapping CF₃-allyl radical intermediates by heteroatom radicals or Cu(II) species might be responsible for the selectivity. When the coupling reaction of CF₃-allyl radical is faster than its isomerization (between **int.I** and **int.II**), poor stereoselectivity might be expected. Otherwise, high stereoselectivity will be observed.

With this in mind, we hold that the coupling reaction of Cu(II)/SCN[−] with allylic radical should be faster than that of Cu(II)/N₃[−], resulting in poor stereoselectivity in products **3**. To test this hypothesis, the competition reaction was conducted, and results indicated that the reaction of thiocyanation is indeed much faster than azidation eq 1.



In conclusion, we have developed a novel copper-catalyzed trifluoromethylazidation and trifluoromethylthiocyanation of allenes. Various CF₃-allyl azides and thiocyanates were obtained with good yields and regioselectivities, which can be readily transformed to some valuable compounds. Further investigation into the mechanism and application is in progress.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures, characterization, and additional data. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b01677.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: yejx@ecust.edu.cn.

*E-mail: gliu@mail.sioc.ac.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for financial support from the National Basic Research Program of China (973-2015CB856600), the National Nature Science Foundation of China (21225210, 21421091, and 21472219), and the Science Technology Commission of the Shanghai Municipality (12ZR1453400). We thank Xiaojian Liu at Shanghai Institute of Technology for his initial studies on the project.

■ REFERENCES

- (1) For allyl amines, see: (a) Brown, E. G. *Ring Nitrogen and Key Biomolecules*; Springer: Boston, MA, 1998. For allyl sulfones, see: (b) Patai, S.; Rapoport, Z.; Stirling, C. *The Chemistry of Sulfones and Sulfoxides*; Wiley: New York, 1988. (b) Johannsen, M.; Jorgensen, K. A. *Chem. Rev.* **1998**, *98*, 1689.
- (2) (a) de Souza, A. O.; Júnior, R. R. S.; Ferreira-Júlio, J. F.; Rodriguez, J. A.; Melo, P. S.; Haun, M.; Sato, D. N.; Durán, N. *Eur. J. Med. Chem.* **2001**, *36*, 843. (b) Sauerberg, P.; Olsen, G. S.; Jeppesen, L.; Mogensen, J. P.; Pettersson, I.; Jeppesen, C. B.; Daugaard, J. R.; Galsgaard, E. D.; Ynddal, L.; Fleckner, J.; Panajotova, V.; Polivka, Z.; Pihera, P.; Havranek, M.; Wulff, E. M. *J. Med. Chem.* **2007**, *50*, 1495. (c) Reck, F.; Zhou, F.; Girardot, M.; Kern, G.; Eyermann, C. J.; Hales, N. J.; Ramsay, R. R.; Gravestock, M. B. *J. Med. Chem.* **2005**, *48*, 499.
- (3) (a) Hiyama, T. *Organofluorine Compounds: Chemistry and Applications*; Springer: Berlin, 2000. (b) Ojima, I. *Fluorine in Medical Chemistry and Chemical Biology*; Wiley-Blackwell: Chichester, U.K., 2009. (c) Jeschke, P. *ChemBioChem* **2004**, *5*, 570. (d) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881.
- (4) For some reviews on trifluoromethylation, see: (a) Schlosser, M. *Angew. Chem., Int. Ed.* **2006**, *45*, 5432. (b) Ma, J.-A.; Cahard, D. *J. Fluorine Chem.* **2007**, *128*, 975. (c) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2008**, *108*, PR1. (d) Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.* **2011**, *111*, 4475. (e) Studer, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 8950.
- (5) (a) Serafinowski, P. J.; Brown, C. A. *Tetrahedron* **2000**, *56*, 333. (b) Heiser, U.; Gaertner, U.-T.; Demuth, H.-U. Patent WO2011/110613A1; 2011. (c) Kawano, Y.; Watanabe, T.; Sakai, J.; Watanabe, H.; Nagano, M.; Nishimura, T.; Miyadera, T. *Chem. Pharm. Bull.* **1980**, *28*, 70.
- (6) (a) Chen, P.; Liu, G. *Synthesis* **2013**, *45*, 2919. (b) Ye, Y.; Sanford, M. S. *Synlett* **2012**, *23*, 2005. (c) Egami, H.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2014**, *53*, 8294. (d) Charpentier, J.; Früh, N.; Togni, A. *Chem. Rev.* **2015**, *115*, 650. (e) Liu, X.; Xu, C.; Wang, M.; Liu, Q. *Chem. Rev.* **2015**, *115*, 683.
- (7) For selected examples, see: (a) Parsons, A. T.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2011**, *50*, 9120. (b) Xu, J.; Fu, Y.; Luo, D.; Jiang, Y.; Xiao, B.; Liu, Z.; Gong, T.; Liu, L. *J. Am. Chem. Soc.* **2011**, *133*, 15300. (c) Wang, X.; Ye, Y.; Zhang, S.; Feng, J.; Xu, Y.; Zhang, Y.; Wang, J. *J. Am. Chem. Soc.* **2011**, *133*, 16410. (d) Shimizu, R.; Egami, H.; Hamashima, Y.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 4577. (e) Janson, P. G.; Ghoneim, I.; Ilchenko, N. O.; Szabó, K. J. *Org. Lett.* **2012**, *14*, 2882. (f) Li, Y.; Studer, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 8221. (g) Yasu, Y.; Koike, T.; Akita, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 9567. (h) Zhu, R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2012**, *134*, 12462. (i) Mizuta, S.; Verhoog, S.; Engle, K. M.; Khotavivattana, K. M.; O'Duill, M.; Wheelhouse, K.; Rassias, G.; Médebielle, M.; Gouverneur, V. *J. Am. Chem. Soc.* **2013**, *135*, 2505. (j) Wu, X.; Chu, L.; Qing, F. *Angew. Chem., Int. Ed.* **2013**, *52*, 2198. (k) Egami, H.; Kawamura, S.; Miyazaki, A.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 7841. (l) Feng, C.; Loh, T.-P. *Angew. Chem., Int. Ed.* **2013**, *52*, 12414. (m) Kong, W.; Casimiro, M.; Merino, E.; Nevado, C. *J. Am. Chem. Soc.* **2013**, *135*, 14480. (n) Liu, X.; Xiong, F.; Huang, X.; Xu, L.; Li, P.; Wu, X. *Angew. Chem., Int. Ed.* **2013**, *52*, 6962. (o) Chen, Z.-M.; Bai, W.; Wang, S.-H.; Yang, B.-M.; Tu, Y.-Q.; Zhang, F.-M. *Angew. Chem., Int. Ed.* **2013**, *52*, 9781. (p) Li, L.; Deng, M.; Zheng, S.-C.; Xiong, Y.-P.; Tan, B.; Liu, X.-Y. *Org. Lett.* **2014**, *16*, 504.
- (8) (a) Xiao, Z.; Hu, H.; Ma, J.; Chen, Q.; Guo, Y. *Chin. J. Chem.* **2013**, *31*, 939. (b) Xu, T.; Cheung, C. W.; Hu, X. *Angew. Chem., Int. Ed.* **2014**, *53*, 4910. (c) Iqbal, N.; Jung, J.; Park, S.; Cho, E. J. *Angew. Chem., Int. Ed.* **2014**, *53*, 539. (d) Maji, A.; Hazra, A.; Maiti, D. *Org. Lett.* **2014**, *16*, 4524. (e) Gao, P.; Shen, Y.-W.; Fang, R.; Hao, X.-H.; Qiu, Z.-H.; Yang, F.; Yan, X.-B.; Wang, Q.; Gong, X.-J.; Liu, X.-Y.; Liang, Y.-M. *Angew. Chem., Int. Ed.* **2014**, *53*, 7629. (f) Xu, J.; Wang, Y.-L.; Gong, T.-J.; Xiao, B.; Fu, Y. *Chem. Commun.* **2014**, *50*, 12915.
- (9) (a) Tsuchii, K.; Imura, M.; Kamada, N.; Hirao, T.; Ogawa, A. *J. Org. Chem.* **2004**, *69*, 6658. (b) Yu, Q.; Ma, S. *Chem. - Eur. J.* **2013**, *19*, 13304. (c) Wang, Y.; Jiang, M.; Liu, J.-T. *Adv. Synth. Catal.* **2014**, *356*, 2907.
- (10) (a) Wang, F.; Qi, X.; Liang, Z.; Chen, P.; Liu, G. *Angew. Chem., Int. Ed.* **2014**, *53*, 1881. (b) Wang, F.; Wang, D.; Mu, X.; Chen, P.; Liu, G. *J. Am. Chem. Soc.* **2014**, *136*, 10202. (c) Liang, Z.; Wang, F.; Chen, P.; Liu, G. *J. Fluorine Chem.* **2014**, *167*, 55. (d) Liang, Z.; Wang, F.; Chen, P.; Liu, G. *Org. Lett.* **2015**, *17*, 2438.
- (11) Wang, F.; Zhu, N.; Chen, P.; Ye, J.; Liu, G. *Angew. Chem., Int. Ed.* **2015**, *54*, in press (DOI: 10.1002/anie.201503412).
- (12) When Tempo was used as radical scavenger, the reaction provided adducts of tempo-CF₃ and tempo-allyl, which suggests the possibility of the CF₃ and allyl radical in a catalytic cycle. For details, see the Supporting Information.