



Novel autophagy modulators: Design and synthesis of (+)-epogymnolactam analogues and structure-activity relationship

Kazuki Ueda, Yuji Okado, Kengo Shigetomi, Makoto Ubukata*

Graduate School of Agriculture, Hokkaido University, Kita-9, Nishi-9, Kita-ku, Sapporo 060-8589, Japan

ARTICLE INFO

Keywords:

Epogymnolactam
Autophagy inducer
Autophagy inhibitor
Structure activity relationship
NIH3T3 cells

ABSTRACT

(+)-Epogymnolactam (**1**) was discovered as a novel autophagy inducer from a culture of *Gymnopus* sp. in our laboratory. To determine structure-activity relationships among (+)-epogymnolactam analogues comparing with cerulenin (**2**), we synthesized 5 analogues including (–)-epogymnolactam (**3**) having each different functional group, and 3 analogues with different side-chain lengths. Five analogues, **3**, **4**, **5**, **6**, and **7** did not significantly increase the ratio of LC3-II to LC3-I as an autophagy marker in NIH3T3 cells. These results suggest that presence and stereochemistry of (2*R*,3*S*)-epoxy group and cyclic *syn*-form (**1b**) of **1** are important for the activity as autophagy inducer. Hexyl analogue (**8**) as well as **1** having butyl side-chain dose-dependently increased the ratio of LC3-II to LC3-I, whereas octyl analogue (**9**) and **2** rather decreased the ratio. Decyl analogue (**10**) did not give a change in the ratio. Although **8** seemed to be an excellent autophagy inducer, it dose-dependently increased SQSTM1 (p62) as in the case of **2**, whereas **1** showed a slight dose-dependent decrease of p62 as an index of autophagic protein degradation. These observations suggest that **8** is an autophagy modulator with different molecular target from **1** or **2**.

1. Introduction

Autophagy inducer and autophagy inhibitor are important for understanding precise mechanisms of autophagy and developing therapeutic agents for such diseases as neurodegenerative diseases, cancer, and infectious diseases.¹ (+)-Epogymnolactam (**1**) was discovered as a novel autophagy inducer from a culture of *Gymnopus* sp. in our laboratory.² Total synthesis of **1** was also achieved to confirm its absolute structure³ and to evaluate further biological properties compared to cerulenin (**2**),⁴ an inhibitor of fatty acid synthesis. Synthetic **1** showed the same activity as natural **1** as autophagy inducer. Although the structure of **1** was similar to that of **2**, major tautomer of **1** seemed to be different from that of **2** as shown in Fig. 1 and Supplemental Table S1. In addition, a contrasting autophagic response was observed in NIH3T3 cells. Cerulenin (**2**) dose-dependently decreased LC3-II/LC3-I, a ratio of lipidate LC3 (LC3-II) to LC3-I as an autophagic marker, and increased p62/actin, a ratio of SQSTM1 (ubiquitin-binding protein p62) to actin, as an index of autophagic degradation.⁵ Thus, cerulenin (**2**) was thought to be an autophagy inhibitor, which is discussed in more detail in the next section.

We, therefore, designed and synthesized 8 analogues of **1**, and evaluated their autophagic activities compared to **1** and **2** in NIH3T3 cells. In this paper, we describe syntheses and biological evaluations of

(+)-epogymnolactam analogues (**3**–**10**) to ascertain the structural requirements of **1** as an autophagy-inducing agent. We also propose hexyl analogue **8** as a new autophagy modulator and discuss several suggestions on the relationship between autophagy and fatty acid synthesis in mammalian cells.

2. Results and discussion

To determine the importance of the stereochemistry of **1** as autophagy inducer, we first synthesized (–)-epogymnolactam (**3**). Epoxy acetate **12** was obtained from *cis*-4-benzyloxy-2-buten-1-ol **11** in 98% ee using Sharpless epoxidation followed by lipase kinetic resolution. Hydrolysis of **12** afforded enantiopure epoxy alcohol **13**, and **3** was synthesized from **13** according to essentially the same method as the total synthesis of **1**³ (Scheme 1).

(+)-Epogymnolactam (**1**) existed mainly as a cyclic *syn*-form (**1b**) in CD₃OD and **1b** was formed even in CDCl₃, whereas NMR of **2** in CDCl₃ showed *keto* form as shown in Fig. 1 and Supplemental Table S1. To determine the importance of *syn* form (**1b**) for autophagy-inducing activity, three chain analogues (**4**–**6**) and *anti*-form analogue (**7**) were synthesized.

Ammonolysis of γ -octalactone **14** with liquid ammonia in a pressure vessel⁶ gave amide alcohol **15** in 88%. Dess-Martin oxidation of **15**

* Corresponding author.

E-mail address: m-ub@for.agr.hokudai.ac.jp (M. Ubukata).

<https://doi.org/10.1016/j.bmc.2018.09.013>

Received 3 July 2018; Received in revised form 13 September 2018; Accepted 15 September 2018

0968-0896/© 2018 Elsevier Ltd. All rights reserved.






Fig. 1. Structures of (+)-epogymnolactam (1) and cerulenin (2). Both cyclic *syn*-form **1b** and *anti*-form **1c** as well as *keto* form **1a** were observed in CD₃OD or even in CDCl₃, whereas only *keto* form of **2** was observed in CDCl₃. Ring-chain tautomerism of **2** was observed in CD₃OD. The tautomeric ratio changed into different ratio with time.



afforded deepoxy analogue **4** as shown in [Scheme 2](#). Deepoxy analogue **4** existed only as a chain form in CDCl₃ ([Supplementary Figs. S1 and S2](#)) by releasing from the steric constraint by the epoxy ring.

Epoxy amide alcohol (**5**) was synthesized as the synthetic precursor³ of **1** ([Scheme 3](#)). Treatment of lactone **16** with (CH₃)₂NH gave **17** which was subjected to Dess-Martin oxidation to give *N,N*-dimethyl analogue (**6**) as indicated in [Scheme 3](#). Chain forms of **5** and **6** were confirmed by their NMR in CDCl₃ as shown in [Supplementary Figs. S3–S6](#).

Cyclic *O*-methyl analogue **7** was prepared by treating **1** with camphor sulfonic acid (CSA) in CH₃OH as shown in [Scheme 3](#). In order to avoid steric and electronic repulsion, CH₃OH attacked from the opposite side of the epoxy group in the carbocation intermediate to give only *anti* form (**7**) as in the case of previously reported observation.⁷ ¹H and ¹³C NMR spectra of **7** indicated its cyclic *anti*-form as shown in [Supplementary Figs. S7 and S8](#).


All these analogues (**3–7**) did not enhance LC3-II/LC3-I, the ratios of LC3-II to LC3-I, and LC3-II/actin, the ratios of LC3-II to actin in NIH-3T3 cells as shown in [Fig. 2](#). These observations suggest that the unknown molecular target of **1** recognizes the absolute configuration of the epoxy group and the cyclic structure of the *syn*-form (**1b**).

In the case of ring-chain tautomerism of **1**, the major tautomer is supposed to be *syn*-form (**1b**) in polar solution, because amide nitrogen of **1a** attacks C4 carbonyl from the opposite side of the epoxy group to avoid steric and electronic repulsion ([Supplemental Table S1](#)). These

considerations suggest that the *syn*-form (**1b**) is the active chemical species in the active site of the unknown primary molecular target. Attempts to synthesize *syn*-form analogue corresponding to **7** have not yet been successful.

Next, we synthesized hexyl, octyl, and decyl analogues (**8–10**) to investigate the effect of side-chain length on biological responses in NIH-3T3 cells. Aldehyde **18**, a synthetic intermediate³ of **1**, was reacted with allyltributyltin in the presence of SnCl₄ to afford homoallylic alcohol **19** and its chlorohydrin.⁸ Treatment of the by-product chlorohydrin with K₂CO₃ gave an additional **19**, and the overall yield of **19** was 71%. Selective hydrogenolysis of benzyl group of **19** was accomplished by using lithium 4,4'-di-*tert*-butylbiphenylide (LiDBB)^{9,10} to give **20** in 61%. Oxidation of **20** with 1,5-dimethyl-9-azanoradamantane *N*-oxyl (DMN-AZADO)¹¹ in combination with [bis(acetoxyl)iodo]benzene (BAIB)¹² gave lactone **21**, which was treated with NH₃ to afford the desired amide **22**. Olefin cross-metathesis of **22** with pent-1-ene gave **23** using Grubbs 2nd-generation catalyst.^{13,14} Chemoselective hydrogenation of **23** using Pd/C-ethylenediamine (Pd/C(en)) catalyst¹⁵ gave **24** which was converted into desired hexyl analogue **8** by Dess-Martin oxidation as shown in [Scheme 4](#). Hexyl analogue **8** mainly existed as a chain form in CDCl₃ as shown in [Supplementary Figs. S9 and S10](#).

Similarly, decyl analogue **10** was synthesized by using 1-nonene as an alkene as shown in [Scheme 5](#). NMR analysis indicated that **10** exists



Scheme 3. Synthesis of epoxy amide alcohol (5), *N,N*-dimethyl analogue (6), and *O*-methyl analogue (7).

as a chain form in CDCl₃ as shown in Supplementary Figs. S13 and S14. Octyl analogue **9** was prepared by chemoselective hydrogenation of cerulenin **2** using Pd/C(en) (Scheme 5). Major tautomer of **9** in CDCl₃ was chain form, while cyclic *syn*- and *anti*-forms were also observed as shown in Supplementary Figs. S11 and S12.

Fig. 3 shows the profiles of LC3-II/LC3-I and LC3-II/actin as autophagic markers after stimulation with **1**, **8**, **9**, **2**, and **10** in NIH3T3 cells. Except for **1**, only hexyl analogue **8** increased the relative amount of LC3-II to LC3-I or actin in a dose-dependent manner. In contrast, octyl analogue **9** and cerulenin **2** were suggested to cause dose-

dependent down-regulation, and decyl analogue **10** did not alter the relative amount of LC3-II.

Our preliminary experiments suggested that cerulenin (**9**) showed cytotoxicity to NIH3T3 cells at these concentrations. We therefore evaluated cytotoxicity of **1**, **2**, **8**, **9**, and **10** in NIH3T3 cells. Only hexyl analogue **8** showed lower cytotoxicity than **2**, **9**, and **10** as in the case of **1** (Fig. 4). Inhibition of fatty acid synthase with **2** induced apoptosis in multiple myeloma cells¹⁶ and human breast cancer cells.¹⁷ We observed significant cell shrinkage as a signal to apoptosis in NIH 3T3 cells¹⁸ treated with **9** at 50 or 100 μM for 4 h, while **1** did not cause significant





Fig. 2. Evaluation of importance of functional groups on autophagy-inducing activity of (+)-epogymnolactam (**1**). NIH3T3 cells were treated for 4 h with DMSO as the vehicle (control), 5 μM rapamycin (positive control), or indicated concentrations of **1**, **3**, **4**, **5**, **6**, or **7**. LC3-II/LC3-I: relative ratio of LC3-II to LC3-I. LC3-II/actin: relative ratio of LC3-II to actin. The data shown represent the mean ± SD of triplicate assay. All significant correlation values (p < 0.05) are given with the indicated significance (*p < 0.05, **p < 0.01) compared to the DMSO control.




Scheme 4. Synthesis of hexyl analogue (8).

cell shrinkage at these concentrations but **8** showed cell shrinkage at 100 μ M (data not shown).

NMR experiments showed that (+)-epogymnolactam (**1**) existed primarily as a cyclic *syn*-form in CD₃OD and a considerable amount of the *syn*-form was observed even in CDCl₃, while hexyl analogue **8**

existed as a chain form in CDCl₃. Despite the close resemblance of the biological properties of **1** and **8**, the physicochemical properties of both are different.

In addition to LC3-II, SQSTM1 (p62) can also be used as autophagy marker to especially determine autophagy flux.⁵ Thus, p62/actin, the



Scheme 5. Synthesis of octyl analogue (9) and decyl analogue (10).




Fig. 3. Effect of side-chain length of (+)-epogymnolactam analogue on autophagy-inducing activity. NIH3T3 cells were treated for 4 h with DMSO as the vehicle (control), 5 μM rapamycin (positive control) or indicated concentrations of **1**, **8**, **9**, **2**, or **10**. LC3-II/LC3-I: relative ratio of LC3-II to LC3-I. LC3-II/actin: relative ratio of LC3-II to actin. The data shown represent the mean \pm SD of triplicate assay. All significant correlation values ($p < 0.05$) are given with the indicated significance (* $p < 0.05$, ** $p < 0.01$) compared to the DMSO control.

up-regulation or degradation ratio of p62 to actin was evaluated in NIH3T3 cells treating with **1** and **8**. (+)-Epogymnolactam (**1**) did not show significant upregulation or downregulation of p62 after 4 h, whereas hexyl analogue **8** upregulated p62 in a dose-dependent manner as indicated in Fig. 5. Octyl analogue **9** also upregulated as in the case of **2**, but dose-dependency was not observed, presumably due to its cytotoxicity.

In conclusion, the study on the structural requirement of **1** indicated that presence and stereochemistry of (2*R*,3*S*)-epoxy group, the molecular shape of the *syn*-form (**1b**), and butyl side chain are critical for inducing autophagy in NIH3T3 cells. We also found that hexyl analogue **8** is a novel autophagy modulator that differs from **1** as an autophagy inducer and **2** as an autophagy inhibitor. SQSTM1 protein (p62) serves as a link between LC3 and ubiquitinated substrates. SQSTM1 and SQSTM1-bound polyubiquitinated proteins become incorporated into the completed autophagosome and are degraded in autolysosomes, thus serving as an index of autophagic degradation. Inhibition of autophagy correlates with increased levels of p62 in mammals, suggesting that steady state levels of the protein reflect the autophagic status. In contrast, decreased p62 levels are associated with autophagy activation.⁵ These considerations suggest that **8** could be an inhibitor of autolysosome formation. However, many cellular-stresses can induce the formation of ubiquitinated protein aggregates. The p62-positive aggregates are also formed by proteasome inhibition or puromycin treatment and can be found in cells exposed to rapamycin for extended periods where the rates of autophagy are elevated.⁵ Because **1** upregulated p62 at 6 h exposure where the rate of autophagy is promoted

(data not shown), the upregulation of p62 at 4 h exposure with **8** can be explained by cellular stress caused by this compound. Thus, we concluded that **1**, **9** (or **2**), and **8** are different types of autophagy modulators which might be useful to determine the relationships between fatty acid synthesis and autophagy induction.

Zimmerman et al. reported that very long chain fatty acid (VLCFA) synthesis has important implications for autophagy and cell homeostasis in yeast.¹⁹ After **1** was discovered as an autophagy inducer,¹ Régnacq et al. reported that the effect of **2** was biphasic that moderate doses (2 or 4 μM) restored autophagy and high doses (20 or 40 μM) inhibited autophagy in lipid droplets (LDs)-deficient yeast. The biphasic effect of **2** was specific to nitrogen-starvation autophagy, and only the inhibitory effect of **2** was observed in rapamycin-induced autophagy in the LDs-deficient mutant.²⁰ Shpilka et al. observed that suppression of autophagy by **2** correlated with a reduction in the number of LDs in yeast cells.²¹ Although these observations suggest that fatty acid synthesis correlates with autophagy in yeast cells, precise mechanism has not been elucidated yet.

The inhibitory activity of **9** on fatty acid synthesis is slightly weaker than that of **2**.²² However, the inhibitory activities of 3-hexenyl analogue (an unsaturated derivative of **8**) and 3-butenyl analogue (an unsaturated derivative of **1**) are much weaker than **2**.^{2,23,24} These observations suggest that both **2** and **9** inhibit basal autophagy and cell proliferation with inhibition of fatty acid synthesis, while **1** and **8** cause different types of autophagic responses together with moderate suppression of cell proliferation without potent inhibition of fatty acid synthesis in NIH3T3 cells.




Fig. 4. Antiproliferation activities of (+)-epogymnolactam (**1**), hexyl analogue (**8**), octyl analogue (**9**), decyl analogue (**10**), and cerulenin (**2**). NIH3T3 cells were treated with DMSO as the vehicle (Control) or indicated concentrations of **1**, **8**, **9**, **2**, or **10**. Relative viable cells were determined using Cell Counting Kit-8. The data shown represent the mean \pm SD of triplicate assay.




Fig. 5. Effect of (+)-epogymnolactam (1) on autophagy flux compared to that of hexyl analogue (8) or octyl analogue (9). NIH3T3 cells were treated for 4 h with DMSO as the vehicle (control), 5 μM rapamycin (positive control) or indicated concentrations of 1, 8, 9, or 2. p62/actin: relative ratio of SEQSTM1 (p62) to actin. The data shown represent the mean ± SD of triplicate assay. All significant correlation values ($p < 0.05$) are given with the indicated significance (* $p < 0.05$, ** $p < 0.01$) compared to the DMSO control.

In hepatocytes, autophagy regulates lipid content because: inhibition of autophagy increased triacylglycerol (TG) store and LDs; loss of autophagy decreased TG breakdown; TGs and LD structural proteins colocalized with autophagic compartments; and LC3 associated with LDs.²⁵

Mobilization of cholesterol and fatty acids from cholesteryl ester (CE) and TG stores of all cells requires lipolytic enzymes.²⁶ Kobayashi et al. demonstrated that a 1:1 mixture of dinapinones A1 and A2 (DPA_{mix}) enhanced neutral lipid degradation together with induction of autophagy in Chinese hamster ovary-K1 cells.²⁷ These results suggest interrelationship between autophagy and TG breakdown in mammalian cells.

Thus, the use of (+)-epogymnolactam (1) and hexyl analogue (8) in combination with cerulenin (2) or octyl analogue (9) may be useful for determining relationship between autophagy and TG breakdown or fatty acid synthesis. These compounds may be also useful to develop therapeutic agents for dyslipidemia, and provide a new perspective on understanding autophagy induction.

Further systematic studies are necessary to determine the effect of lipid metabolism on autophagy induction in mammalian cells and to determine the molecular targets of 1 and 8.

3. Experimental procedure

3.1. Chemicals

All commercially available chemicals of highest purity were used without further purification. (+)-Epogymnolactam (1) was synthesized according to the previously reported procedure.³ Cerulenin (2) was purchased from Wako Pure Chemical Co. (now FUJIFILM Wako Pure Chemical Co., Japan). Rapamycin as an autophagy inducer was obtained from Cayman Chemical (MI, USA). Thin layer and column chromatography were performed with Merck Silica Gel 60 F₂₅₄ and Kanto Chemicals Co. (Japan) Silica Gel 60 N (spherical, neutral), respectively. NMR spectra were measured on a JEOL JMN EX-270 FT-NMR spectrometer. Chemical shifts are reported in ppm (δ -scale) using TMS as the internal standard and coupling constants (J) are given in Hz. The carbon number used for NMR assignment was indicated according to the number shown in 1 and 2 of Fig. 1. MS spectra were acquired with FI or FD mode on a JEOL JMS-T100GCV equipment. Optical rotations were measured on a JASCO P-2000 polarimeter.

3.1.1. (2*S*,3*R*)-4-Benzoyloxy-2,3-epoxybutan-1-ol (13, 88% ee)

To 4 Å molecular sieves (2.29 g) in CH₂Cl₂ (32 ml), Ti(OⁱPr)₄ (7.2 ml, 24.3 mmol) was added, and L-(+)-DIPT (5.03 ml, 23.9 mmol) was added dropwise at -25 °C under an argon atmosphere. After stirring for 30 min, allyl alcohol 11 (4.00 g, 22.4 mmol) in CH₂Cl₂ (32 ml) was added dropwise over 1.5 h. To the mixture, *t*-BuOOH in nonane (5.5 M, 8.8 ml) was added dropwise, then the reaction temperature was raised to -20 °C and stirred for 3 days. The reaction mixture was raised to room temperature, quenched with a saturated Na₂S₂O₃ solution

(40 ml), and stirred for 2 h. The resultant solution was filtered through celite, Et₂O was added, and the mixture was washed with brine. The organic layer was dried with anhydrous Na₂SO₄ and evaporated in vacuo. The residue was subjected to silica gel column chromatography (EtOAc: hexane = 1:1) to give epoxy alcohol 13 (2.98 g, 68.4%). The enantiomeric excess of 13 was determined as 85.7% ee from the peak-area ratio by chiral HPLC analysis (DAICEL Chiralpak AD-H (ϕ 0.45 cm × 25 cm), 254 nm, EtOH : hexane = 1:9).

$[\alpha]_D^{25} = -18.4$ (c 1.0, CHCl₃).

¹H NMR (CDCl₃, 270 MHz): δ 3.18–3.31 (2H, m, H-2 and H-3), 3.62–3.75 (4H, m, H-1 and H-4), 4.50–4.64 (2H, m, benzyl), 7.30–7.36 (5H, m, aromatic).

¹³C NMR (CDCl₃, 67.5 MHz): δ 54.7 (C-3), 55.6 (C-2), 60.7 (C-1), 68.0 (C-4), 73.5 (benzyl), 127.8 (aromatic), 128.0 (aromatic), 128.5 (aromatic), 137.4 (aromatic).

FI-MS: $m/z = 194.1$ [M]⁺.

3.1.2. (2*S*,3*R*)-4-Benzoyloxy-2,3-epoxybutan-1-yl acetate (12)

To epoxy alcohol 13 (1.0 g, 5.1 mmol) in vinyl acetate (39.3 ml), porcine pancreatic lipase (PPL) was added, and the reaction mixture was stirred for 14 h. The reaction mixture was filtered through celite and evaporated in vacuo. The residue was subjected to silica gel column chromatography (EtOAc: hexane = 1:2) to give epoxy acetate 12 (0.99 g, 4.17 mmol). The enantiomeric excess of 12 was determined to be 98.1% ee from the peak-area ratio by chiral HPLC analysis (DAICEL Chiralpak AD-H (ϕ 0.45 cm × 25 cm), 254 nm, EtOH : hexane = 1:9).

$[\alpha]_D^{25} = -19.4$ (c 1.0, CHCl₃).

¹H NMR (CDCl₃, 270 MHz): δ 2.10 (3H, s, Ac), 3.22–3.32 (2H, m, H-2 and H-3), 3.60 (1H, dd, $J = 11.6, 5.9$ Hz, H-4a), 3.72 (1H, dd, $J = 11.2, 4.2$ Hz, H-4b), 4.04 (1H, dd, $J = 12.1, 7.0$ Hz, H-1a), 4.32 (1H, dd, $J = 12.4, 3.8$ Hz, H-1b), 4.51–4.64 (2H, m, benzyl), 7.28–7.36 (5H, m, aromatic).

¹³C NMR (CDCl₃, 67.5 MHz): δ 20.7 (–CH₃), 53.0 (C-2), 54.7 (C-3), 62.6 (C-1), 67.8 (C-4), 73.4 (benzyl), 127.8 (aromatic), 127.9 (aromatic), 128.4 (aromatic), 137.5 (aromatic), 170.6 (–C(=O)–).

FI-MS: $m/z = 236.1$ [M]⁺.

3.1.3. (2*S*,3*R*)-4-Benzoyloxy-2,3-epoxybutan-1-ol (enantiopure 13)

To epoxy acetate 12 (499.5 mg, 2.11 mmol) in THF (6.09 ml), CH₃OH (6.09 ml) was added dropwise and then K₂CO₃ (58.0 mg) was added at 0 °C. The reaction mixture was stirred for 2 h at the same temperature. The reaction was quenched with a saturated NH₄Cl solution (40 ml), and the mixture was extracted with Et₂O. The organic layer was dried with anhydrous Na₂SO₄ and evaporated in vacuo. The residue was subjected to silica gel column chromatography (EtOAc: hexane = 2:1) to give enantiopure epoxy alcohol 13 (375.9 mg, 91%).

$[\alpha]_D^{25} = -23.1$ (c 1.0, CHCl₃).

3.1.4. (–)-Epogymnolactam (3)

(2*S*,3*S*)-3-(1-Hydroxypentyl)oxiran-2-carboxamide was derived from enantiopure 13 using essentially the same method as the synthesis

of **1**.³ To the amide (19.5 mg, 0.11 mmol) in CH₂Cl₂ (1.6 ml), Dess-Martin periodinane (67.1 mg, 1.45 eq.) was added at 0 °C under an argon atmosphere. After stirring for 2 h, the reaction mixture was quenched with a mixture of saturated Na₂S₂O₃ solution and saturated NaHCO₃ and extracted with EtOAc. The organic layer was washed with brine, dried with anhydrous Na₂SO₄ and evaporated in vacuo. The residue was subjected to silica gel column chromatography (EtOAc: hexane = 2: 1) to give (-)-epogymnolactam **3** (11.3 mg, 59%).

$[\alpha]_{\text{D}}^{25} = -14.4$ (c 1.1, CH₃OH), [(+)-epogymnolactam **1**: $[\alpha]_{\text{D}}^{20} = +47.8$ (c 1.0, CH₃OH, incubation time is different)].²

¹H NMR (CD₃OD, 270 MHz): *Keto* form: δ 0.88–0.97 (3H, m, H-8), 1.28–1.58 (4H, m, H-6 and H-7), 2.57–2.69 (m, H-5), 3.70 (d, $J = 5.3$ Hz, H-3), 3.89 (d, $J = 5.3$ Hz, H-2); *syn*-form: 0.88–0.97 (3H, m, H-8), 1.28–1.58 (4H, m, H-6 and H-7), 1.69–1.75 (m, H-5), 3.58 (1H, d, $J = 2.6$ Hz, H-3), 3.80 (1H, d, $J = 2.6$ Hz, H-2); *anti*-form: 0.88–0.97 (3H, m), 1.28–1.41 (2H, m), 1.49–1.58 (2H, m), 1.69–1.75 (2H, m), 3.57 (d, $J = 2.6$ Hz, H-3), 3.81 (d, $J = 2.6$ Hz, H-2).

¹³C NMR (CD₃OD, 67.5 MHz): *Keto* form: δ 14.1 (C-8), 23.2 (C-7), 26.1 (C-6), 41.0 (C-5), 55.8 (C-2), 59.4 (C-3), 170.5 (C-1), 205.8 (C-4); *syn*-form: 14.3 (C-8), 24.0 (C-7), 27.0 (C-6), 36.3 (C-5), 50.1 (C-3), 53.1 (C-2), 87.2 (C-4), 174.4 (C-1); *anti*-form: 14.3 (C-8), 23.9 (C-7), 25.9 (C-6), 38.9 (C-5), 54.3 (C-2), 58.1 (C-3), 86.8 (C-4), 173.0 (C-1).

FI-MS: $m/z = 172.1$ [M+H]⁺.

3.1.5. 4-Hydroxyoctanamide (15)

γ -Octalactone **14** (420.0 mg, 3.0 mmol) was mixed with approximately 20 ml of liquid NH₃ in a pressure-resistance reaction tube, and the mixture was stirred at 50 °C for 7 days. The tube was opened at -196 °C, and NH₃ was evaporated at ambient temperature. The resultant white solid was purified by silica gel column chromatography (CHCl₃: CH₃OH = 9: 1) to give amide **15** (411.3 mg, 87%) as a white solid.

¹H NMR (CD₃OD, 270 MHz): δ 0.82 (3H, t, $J = 6.8$ Hz, H-8), 1.31–1.84 (8H, m, H-3, H-5, H-6 and H-7), 2.21–2.41 (2H, m, H-2), 3.47–3.56 (1H, m, H-4).

¹³C NMR (CD₃OD, 67.5 MHz): δ 14.4 (C-8), 23.8 (C-7), 29.0 (C-6), 32.9 (C-2), 34.1 (C-3), 38.1 (C-5), 71.9 (C-4), 179.2 (C-1).

FD-MS: $m/z = 160.1$ [M+H]⁺.

3.1.6. Deepoxy analogue (4)

Dess-Martin periodinane (129.7 mg, 2 eq.) was added to a stirred solution of amide **15** (23.7 mg, 148.8 μ mol) in dry CH₂Cl₂ (8.4 ml) at 0 °C under argon atmosphere. After stirring for 3 h, the reaction was quenched with a solution of saturated aqueous Na₂S₂O₃ and saturated aqueous NaHCO₃. The solution was extracted with Et₂O and organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The resultant residue was subjected to silica gel column chromatography (CHCl₃: CH₃OH = 19: 1) to afford deepoxy analogue **4** (14.1 mg, 60%) as a colorless solid.

¹H NMR (CDCl₃, 270 MHz): δ 0.90 (3H, t, $J = 7.3$ Hz, H-8), 1.31 (2H, sext, $J = 7.3$ Hz, H-7), 1.57 (2H, quin, $J = 7.5$, H-6), 2.43–2.51 (4H, m, H-2 and H-3), 2.78 (2H, t, $J = 6.5$ Hz, H-3) (Supplementary Fig. S1).

¹³C NMR (CDCl₃, 67.5 MHz): δ 13.8 (C-8), 22.3 (C-7), 25.9 (C-6), 29.1 (C-2), 37.4 (C-3), 42.5 (C-5), 174.4 (C-1), 210.2 (C-4) (Supplementary Fig. S2).

HR-FD-MS: $m/z = 158.1185$ [M+H]⁺ calcd. for C₈H₁₆NO₂, found 158.1181.

3.1.7. (2R,3R,4R)-2,3-Epoxy-4-hydroxyoctanamide (5)

Lactone **16**² (low polar diastereomer, 49.6 mg, 0.317 mmol) was dissolved in a solution of NH₃ in CH₃OH (2.0 M, 8 ml) under nitrogen atmosphere and the mixture was stirred for 2.5 h at 0 °C. The reaction mixture was evaporated in vacuo and purified by silica gel column chromatography (CHCl₃: CH₃OH = 9: 1) to afford epoxy amide alcohol **5** (54.0 mg, 98.5%).

$[\alpha]_{\text{D}}^{25} = +49.3$ (c 1.0, CH₃OH).

¹H NMR (CDCl₃, 270 MHz): δ 0.92 (3H, t, $J = 7.2$ Hz, H-8), 1.32–1.75 (6H, m, H-5, H-6 and H-7), 3.13 (1H, dd, $J = 7.8, 4.6$ Hz, H-3), 3.42–3.51 (1H, m, H-4), 3.54 (1H, d, $J = 4.6$ Hz, H-2) (Supplementary Fig. S3).

¹³C NMR (CDCl₃, 67.5 MHz): δ 14.0 (C-10), 22.6 (C-7), 27.0 (C-6), 34.6 (C-5), 54.3 (C-2), 60.1 (C-3), 69.0 (C-4), 170.2 (C-1) (Supplementary Fig. S4).

FI-MS: $m/z = 174.1$ [M+H]⁺.

3.1.8. (2R,3R,4R)-2,3-Epoxy-4-hydroxy-N,N-dimethyloctanamide (17)

Lactone **16** (10.0 mg, 64 μ mol) was dissolved in a solution of (CH₃)₂NH in CH₃OH (2.0 M, 1.5 ml) under argon atmosphere and the mixture was stirred for 2.5 h at 0 °C. The reaction mixture was concentrated in vacuo and purified by silica gel column chromatography (CHCl₃: CH₃OH = 96: 4) to afford **17** (8.3 mg, 65%) as a colorless solid.

¹H NMR (CDCl₃, 270 MHz): δ 0.92 (3H, t, $J = 7.0$ Hz, H-8), 1.26–1.73 (6H, m, H-5, H-6 and H-7), 3.00 (3H, s, NCH₃), 3.09 (1H, dd, $J = 3.8, 4.3$ Hz, H-3), 3.20 (3H, s, NCH₃), 3.58 (1H, d, $J = 3.8$ Hz, H-2), 3.87 (1H, m, H-4).

¹³C NMR (CDCl₃, 67.5 MHz): δ 14.0 (C-8), 22.7 (C-7), 27.2 (C-6), 34.5 (C-5), 35.1 (-N(CH₃)₂), 36.7 (-N(CH₃)₂), 53.8 (C-2), 59.9 (C-3), 71.8 (C-4), 167.2 (C-1).

FI-MS: $m/z = 1201.1$ [M]⁺.

3.1.9. N,N-Dimethyl Analogue (6)

To a stirred solution of **17** (8.2 mg, 41 μ mol) in CH₂Cl₂ (2.4 ml), Dess-Martin periodinane (29.0 mg, 1.5 eq.) was added at 0 °C under argon atmosphere. After stirring for 3 h, the reaction was quenched with a mixture of saturated aqueous Na₂S₂O₃ and saturated aqueous NaHCO₃. The resultant solution was extracted with EtOAc and the organic layer was washed with brine, dried over anhydrous Na₂SO₄, concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc: hexane = 2: 1) to give N,N-dimethyl analogue **6** (7.3 mg, 90%) as a colorless solid.

$[\alpha]_{\text{D}}^{25} = -44.2$ (c 0.7, CHCl₃).

¹H NMR (CDCl₃, 270 MHz): δ 0.89 (3H, t, $J = 7.2$ Hz, H-8), 1.30 (2H, sext, $J = 7.4$ Hz, H-7), 1.48–1.62 (2H, m, H-6), 2.55 (2H, dt, $J = 7.4, 7.4, 2.9$ Hz, H-5), 2.93 (3H, s, NCH₃), 3.13 (3H, s, NCH₃), 3.70 (1H, d, $J = 4.9$ Hz, H-3), 3.82 (1H, d, $J = 5.1$ Hz, H-2) (Supplementary Fig. S5).

¹³C NMR (CDCl₃, 67.5 MHz): δ 13.8 (C-8), 22.1 (C-7), 24.8 (C-6), 35.3 (-N(CH₃)₂), 36.3 (-N(CH₃)₂), 39.6 (C-5), 54.7 (C-2), 58.2 (C-3), 164.3 (C-4), 205.5 (C-1) (Supplementary Fig. S6).

HR-FI-MS: $m/z = 200.1290$ [M+H]⁺ calcd. for C₁₀H₁₈NO₃, found 200.1287.

3.1.10. O-Methyl analogue (7)

To a solution of (+)-epogymnolactam **1** (8.5 mg, 50 μ mol) in CH₃OH (3.0 ml), 10-camphorsulfonic acid (3.0 mg, 0.26 eq.) was added, and the reaction mixture was stirred at 50 °C for 6 h. The resulting mixture was diluted with EtOAc and washed H₂O and brine. The organic layer was dried over anhydrous Na₂SO₄, evaporated in vacuo. The resultant oil was purified by silica gel column chromatography (EtOAc: hexane = 1: 2) to afford O-methyl analogue **7** (2.2 mg, 24%) as a colorless oil.

$[\alpha]_{\text{D}}^{25} = +95.3$ (c 0.29, CHCl₃).

¹H NMR (CDCl₃, 270 MHz): δ 0.93 (3H, t, $J = 7.1$ Hz, H-8), 1.26–1.56 (4H, m, H-6 and H-7), 1.79 (2H, t, $J = 7.7$ Hz, H-5), 3.65 (1H, t, $J = 2.3$ Hz, H-3), 3.79 (1H, d, $J = 2.7$ Hz, H-2) (Supplementary Fig. S7).

¹³C NMR (CDCl₃, 67.5 MHz): δ 13.9 (C-8), 22.7 (C-7), 25.7 (C-6), 34.1 (C-5), 49.4 (C-2), 51.7 (-OCH₃), 56.5 (C-3), 89.6 (C-4), 171.5 (C-1) (Supplementary Fig. S8).

HR-FD-MS: $m/z = 186.1166$ [M+H]⁺ calcd. for C₉H₁₆NO₃, found 186.1130.

3.1.11. (5*R*,6*S*)-7-Benzoyloxy-5,6-epoxyhept-1-en-4-ol (**19**)

To a stirred solution of **18**³ (100 mg, 0.521 mmol) in dry CH₂Cl₂ (12 ml), SnCl₄ (65 μl, 1.1 eq.) was added dropwise at -78 °C under argon atmosphere. After stirring for 1 h, allyltributyltin (208 μl, 1.3 eq.) was added dropwise and the reaction mixture was continuously stirred for another 1.5 h. The reaction mixture was quenched with sat. aqueous NH₄Cl at -78 °C and allowed to room temperature. The solution was extracted with Et₂O and the organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The resultant residue was purified by silica gel column chromatography (EtOAc : hexane = 1 : 3) to give a mixture of homoallylic alcohol **19** (74.3 mg) and its chlorohydrin (20 mg). To a solution of chlorohydrin (20 mg) in CH₃OH (1 ml), K₂CO₃ (15.3 mg, 1.5 eq.) was added at 0 °C, and the reaction mixture was allowed to room temperature. After 4 days, additional K₂CO₃ in CH₃OH (1 ml) was added and the reaction mixture was stirred for 5 h. The resultant mixture was quenched with saturated aqueous NH₄Cl and extracted with Et₂O. The organic layer was washed with brine, dried over Na₂SO₄, concentrated in vacuo, and the residue was purified by silica gel column chromatography (EtOAc: hexane = 1 : 3) to afford an additional **19** (12.2 mg). The total yield of **19** was 71%.

¹H NMR (CDCl₃, 270 MHz): δ 2.35–2.56 (2H, m, H-3), 2.96 (1H, dd, *J* = 8.1 and 4.3 Hz, H-5), 3.23–3.29 (1H, m, H-6), 3.47–3.55 (1H, m, H-4), 3.66 (1H, dd, *J* = 10.5, 6.7 Hz, H-7a), 3.81–3.87 (1H, dd, *J* = 10.7, 6.1 Hz, H-7b), 4.58 (2H, m, benzyl), 5.14–5.22 (2H, m, H-1), 5.83–5.93 (1H, m, H-2), 7.30–7.37 (5H, m, aromatic).

¹³C NMR (CDCl₃, 67.5 MHz): δ 39.4 (C-5), 54.1 (C-2), 57.9 (C-3), 68.5 (C-1), 69.0 (C-4), 73.6 (benzyl), 118.3 (C-7), 127.9 (aromatic), 128.6 (aromatic), 133.5 (C-6), 137.2 (aromatic).

FI-MS: *m/z* = 234.1 [M]⁺.

3.1.12. (2*S*,3*R*)-2,3-Epoxyhept-6-en-1,4-diol (**20**)

Lithium wire (20.9 mg, 3.02 mmol) was hammered out into foil, cut into several small pieces and washed with hexane. The pieces were then dipped, one by one, in CH₃OH until the surface turns shiny, then immediately transferred to THF. The lithium chunks were placed in a Schlenk flask containing DBB (534.8 mg, 2.01 mmol). The flask was evacuated and purged with argon several times. Dry THF (12 ml) was added, and the contents of the flask were sonicated between 0 and 20 °C for 3 h. The resultant deep forest green solution of radical anion (4.7 ml) was then added dropwise to a solution of **19** (47.0 mg, 0.201 mmol) in dry THF (5.0 ml) at -78 °C under argon atmosphere. The mixture was stirred for 1.5 h and quenched with sat. aqueous NH₄Cl at -78 °C and allowed to reach room temperature. The resultant solution was extracted with Et₂O and the organic layer was washed with brine, dried over Na₂SO₄, concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc: hexane = 2 : 1) to give a diastereomeric mixture of diol **20** (17.6 mg, 61%) as a colorless oil.

¹H NMR (CDCl₃, 270 MHz): δ 2.28–2.58 (2H, m, H-5), 2.97–3.08 (1H, dd, *J* = 7.8, 4.3 Hz, H-3), 3.20–3.76 (3H, m, H-1a, H-2 and H-4), 3.84–4.01 (1H, m, H-1b), 5.12–5.24 (2H, m, H-7), 5.72–5.96 (1H, m, H-6).

¹³C NMR (CDCl₃, 67.5 MHz): δ 39.8 (C-5), 55.8 (C-2), 58.3 (C-3), 60.7 (C-1), 68.8 (C-4), 119.0 (C-7), 133.1 (C-6).

FD-MS: *m/z* = 145.1 [M+H]⁺.

3.1.13. (1*R*,5*R*)-4-(Prop-2-enyl)-3,6-dioxabicyclo[3.1.0]hexan-2-one (**21**)

PhI(OAc)₂ (462 mg, 3.0 eq.) was added to a stirred solution of **20** (68.8 mg, 0.478 mmol) and DMN-AZADO (4.0 mg, 0.05 eq.) in CH₃CN (4.82 ml) and aqueous phosphate buffer (0.54 ml, 0.2 M, pH 6.8) at 0 °C. After stirring for 5 h, the mixture was quenched with saturated aqueous Na₂S₂O₃. The solution was extracted with EtOAc and the organic layer was washed with brine, dried over Na₂SO₄, concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc: hexane = 1 : 3) to afford a diastereomeric mixture of lactone

21 (50.0 mg, 75%) as a colorless oil.

¹H NMR (CDCl₃, 270 MHz): δ 2.43–2.64 (2H, m, allyl), 3.79 (1H, t, *J* = 2.7 Hz, H-5), 4.03 (d, *J* = 2.4 Hz, H-1 of major diastereomer), 4.11 (dd, *J* = 2.6, 1.5 Hz, H-1 of minor diastereomer), 4.50 (dt, *J* = 7.1, 1.4 Hz, H-4 of minor diastereomer), 4.64 (t, *J* = 6.2 Hz, H-4 of major diastereomer), 5.17–5.30 (2H, m, olefinic), 5.70–5.92 (1H, m, olefinic).

¹³C NMR (CDCl₃, 67.5 MHz): δ 34.0 and 36.1 (C-5), 49.8 and 50.6 (C-3), 56.2 and 57.6 (C-2), 78.1 and 78.4 (C-4), 119.3 and 120.4 (C-7), 129.8 and 131.2 (C-6), 170.1 (C-1).

FI-MS: *m/z* = 140.1 [M]⁺.

3.1.14. (2*R*,3*R*)-2,3-Epoxy-4-hydroxyhept-6-enamide (**22**)

The diastereomeric mixture of **21** (9.6 mg, 0.067 mmol) was dissolved in a solution of NH₃ in CH₃OH (2.0 M, 1.3 ml) under argon atmosphere and the mixture was stirred at 0 °C for 3 h. The resulting solution was concentrated in vacuo and purified by silica gel column chromatography (CHCl₃: CH₃OH = 93 : 7) to afford a diastereomeric mixture of amide **22** (10.3 mg, 96%) as a colorless solid.

¹H NMR (CDCl₃, 270 MHz): δ 2.33–2.55 (2H, m, H-5), 3.14–3.22 (1H, dd, *J* = 8.1, 4.6 Hz, H-3), 3.46–3.66 (2H, m, H-2 and H-4), 5.14–5.23 (2H, m, H-7), 5.72–5.97 (1H, m, H-6).

¹³C NMR (CDCl₃, 67.5 MHz): δ 38.2 and 39.3 (C-5), 54.3 and 54.5 (C-2), 59.6 and 60.8 (C-3), 67.9 and 68.5 (C-4), 118.4 and 118.8 (C-7), 132.7 and 133.1 (C-6), 169.9 and 170.6 (C-1).

FD-MS: *m/z* = 158.1 [M+H]⁺.

3.1.15. (2*R*,3*R*)-2,3-Epoxy-4-hydroxydec-6-enamide (**23**)

Grubbs 2nd generation catalyst (3.1 mg, 0.1 eq.) was added to a stirred solution of **22** (5.7 mg, 36.3 μmol) and 1-pentene (70 mg) in dry and degassed CH₂Cl₂ (1.0 ml) under argon and the resulting mixture was heated to reflux for 24 h. The mixture was concentrated in vacuo and purified by silica gel column chromatography (EtOAc: hexane = 4 : 1) to afford a diastereomeric mixture of amide **23** (4.4 mg, 61%) as a colorless solid.

¹H NMR (CDCl₃, 270 MHz): δ 0.96 (3H, t, *J* = 7.2 Hz, H-10), 1.23–1.43 (2H, m, H-9), 1.98–2.11 (2H, m, H-8), 2.17–2.51 (2H, m, H-5), 3.14–3.25 (1H, m, H-3), 3.47–3.65 (2H, m, H-2 and H-4), 5.32–5.65 (2H, m, H-6 and H-7), 6.14–6.22 (1H, s).

¹³C NMR (CDCl₃, 67.5 MHz): δ 13.6 (C-10), 22.4 (C-9), 34.7 (C-8), 37.1 and 38.1 (C-5), 54.2 and 54.3 (C-2), 59.5 and 60.8 (C-3), 68.6 and 69.2 (C-4), 123.5 and 123.9 (C-7), 135.3 and 135.9 (C-6), 169.1 and 169.6 (C-1).

FD-MS: *m/z* = 200.1 [M+H]⁺.

3.1.16. (2*R*,3*R*)-2,3-Epoxy-4-hydroxydecanamide (**24**)

To a solution of **23** (4.5 mg, 22.6 μmol) in THF (0.7 ml), Pd/C(en) (5.0 mg) was added and the mixture was stirred vigorously overnight under hydrogen atmosphere. The resulting solution was filtered through celite pad, concentrated and the residue was purified by silica gel column chromatography (EtOAc: hexane = 2 : 1) to afford a diastereomeric mixture of amide **24** (3.5 mg, 77%) as a colorless solid.

¹H NMR (CDCl₃, 270 MHz): δ 0.88 (3H, t, *J* = 7.2 Hz, H-10), 1.28–1.70 (10H, m, CH₂), 3.11–3.21 (1H, m, H-3), 3.43–3.60 (2H, m, H-2 and H-4).

¹³C NMR (CDCl₃, 67.5 MHz): δ 14.0 (C-10), 22.5 and 22.6 (C-9), 24.7 and 24.8 (C-8), 29.1 and 29.2 (C-7), 31.6 and 31.7 (C-6), 33.6 and 34.8 (C-5), 54.2 and 54.8 (C-2), 60.0 and 61.4 (C-3), 69.3 and 70.0 (C-4), 169.2 and 169.9 (C-1).

FD-MS: *m/z* = 202.2 [M+H]⁺.

3.1.17. Hexyl analogue (**8**)

To a stirred solution of **24** (3.4 mg, 16.9 μmol) in dry CH₂Cl₂ (1.0 ml), Dess-Martin periodinane (12.2 mg, 1.7 eq.) was added at room temperature under argon atmosphere. After stirring for 3 h, the reaction mixture was purified by silica gel column chromatography (EtOAc : hexane: toluene = 3 : 1 : 1) to afford hexyl analogue **8** (2.1 mg, 62%) as a

colorless solid.

^1H NMR (CDCl_3 , 270 MHz): δ 0.88 (3H, t, J = 7.0 Hz, H-10), 1.28–1.61 (8H, m), 2.48–2.68 (2H, m, H-5), 3.73 (1H, d, J = 5.4 Hz, H-3), 3.88 (1H, d, J = 5.1 Hz, H-2) (Supplementary Fig. S9).

^{13}C NMR (CDCl_3 , 67.5 MHz): δ 14.0 (C-10), 22.4 (C-9), 23.1 (C-8), 28.7 (C-7), 31.4 (C-6), 41.1 (C-5), 55.3 (C-2), 58.3 (C-3), 167.3 (C-1), 202.7 (C-4) (Supplementary Fig. S10).

HR-FD-MS: m/z = 200.1294 $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{10}\text{H}_{18}\text{NO}_3$, found 200.1287.

3.1.18. (2R,3R)-2,3-Epoxy-4-hydroxytetradec-6-enamide (25)

Grubbs 2nd generation catalyst (2.1 mg, 0.1 eq.) was added to a stirred solution of **22** (3.8 mg, 24.3 μmol) and 1-nonene (97.2 mg) in dry and degassed CH_2Cl_2 (2.0 ml) under argon and the mixture was heated to reflux for 24 h. The resulting mixture was concentrated in vacuo and purified by silica gel column chromatography (CHCl_3 : CH_3OH = 19: 1) to afford amide **25** (2.7 mg, 43%) as a colorless solid.

^1H NMR (CDCl_3 , 270 MHz): δ 0.88 (3H, t, J = 6.8 Hz, H-14), 1.27–1.61 (10H, m, CH_2), 1.98–2.06 (2H, m, H-8), 2.25–2.32 (2H, m, H-5), 3.18–3.23 (1H, m, H-3), 3.55–3.57 (2H, m, H-2 and H-4), 5.34–5.63 (2H, m, H-6 and H-7).

^{13}C NMR (CDCl_3 , 67.5 MHz): δ 14.1 (C-14), 22.7 (C-13), 29.15 (C-11), 29.17 (C-9), 29.3 (C-10), 31.8 (C-12), 32.6 (C-8), 37.1 (C-5), 54.2 (C-2), 60.8 (C-3), 69.2 (C-4), 123.2 (C-6), 136.3 (C-7), 169.1 (C-1).

HR-FD-MS: m/z = 256.1917 $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{14}\text{H}_{26}\text{NO}_3$, found 256.1913.

3.1.19. (2R,3R)-2,3-Epoxy-4-hydroxytetradecanamide (26)

Pd/C(en) (6.0 mg) was added to a solution of **25** (3.0 mg, 11.7 μmol) in THF (0.7 ml) and the reaction mixture was stirred vigorously at room temperature under H_2 atmosphere. The resulting mixture was filtered through celite pad and concentrated in vacuo. The residue was purified by silica gel column chromatography (CHCl_3 : CH_3OH = 19: 1) to give amide **26** (2.1 mg, 69%) as a colorless solid.

^1H NMR (CDCl_3 , 270 MHz): δ 0.88 (3H, t, J = 6.8 Hz, H-14), 1.25–1.58 (20H, m, CH_2), 3.16–3.21 (1H, m, H-3), 3.49–3.59 (2H, m, H-2 and H-4).

^{13}C NMR (CDCl_3 , 67.5 MHz): δ 14.1 (C-14), 22.7 (C-13), 24.7 (C-6), 29.3 (C-11), 29.4 (C-8), 29.55 (C-9), 29.57 (C-10), 29.7 (C-7), 31.9 (C-12), 33.6 (C-5), 54.7 (C-2), 61.4 (C-3), 70.0 (C-4), 169.0 (C-1).

HR-FD-MS: m/z = 258.2076 $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{14}\text{H}_{28}\text{NO}_3$, found 258.2069.

3.1.20. Decyl analogue (10)

To a stirred solution of **26** (1.9 mg, 7.4 μmol) in dry CH_2Cl_2 (1.0 ml), Dess-Martin periodinane (5.3 mg, 1.7 eq.) was added at room temperature under argon atmosphere. After stirring for 3 h, the reaction mixture was purified by silica gel column chromatography (CHCl_3 : CH_3OH = 19: 1) to afford decyl analogue **10** (1.3 mg, 69%) as a colorless solid. In the NMR of **10** in CDCl_3 , only the keto form was observed.

^1H NMR (CDCl_3 , 270 MHz): δ 0.88 (3H, t, J = 6.7 Hz, H-14), 1.26 (14H, br s, CH_2), 1.56–1.66 (2H, m, H-6), 2.58 (2H, dt, J = 7.6, 7.6 and 5.1 Hz, H-5), 3.73 (1H, d, J = 5.1 Hz, H-3), 3.87 (1H, d, J = 5.4 Hz, H-2) (Supplementary Fig. S13).

^{13}C NMR (CDCl_3 , 67.5 MHz): δ 14.1 (C-14), 22.7 (C-13), 23.1 (C-6), 29.0 (C-7), 29.2 (C-8), 29.3 (C-11), 29.4 (C-10), 29.5 (C-9), 31.9 (C-12), 41.1 (C-5), 55.3 (C-2), 58.3 (C-3), 167.4 (C-1), 202.6 (C-4) (Supplementary Fig. S14).

HR-FD-MS: m/z = 256.1917 $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{14}\text{H}_{26}\text{NO}_3$, found 256.1913.

3.1.21. Octyl analogue (9)

Pd/C(en) (1.0 mg) was added to a solution of cerulenin **2** (4.3 mg, 19.3 μmol) in THF (0.5 ml) and the reaction mixture was stirred at room temperature overnight under hydrogen atmosphere. The resulting

solution was filtered through celite pad and concentrated in vacuo. The residue was purified by silica gel column chromatography (EtOAc: hexane = 1: 1) to afford octyl analogue **9** (4.4 mg, quant.) as a colorless solid.

$[\alpha]_{\text{D}}^{25}$ = +17.2 (c 0.45, CHCl_3).

^1H NMR (CDCl_3 , 270 MHz): δ 0.85–0.91 (3H, m, H-12), 1.26 (10H, br s, CH_2), 1.55–1.63 (2H, m, H-6), 1.70–1.87 (m, H-5 of cyclic form), 2.48–2.68 (m, H-5 of keto-form), 3.61 (t, J = 2.3 Hz, H-3 of major cyclic form), 3.65 (t, J = 2.4 Hz, H-3 of minor cyclic form), 3.73 (d, J = 5.1 Hz, H-3 of keto-form), 3.80–3.83 (m, H-2 of cyclic form), 3.87 (d, J = 5.4 Hz, H-2 of keto-form) (Supplementary Fig. S11).

^{13}C NMR (CDCl_3 , 67.5 MHz): keto form δ 14.1 (C-12), 22.6 (C-11), 23.1 (C-6), 29.0 (C-9 and C-8), 29.1 (C-7), 31.8 (C-10), 41.1 (C-5), 55.3 (C-2), 58.3 (C-3), 167.3 (C-1), 202.7 (C-4); *syn*- and *anti*-form δ 14.1 (C-12), 22.6 (C-11), 23.6 (C-6), 29.1 (C-9), 29.3 and 29.4 (C-8), 29.6 (C-7), 31.8 (C-10), 37.1 and 37.5 (C-5), 51.5 and 53.9 (C-2), 57.5 and 57.8 (C-3), 86.3 (C-4), 171.4 (C-1) (Supplementary Fig. S12).

HR-FD-MS: m/z = 228.1595 $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{12}\text{H}_{22}\text{NO}_3$, found 228.1599.

3.2. Monitoring of autophagy-inducing activity

The autophagy-inducing activity of each compound was evaluated by slightly modifying the method reported previously.² Briefly, Mouse embryo fibroblast NIH 3T3 cells were inoculated at 3.3×10^5 cells/ml and cultured at 37 °C under 5% CO_2 atmosphere. After 24 h, medium change was made in each well, and the DMSO solution of each compound was added at final concentrations of 25 μM , 50 μM , and 100 μM . The cells were incubated at 37 °C for 4 h, and adherent cells were harvested in 1 ml of PBS (–) and collected by centrifugation. The collected cells were washed twice with PBS (–) and sonicated in RIPA Lysis Buffer (Santa Cruz: sc-24948). After centrifugation, the supernatant was collected and stored at –20 °C until use. Proteins were separated by SDS-PAGE and immunoblot analysis was performed with anti-LC3 antibody (Cell Signaling Technology), anti-SQSTM1/p62 antibody (Funakoshi, Japan) and anti-actin antibody (Cosmo Bio, Sigma-Aldrich). The specific signals from LC3, SQSTM1 (p62), and actin were quantified with the enhanced chemiluminescence reagents (NEL105, Perkin Elmer) by luminescent analyzer LAS-3000 (FUJI FILM, Japan) or ChemiDoc MP (Bio-Rad). As indicated in reference 2, it was difficult to visually judge by conventional immunoblotting analysis. Thus, the relative signal ratio (LC3-II/LC3-I, LC3-II/actin, or p62/actin) was determined from the quantified specific signal of each band in LAS-3000 or ChemiDoc MP to evaluate the autophagy inducing activity or autophagy inhibiting activity.

3.3. Anti-proliferative activity

Anti-proliferative activity was performed with slight modification of the previously reported method.² Using the trypan blue exclusion assay, the number of viable cells of NIH3T3 cells was adjusted to 0.53×10^4 cells/ml. The cells were treated with each compound in DMSO to the final concentrations of 25 μM , 50 μM and 100 μM , and plated on a 96-well plate. The DMSO concentration in all wells was set to 0.1%. After 4 h incubation at 37 °C in a CO_2 incubator, 10 μl of Cell Counting Kit-8 (DOJINDO Laboratories, Japan) was added. After incubating for 14 h, the relative number of viable cells was expressed as the optical density (OD_{450}) value determined by a microplate reader (Sunrise Remote, Tecan Group Ltd., Switzerland).

Acknowledgements

This work was supported by JSPS KAKENHI (Grant-in-Aid for Scientific Research (C)) Grant Number JP 26450135, Japan.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bmc.2018.09.013>.

References

- Mizushima N, Levine B, Cuervo AN, Klionsky DJ. *Nature*. 2008;451:1069.
- Mitsuhashi S, Shindo C, Shigetomi K, Miyamoto T, Ubukata M. *Phytochemistry*. 2015;114:163.
- Okado Y, Shigetomi K, Mitsuhashi S, Ubukata M. *J Antibiot*. 2015;68:721.
- Kawaguchi A, Tomoda H, Nozoe S, Omura S, Okuda S. *J Biochem*. 1982;92:7.
- Klionsky DJ, et al. *Autophagy*. 2016;12:1.
- Taylor SK, Ide ND, Silver ME, Stephan ML. *Synth Commun*. 2001;31:2391.
- Kuramochi K, Mizushima Y, Nagata S, Sugawara F, Sakaguchi K, Kobayashi S. *Bioorg Med Chem*. 1983;2004:12.
- Wang S, Howe GP, Mahal RS, Procter G. *Tetrahedron Lett*. 1992;33:3351.
- (a) Freeman PK, Hutchinson LL. *Tetrahedron Lett*. 1849;1976:17;
- (b) Freeman PK, Hutchinson LL. *J Org Chem*. 1924;1980:45.
- Donohoe TJ, House D. *J Org Chem*. 2002;67:5015.
- Doi R, Shibuya M, Murayama T, Yamamoto Y, Iwabuchi Y. *J Org Chem*. 2015;80:401.
- Kumazaki H, Nakajima R, Bessho Y, Yokoshima S, Fukuyama T. *Synlett*. 2015;26:2131.
- Scholl M, Ding S, Lee CW, Grubbs RH. *Org Lett*. 1999;1:953.
- Mukherjee JP, Sil S, Pahari K, Chattopadhyay SK. *Synthesis*. 2016;48:1181.
- Sajiki H, Hattori K, Hirota K. *Chem Eur J*. 2000;6:2200.
- Okawa Y, Hideshima T, Ikeda H, et al. *Br J Haematol*. 2008;141:659.
- Thupari JN, Pinn ML, Kuhajda FP. *Biochem Biophys Res Commun*. 2001;285:217.
- Friis MB, Friborg CR, Schneider L, et al. *J Physiol*. 2005;567:427.
- Zimmermann C, Santos A, Gable K, et al. *Cell Rep*. 2013;5:1036.
- Régnacq M, Voisin P, Sere YY, et al. *BBRC*. 2016;477:33.
- Shpilka T, Welter E, Borovsky N, et al. *EMBO J*. 2015;34:2117.
- Lawrence DS, Zilfou JT, Smith CD. *J Med Chem*. 1999;42:4932.
- Shimazawa R, Ogawa Y, Morisaki N, Funabashi H, Kawaguchi A, Iwasaki S. *Chem Pharm Bull*. 1992;40:2954.
- Morisaki N, Funabashi H, Shimazawa R, et al. *Eur J Biochem*. 1993;211:111.
- Singh R, Kaushik S, Wang Y, et al. *Nature*. 2009;458:1131.
- Buchebner M, Pfeifer T, Rathke N, et al. *J Lipid Res*. 2010;51:2896.
- Kobayashi K, Ohte S, Ohshiro T, Ugaki N, Tomoda H. *Sci. Rep*. 2018;8:12099.