


Copper-Catalyzed Synthesis of α -Trifluoromethylthio-Substituted KetonesYangjie Huang,[†] Xing He,[†] Xiaoxi Lin, Mingguang Rong, and Zhiqiang Weng*

Department of Chemistry, Fuzhou University, Fuzhou 350108, China

Supporting Information

ABSTRACT: The CF₃S-substituted moiety serves as an important structural element in many bioactive molecules. A versatile copper catalyst that allowed for trifluoromethylthiolation of primary and secondary α -bromoketones is described. The reaction with readily available elemental sulfur and CF₃SiMe₃ afforded a broad scope and moderate to good yields of α -trifluoromethylthio-substituted ketones.

This procedure represents a very operationally simple yet powerful strategy for the synthesis of α -trifluoromethylthio-substituted ketones, a useful and versatile class of synthetic synthons.



α -Trifluoromethyl-substituted carbonyl compounds are valuable synthons for the preparation of numerous derivatives containing the CF₃ group that are of increasing interest in pharmaceutical, agricultural, and material sciences.¹ With their broad application considered, the development of novel and more efficient synthetic methods for α -CF₃-substituted carbonyl compounds has been the focus of extensive research interest.^{2,3}

However, whereas efficient methods for synthesis of α -CF₃-substituted carbonyl compounds have been reported,⁴ the development of preparation of α -CF₃S-substituted analogues is much less explored (Scheme 1). Compared to other fluorinated

reactions of ketone or benzoylacetate ethyl ester with trifluoromethylsulfenyl chloride to give the α -CF₃S-substituted ketones.^{11,12} Recently, the Shen and the Rueping groups independently described the electrophilic trifluoromethylthiolation of β -ketoesters to afford the corresponding α -trifluoromethylthiolated carbonyl compounds.^{6,13} Most recently, Li and Zard reported trifluoromethylthiolation of α -bromoketones with O-octadecyl-S-trifluorothiolcarbonate.¹⁴ However, some of these methods show modest substrate scope and require extremely toxic and corrosive CF₃SCl or electrophilic-type trifluoromethylthiolation reagent and are limited in their functional group compatibility.


Nonetheless, elegant work from the Qing group on copper-catalyzed oxidative trifluoromethylthiolation of aryl boronic acids with CF₃SiMe₃ and elemental sulfur^{10a,15} has raised anticipation that efficient trifluoromethylthiolation could be realized by using readily available and inexpensive catalysts and fluorinated reagents.

As part of our ongoing interest in the development of efficient methods for trifluoromethylthiolation¹⁶ and trifluoromethylselenolation,¹⁷ we herein report copper-catalyzed trifluoromethylthiolation of α -bromoketones with elemental sulfur and CF₃SiMe₃ to synthesize α -trifluoromethylthio-substituted ketones.

Our initial investigation commenced with the treatment of 2-bromo-1-*p*-tolylethanone **1a**, elemental sulfur, and CF₃SiMe₃ and KF with CuI (20 mol %)/phen (20 mol %) as catalyst in CH₂Cl₂ as the solvent at 40 °C (Table 1). The reaction occurred leading to the formation of the trifluoromethylthiolated ketone **2a** in 25% yield (entry 1).

To determine the efficient copper catalyst for this one-pot synthesis, we then examined a variety of copper salts. When CuSCN was used as a catalyst, a small improvement in the yield (39%) was observed (entry 2). When Cu(MeCN)₄PF₆, CuF₂,

Scheme 1. Methods for Synthesis of α -CF₃S-Substituted Ketones




alkyl groups (SF₅-, π_x = 1.23; CF₃O-, π_x = 1.04; CF₃-, π_x = 0.88), the CF₃S- group shows the highest lipophilicity value (π_x = 1.44).^{5,6} It is well-known that the introduction of trifluoromethylthio group into organic molecules such as pharmaceuticals and agrochemicals greatly enhances their bioavailability.^{7–10}

Therefore, new methods for incorporation of CF₃S-substituents into carbonyl compounds would be broadly useful in synthetic chemistry. Haas and Kolasa initially reported the

Received: May 6, 2014

Published: June 4, 2014

Table 1. Optimization of Copper-Catalyzed Trifluoromethylthiolation of 2-Bromo-4'-methylacetophenone^a



entry	[Cu]	ligand	solvent	temp (°C)	time (h)	yield ^b (%)
1	CuI	phen	CH ₂ Cl ₂	40	16	25
2	CuSCN	phen	CH ₂ Cl ₂	40	16	39
3	Cu(MeCN) ₄ PF ₆	phen	CH ₂ Cl ₂	40	16	63
4	CuF ₂	phen	CH ₂ Cl ₂	40	16	68
5	Cu(TFA) ₂	phen	CH ₂ Cl ₂	40	16	63
6	Cu(OTf) ₂	phen	CH ₂ Cl ₂	40	16	92
7			CH ₂ Cl ₂	40	16	2
8	Cu(OTf) ₂	bpy	CH ₂ Cl ₂	40	16	58
9	Cu(OTf) ₂	tmeda	CH ₂ Cl ₂	40	16	0
10	Cu(OTf) ₂	dmcda	CH ₂ Cl ₂	40	16	9
11	Cu(OTf) ₂	Me ₃ py	CH ₂ Cl ₂	40	16	1
12	Cu(OTf) ₂	phen	CH ₃ CN	40	16	5
13	Cu(OTf) ₂	phen	CH ₃ OH	40	16	0
14	Cu(OTf) ₂	phen	DMF	40	16	1
15	Cu(OTf) ₂	phen	DMF	80	16	1
16	Cu(OTf) ₂	phen	DMSO	40	16	1
17	Cu(OTf) ₂	phen	THF	40	16	9
18	Cu(OTf) ₂	phen	toluene	40	16	0
19	Cu(OTf) ₂	phen	CH ₂ Cl ₂	30	16	57
20	Cu(OTf) ₂	phen	CH ₂ Cl ₂	50	16	74
21	Cu(OTf) ₂	phen	CH ₂ Cl ₂	40	8	48

^aReaction conditions: [Cu] (0.010 mmol), ligand (0.010 mmol), 2-bromo-4'-methylacetophenone (0.050 mmol), CF₃SiMe₃ (0.20 mmol), S₈ (0.20 mmol), KF (0.20 mmol), solvent (1.0 mL), N₂; phen = 1,10-phenanthroline, bpy = 2,2'-bipyridine, tmeda = N,N,N',N'-tetramethylethylenediamine, dmcda = *trans*-N,N'-dimethyl-1,2-cyclohexanediamine, Me₃py = 2,4,6-trimethylpyridine. ^bThe yield was determined by ¹⁹F NMR spectroscopy with PhOCF₃ as internal standard.


and Cu(TFA)₂ (with phen as ligands) were employed, a moderate yield of product was obtained (entries 3–5). Notably, the use of Cu(OTf)₂ with phen as ligand produced the best results (92% yield; entry 6). It is also noteworthy that the products derived from the addition of the CF₃ to the carbonyl group^{1a,18} were not detected by using these conditions. Cu(OTf)₂ was therefore established as the preferred Cu source for the copper-catalyzed trifluoromethylthiolation. Only a trace amount of product was formed in the absence of a copper source and ligand (entry 7). The result clearly demonstrated the importance of both the catalyst and ligands in this transformation.

Although bipyridine has been shown previously to be a powerful ligand for copper-mediated trifluoromethylthiolation with aryl halides,^{16a} bpy was not so efficient as a ligand in this case, affording a less satisfactory yield (entry 8). Changing to other nitrogen ligands, such as tmeda, dmcda, or Me₃py, only retarded the reaction (entries 9–11). Additionally, the choice of solvent influenced the yield. The use of other solvents, e.g., CH₃CN, CH₃OH, DMF, DMSO, THF, or toluene (entries 12–18), was found to be detrimental. The effect of temperature was also investigated. Reactions performed either at lower temperatures (30 °C) or elevated temperature (50 °C) produced lower yields (entries 19 and 20). Furthermore,

reducing the reaction time to 8 h resulted in incomplete conversion (entry 21).

Having identified these optimal conditions, we next examined the substrate scope for this new reaction. A variety of aromatic α -bromoketones were surveyed to synthesize different trifluoromethylthiolated ketones (Scheme 2).

Scheme 2. Copper-Catalyzed Trifluoromethylthiolation of α -Bromoketones^a



^aReaction conditions: Cu(OTf)₂ (0.050 mmol), phen (0.050 mmol), α -bromoketones (0.25 mmol), CF₃SiMe₃ (1.0 mmol), S₈ (1.0 mmol), KF (1.0 mmol), CH₂Cl₂ (1.0 mL), N₂. Yields of isolated products are shown. ^bThe yield was determined by ¹⁹F NMR spectroscopy with PhOCF₃ as internal standard.


α -Bromoketones bearing methyl, dimethyl, and phenyl substitutions at the aromatic rings reacted with S₈ and CF₃SiMe₃ to give the products 2a–c, respectively, in 68–87% yields. The unfunctionalized aromatic α -bromoketones, 2-bromo-1-phenylethanone, and 2-bromo-1-(naphthalen-2-yl)ethanone also underwent trifluoromethylthiolation to afford the desired products 2d and 2e in good yields (72% and 75%, respectively).

Variation of electronic properties of the aryl substituents in the aromatic α -bromoketones had a dramatic effect on the reaction efficiency. α -Bromoketone substrates with a methoxy substituent at the *ortho*-, *meta*-, or *para*-position of the aromatic rings provided the corresponding products 2f–h in good yields (80%, 78% and 88%, respectively). Furthermore, α -bromoketone possessing a dimethylamino substituent in the *para*-position of the aryl groups also provided the desired product 2i, exclusively (70% yield). Additionally, α -bromoketone having nitro or nitrile substituents at the *meta*- or *para*-position of the aryl groups could also be transferred through this protocol, although reduced yields of the corresponding products 2j,k were observed (17% and 13%, respectively), with the trifluoromethylated alcohol resulting from the addition of CF₃ to the carbonyl group as side product. Notably, substrates bearing halogenated arenes were also accommodated and furnished the desired products 2l–n in moderate yields (20–69%), thereby providing possibilities for subsequent chemical transformations. Heteroaryl groups, such as coumarin group, could also be tolerated to give the trifluoromethylthiolated

product **2o** in 30% yield (^{19}F NMR). Interestingly, the trifluoromethylthiolation of an aliphatic α -bromoketone was possible and proceeded in excellent yield (**2p**, 85%), thus enhancing the scope of our reaction.

The scope was then extended to the use of secondary bromides (Scheme 3). The reaction with 2-bromopropiophe-

Scheme 3. Copper-Catalyzed Trifluoromethylthiolation of Secondary α -Bromoketones^a




^aReaction conditions: $\text{Cu}(\text{OTf})_2$ (0.075 mmol), phen (0.075 mmol), α -bromoketones (0.25 mmol), CF_3SiMe_3 (1.0 mmol), S_8 (1.0 mmol), KF (1.0 mmol), CH_2Cl_2 (1.0 mL), N_2 . Yields of isolated products are shown.

none, 4'-(benzyloxy)-2-bromopropiophenone, and 2-bromo-1-(3-chlorophenyl)-1-propanone led to the formation of desired products **2q–s** in good yields (77%, 80%, and 64%, respectively), albeit with the higher catalyst loading of CuI (30 mol %)/phen (30 mol %). Moreover, 2-bromo-1-(4-bromophenyl)-1-propanone smoothly underwent trifluoromethylthiolation to afford product **2t** in 71% yield with remarkable chemoselectivity, leaving the $\text{C}(\text{sp}^2)\text{--Br}$ bond untouched.

To prove the practicality of this protocol for large-scale synthesis, we prepared **2a** on a gram scale under the optimized reaction conditions (Scheme 4). The trifluoromethylthiolation of **1a** took place satisfactorily, affording the expected product **2a** in 72% yield.

Scheme 4. Gram-Scale Synthesis of α -Trifluoromethylthio-Substituted Ketones




In a synthetic application of the present reaction, we attempted to prepare alcohol and pyrazole derivatives, and the results are summarized in Scheme 5. The trifluoromethylthiolated ketone **2a** was treated with NaBH_4 to deliver alcohol **3** in 90% yield or alternatively reacted with PhMgBr to generate tertiary alcohol **4** in 72% yield.

The utility of the reaction is further illustrated by the synthesis of trifluoromethylthio-substituted pyrazoles, which is of great interest in medicinal chemistry.^{4b,19} Reaction of **2a** with dimethylformamide acetal followed by hydrazine gave the desired trifluoromethylthio-substituted pyrazole **5** in 88% overall yield, thus demonstrated herein that this new protocol provides a convenient methodology to access these biologically interesting scaffolds (Scheme 5).


In an attempt to find out whether any radical intermediates are formed in trifluoromethylthiolation, a radical scavenger, 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO), was added to

Scheme 5. Synthetic Utility of **2a**



the reaction under the optimized conditions (Scheme 6). Surprisingly, the reaction produced the desired product **2a** in

Scheme 6. Inhibition Experiment with TEMPO



only 19% yield together with the TEMPO- CF_3 adduct **6** in 56% yield by ^{19}F NMR analysis. This observation indicated that a CF_3 radical intermediate might be involved in the reactions.

In summary, a new approach to the direct construction of α -trifluoromethylthio-substituted ketones from readily available α -bromoketones, elemental sulfur, and CF_3SiMe_3 has been achieved using copper catalysis. This method provides a broad scope and moderate to good yields of the trifluoromethylthiolated products, and a variety of functional groups are compatible with these reaction conditions. The present reaction, therefore, is anticipated to be a powerful protocol for the synthesis of α -trifluoromethylthio-substituted ketones, which are important synthons for the preparation of numerous derivatives containing the CF_3S group. Further studies of this and related copper-catalyzed chemistry are in progress.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and spectral data for products and mechanistic study experiments. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: zweng@fzu.edu.cn.

Author Contributions

[†]These authors contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by National Natural Science Foundation of China (21072030), Research Fund for the Doctoral Program of Higher Education of China (No. 20123514110003), the SRF for ROCS, SEM, China (2012-1707), and Fuzhou University (022318, 022494).

■ REFERENCES

- (1) (a) Prakash, G. K. S.; Yudin, A. K. *Chem. Rev.* **1997**, *97*, 757–786. (b) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2004**, *104*, 6119–6146. (c) Ma, J.-A.; Cahard, D. *Chem. Rev.* **2008**, *108*, PR1–PR43. (d) Billard, T.; Langlois, B. R. *Eur. J. Org. Chem.* **2007**, 891–897. (e) Macé, Y.; Magnier, E. *Eur. J. Org. Chem.* **2012**, 2479–2494. (f) Shibata, N.; Matsnev, A.; Cahard, D. *Beilstein J. Org. Chem.* **2010**, *6* (65). (g) Ma, J.-A.; Cahard, D. *J. Fluorine Chem.* **2007**, *128*, 975–996.
- (2) (a) Miura, K.; Taniguchi, M.; Nozaki, K.; Oshima, K.; Utimoto, K. *Tetrahedron Lett.* **1990**, *31*, 6391–6394. (b) Mikami, K.; Tomita, Y.; Ichikawa, Y.; Amikura, K.; Itoh, Y. *Org. Lett.* **2006**, *8*, 4671–4673. (c) Sato, K.; Yuki, T.; Yamaguchi, R.; Hamano, T.; Tarui, A.; Omote, M.; Kumadaki, I.; Ando, A. *J. Org. Chem.* **2009**, *74*, 3815–3819. (d) Umemoto, T.; Ishihara, S. *J. Am. Chem. Soc.* **1993**, *115*, 2156–2164. (e) Itoh, Y.; Mikami, K. *Org. Lett.* **2005**, *7*, 649–651. (f) Itoh, Y.; Mikami, K. *Org. Lett.* **2005**, *7*, 4883–4885. (g) Itoh, Y.; Houk, K. N.; Mikami, K. *J. Org. Chem.* **2006**, *71*, 8918–8925. (h) Itoh, Y.; Mikami, K. *J. Fluorine Chem.* **2006**, *127*, 539–544. (i) Petrik, V.; Cahard, D. *Tetrahedron Lett.* **2007**, *48*, 3327–3330. (j) Tomita, Y.; Ichikawa, Y.; Itoh, Y.; Kawada, K.; Mikami, K. *Tetrahedron Lett.* **2007**, *48*, 8922–8925. (k) Noritake, S.; Shibata, N.; Nakamura, S.; Toru, T.; Shiro, M. *Eur. J. Org. Chem.* **2008**, 3465–3468. (l) Umemoto, T.; Adachi, K. *J. Org. Chem.* **1994**, *59*, 5692–5699. (m) Matoušek, V.; Togni, A.; Bizet, V.; Cahard, D. *Org. Lett.* **2011**, *13*, 5762–5765.
- (3) (a) Ma, J.-A.; Cahard, D. *J. Org. Chem.* **2003**, *68*, 8726–8729. (b) Kieltsch, I.; Eisenberger, P.; Togni, A. *Angew. Chem., Int. Ed.* **2007**, *46*, 754–757. (c) Noritake, S.; Shibata, N.; Nomura, Y.; Huang, Y.; Matsnev, A.; Nakamura, S.; Toru, T.; Cahard, D. *Org. Biomol. Chem.* **2009**, *7*, 3599–3604. (d) Deng, Q.-H.; Wadepohl, H.; Gade, L. H. *J. Am. Chem. Soc.* **2012**, *134*, 10769–10772.
- (4) (a) Pham, P. V.; Nagib, D. A.; MacMillan, D. W. C. *Angew. Chem., Int. Ed.* **2011**, *50*, 6119–6122. (b) Morandi, B.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 9085–9088. (c) Zhang, C.-P.; Wang, Z.-L.; Chen, Q.-Y.; Zhang, C.-T.; Gu, Y.-C.; Xiao, J.-C. *Chem. Commun.* **2011**, *47*, 6632–6634. (d) Novák, P.; Lishchynskiy, A.; Grushin, V. V. *J. Am. Chem. Soc.* **2012**, *134*, 16167–16170. (e) Deb, A.; Manna, S.; Modak, A.; Patra, T.; Maity, S.; Maiti, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 9747–9750.
- (5) Kirsch, P. *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*; Wiley-VCH: Weinheim, 2004; p xii, 308 p.
- (6) Bootwicha, T.; Liu, X.; Pluta, R.; Atodiresei, I.; Rueping, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 12856–12859.
- (7) Filler, R. *Biomedical Aspects of Fluorine Chemistry*; Kodansha: Tokyo, 1982.
- (8) Yagupol'skii, L. M.; Ilchenko, A. Y.; Kondratenko, N. V. *Russ. Chem. Rev.* **1974**, *43*, 32–47.
- (9) (a) Tlili, A.; Billard, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 6818–6819. (b) Wang, H.; Vicic, D. A. *Synlett* **2013**, *24*, 1887–1898.
- (10) (a) Chen, C.; Xie, Y.; Chu, L.; Wang, R.-W.; Zhang, X.; Qing, F.-L. *Angew. Chem., Int. Ed.* **2012**, *51*, 2492–2495. (b) Baert, F.; Colomb, J.; Billard, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 10382–10385. (c) Alazet, S.; Zimmer, L.; Billard, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 10814–10817. (d) Zhang, C.-P.; Vicic, D. A. *J. Am. Chem. Soc.* **2012**, *134*, 183–185. (e) Zhang, C.-P.; Vicic, D. A. *Chem.—Asian J.* **2012**, *7*, 1756–1758. (f) Chen, C.; Chu, L.; Qing, F.-L. *J. Am. Chem. Soc.* **2012**, *134*, 12454–12457. (g) Yang, Y.-D.; Azuma, A.; Tokunaga, E.; Yamasaki, M.; Shiro, M.; Shibata, N. *J. Am. Chem. Soc.* **2013**, *135*, 8782–8785. (h) Deng, Q.-H.; Rettenmeier, C.; Wadepohl, H.; Gade, L. H. *Chem.—Eur. J.* **2014**, *20*, 93–97. (i) Vinogradova, E. V.; Müller, P.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2014**, *53*, 3125–3128.
- (11) Bayreuther, H.; Haas, A. *Chem. Ber.* **1973**, *106*, 1418–1422.
- (12) Kolasa, A. *J. Fluorine Chem.* **1987**, *36*, 29–40.
- (13) (a) Shao, X.; Wang, X.; Yang, T.; Lu, L.; Shen, Q. *Angew. Chem., Int. Ed.* **2013**, *52*, 3457–3460. (b) Wang, X.; Yang, T.; Cheng, X.; Shen, Q. *Angew. Chem., Int. Ed.* **2013**, *52*, 12860–12864.
- (14) Li, S.-G.; Zard, S. Z. *Org. Lett.* **2013**, *15*, 5898–5901.
- (15) Chu, L.; Qing, F.-L. *Acc. Chem. Res.* **2014**, *47*, 1513–1522.
- (16) (a) Weng, Z.; He, W.; Chen, C.; Lee, R.; Tan, D.; Lai, Z.; Kong, D.; Yuan, Y.; Huang, K.-W. *Angew. Chem., Int. Ed.* **2013**, *52*, 1548–1552. (b) Tan, J.; Zhang, G.; Ou, Y.; Yuan, Y.; Weng, Z. *Chin. J. Chem.* **2013**, *31*, 921–926. (c) Zhu, P.; He, X.; Chen, X.; You, Y.; Yuan, Y.; Weng, Z. *Tetrahedron* **2014**, *70*, 672–677.
- (17) (a) Chen, C.; Hou, C.; Wang, Y.; Hor, T. S. A.; Weng, Z. *Org. Lett.* **2014**, *16*, 524–527. (b) Chen, C.; Ouyang, L.; Lin, Q.; Liu, Y.; Hou, C.; Yuan, Y.; Weng, Z. *Chem.—Eur. J.* **2014**, *20*, 657–661.
- (18) Singh, R. P.; Shreeve, J. M. *Tetrahedron* **2000**, *56*, 7613–7632.
- (19) (a) Benbow, J. W.; Lou, J.; Pfefferkorn, J. A.; Tu, M. M. US2010/0063063, 2010. (b) Hoffmann, M. G.; Helmeke, H.; Willms, L.; Auler, T.; Kehne, H.; Hills, M.; Feucht, D. US2005/0209106, 2005. (c) Bargamova, M. D.; Motsishkate, S. M.; Knunyants, I. L. *Bull. Russ. Acad. Sci.* **1991**, *39*, 2338–2344.