This article was downloaded by: [University of North Carolina] On: 07 October 2014, At: 02:44 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn19</u>

Amino Acid Nucleic Acids: Synthesis and Hybridization Properties of a Novel Class of Antisense Oligonucleotides

Kanda S. Ramasamy ^a & Wilfried Seifert ^a ^a Research Division, ICN Pharmaceuticals, Inc., 3300 Hyland Avenue, Costa Mesa, CA, 92626 Published online: 16 Aug 2006.

To cite this article: Kanda S. Ramasamy & Wilfried Seifert (1997) Amino Acid Nucleic Acids: Synthesis and Hybridization Properties of a Novel Class of Antisense Oligonucleotides, Nucleosides and Nucleotides, 16:7-9, 1519-1522, DOI: <u>10.1080/07328319708006220</u>

To link to this article: <u>http://dx.doi.org/10.1080/07328319708006220</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

AMINO ACID NUCLEIC ACIDS: SYNTHESIS AND HYBRIDIZATION PROPERTIES OF A NOVEL CLASS OF ANTISENSE OLIGONUCLEOTIDES

Kanda S. Ramasamy* and Wilfried Seifert

Research Division, ICN Pharmaceuticals, Inc., 3300 Hyland Avenue, Costa Mesa, CA 92626

Abstract: Oligonucleotides containing novel phoshoramidite 12 were synthesized and studied for their hybridization properties for the first time. Interestingly, these modified oligonucleotides showed a remarkable resistance to exonuclease.

Oligonucleotides that specifically recognize messenger RNA¹ present unique opportunities for the treatment of viral diseases, cancer, and for the study of genetic disorders. In order to be pharmacologically useful, the oligonucleotides must have resistance to cellular nucleases and penetrate through cell membrane. To meet these properties several derivatives have been synthesized² and examined their biophysical characteristics.

Our interest to design a novel class of antisense oligonucleotides has lead us to envision amino acid nucleic acids (AANA) (i.e., molecules where the individual nucleobases are linked to an amino alcohol through an amide bond and phosphodiester backbone). This isosteric oligonucleotide analogue derived from serinol is shown in Figure 1, (2). In this novel class of molecule, we believe, the formation of an intramolecular hydrogen bonding between the amide "NH" of the acetamide tether and the phosphate "O" is necessary for AANA to adopt a constrained conformation (see Figure 1, structure 2). Thus, the constrained monomers when incorporated into oligonucleotides, may have a force greater than a single hydrogen bond and favor the formation of binding competent nucleic acids. Furthermore, the hydrogen bonding could neutralize the negative of the phosphodiester backbone and this neutralization is expected to enhance the uptake of the AANAs relative to DNAs, rendering them potentially more useful as therapeutic agents.

In this note, we report the synthesis of a novel oligonucleotide building block 12 and biophysical properties of oligonucleotides containing 12. Optically active amino acid nucleoside³ analogue bearing thymine was prepared from <u>O</u>-benzyl-D-serine by the route shown in Scheme 1. Thymine acetic acid (7) was prepared by the literature procedure.⁴ Treatment of <u>O</u>-benzyl-D-serine with di-*tert*-butyl dicarbonate provided N^{α}-BOC-<u>O</u>-benzyl-D-



Figure 1



Scheme 1^a

^a(a) DBDC/THF/water (93%); (b) i. IBCF/TEA/THF; ii.NaBH₄THF/Water; (82%); (c) (IbO)₂O/Py (84%); (d) i.TFA/CH₂Cl₂, ii. EDC/NMM/HOBt/7/DMF (94%); (c) Pd(OH)₂/Cyclohexene (91%); (f) DMTCl/TEA/Py (88%); (g) 1N NaOH/THF/MeOH (99%); (h) EtN(iPr)₂/P(Cl)(iPr)₂OCH₂CH₂CN (2 equiv)/CH₂Cl₂ (83%). DBDC = di-*tert*-butyl dicarbonate; IBCF = isobutyl chloroformate; NMM = N-methylmorpholine; HOBt = 1-hydroxybenzotriazole; T = Thymine.

serine (4) in quantitative yield. Reduction of the α -carboxyl group of 4 using mixed anhydride method⁵ furnished the corresponding alcohol (5). Exposure of the alcohol (5) to isobutyric anhydride in dry pyridine for 8 h, followed by usual workup produced the diprotected alcohol (6). Removal of the t-BOC group with trifluoroacetic acid followed by coupling of the TFA salt with thymine acetic acid (7) in the presence of 1-ethyl-3-(3dimethylaminopropyl)-carbodiimide hydrochloride (EDC), as coupling reagent, gave the

Sequences	Tm(°C)		ΔTm /modification (°C)	
	DNA	RNA	DNA	RNA
5' GAA AGG AAG CGG AGA GAT 3'				
5' ATC TCT CCG CTT CCT TTC 3'	58.29	66.8		
5' ATC TCT CCG CTT CCT TtC 3'	60.34	69.03	+2.05	+2.23
5' ATC TCT CCG CTT CCt ttC 3'	56.79		- 0.50	
5' AtC TCT CCG CTT CCT TTC 3'	60.52	68.90	+2.23	+2.1
5' ATC TCT CCG C(T CCT TTC 3'	48.45	58.02	- 9.84	- 8.78
5' AtC TCT CCG CTT CCT TtC 3'	56.85	65.8	~ 0.72	- 0.50

Table 1: Tm values Of Modified Oligonucleotides

The letter "t" denotes modified thymidine.

amino alcohol nucleoside (8). Hydrogenation of 8 with $Pd(OH)_2$ in methanol containing cyclohexene furnished 1-Q-isobutyryl-2-(thyminylacetyl)amino-3-hydroxy-D-propanol (9).⁶ Dimethoxytritylation⁷ of 9 followed by the removal of the isobutryl group in 10 with 1N NaOH solution and phosphitylation⁸ of 11 afforded amino acid nucleic acid amidite synthom 12 as colorless powder.

Incorporation of the building block 12 into oligonucleotide sequences was accomplished using ABI 394 DNA synthesizer and protocol,⁹ and the coupling efficiency was found to be higher than 99%. Enzymatic degradation and subsequent HPLC analysis of the modified oligonucleotides indicated the expected ratios of the nucleoside components.¹⁰ The binding behaviour of these modified oligonucleotides was assayed by examining their ultraviolet (UV) absorbance verses temperature profiles.¹¹ An 18mer oligonucleotide (5'-ATCTCTCCGCTTCCTTTC-3') was modified with 12 at different locations and hybridized to complementary DNA or RNA. Incorporation of 12 at the 3'-end of oligonucleotide increased the duplex stability by a Δ Tm of 2.05 °C per modification. The same type of duplex stability was observed when 12 is substituted at 5'-end too. On the contrary, incorporation of 12 in the middle of the duplex caused a significant reduction in binding (Δ Tm= -9.84 °C per modification). The presence of 12 at 3'-end as well as at 5'-end had very little effect ($\Delta Tm = -$ 0.72 °C) on duplex stability. When the fully modified (4',5' linked) poly T (AANA) 18mer was heated with its complement poly dA (3',5' linked), a steady increase in absorbance (260 nm) is observed but no cooperative transitions are detectable. This confirms the lack of association of poly T (AANA) having 4',5' linkage with poly dA in aqueous solution. Interestingly enough, incorporation of three modifications at the 3'-end of the oligonucleotides, increased their exonuclease stability¹² by 50-fold than natural DNA. On the other hand, 5'-modified AANAs do not show any improved exonuclease property.

It is evident from this study that 4',5' linked AANA forms a weaker duplex than normal DNA. The weaker binding of oligonucleotides containing D-serine amino acid is an indicative that the monomer 12 is not adopting a constrained conformation that of ribose sugar puckering. Although oligonucleotides containing amino acid nucleic acid building block exhibited better stability towards nucleases, their inability to bind strongly especially under physiological conditions, may precludes their use as antisense compounds.

REFERENCES

- (a) Zamecnik, P. C.; Stephenson, M. L. Proc. Natl. Acad. Sci. U. S. A. 1978, 75, 280.
 (b) Goodchild, J. Bioconjugate Chem. 1990, 1, 165.
- 2. Ramasamy, K.; Seifert, W. Bioorg. & Med. Chem. Lett., 1996, 6, 1799 and references cited there in.
- 3. Nucleobases attached to serinol via acetamide linker. See Fig. 1 and structure (2).
- 4. Egholm, M.; Buchardt, O.; Nielsen, P. E.; Berg, R. H. J. Am. Chem. Soc. 1992, 114, 1895.
- 5. Ramasamy, K.; Olsen, R. K.; Emery, T. Synthesis 1982, 42.
- 6. All the compounds synthesized were confirmed by ¹HNMR and elemental analysis.
- 7. Schaller, H.; Weimann, G.; Lerch, B.; Khorana, H. G. J. Am. Chem. Soc. 1963, 85, 3821.
- 8. Karpyshev, N. N. Russ. Chem. Rev. 1988, 57, 886.
- 9. Atkinson, T.; Smith, M. in Oligonucleotide Synthesis: A practical Approach; Gait, M. J., Ed.; IRL: Oxford; 1984; pp 35-81.
- 10. Oligonucleotides were digested with a mixture of spleen phosphodiesterase, snake venom phosphodiesterase, and bacterial alkaline phosphatase to provide individual nucleosides which were analyzed by HPLC.
- 11. Freier, S. M.; Albergo, D. D.; Turner, D. H. Biopolymers 1982, 22, 1107.
- 12. For experimental procedure see: Svendsen, M. L.; Wengel, J. Tetrahedron 1993, 49, 11341.