



Communication

Subscriber access provided by Caltech Library Services

Completely regioselective direct C–H functionalization of benzo[b]thiophenes using a simple heterogeneous catalyst

Dan-Tam Daniel Tang, Karl D Collins, and Frank Glorius

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/ja403130g • Publication Date (Web): 06 May 2013 Downloaded from http://pubs.acs.org on May 7, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Completely regioselective direct C–H functionalization of benzo[*b*]thiophenes using a simple heterogeneous catalyst

Dan-Tam D. Tang, Karl D. Collins and Frank Glorius*

Westfälische Wilhelms-Universität Münster, Organisch-Chemisches Institut, Corrensstraße 40, 48149, Münster, Germany. *Supporting Information Placeholder*

ABSTRACT: The first completely selective C3 C–H arylation of benzo[*b*]thiophenes is reported, demonstrating previously unexploited reactivity of palladium. Benzo[*b*]thiophenes are coupled with readily available aryl chlorides using a ligand free, dual catalytic system of heterogeneous Pd/C, and CuCl. The reaction is operationally simple, insensitive to air and moisture, and provides valuable products with complete selectivity. Significant investigation into the nature of the active catalytic species and mechanistic considerations are discussed.

Despite significant advances in homogeneous catalysis in recent times, this knowledge has at large proved resistant to development in an industrial environment, with the majority of industrial catalytic processes continuing to employ heterogeneous catalysts.¹ The high costs associated with homogenous catalysis due to the use of unrecoverable metals and ligands, as well as practical limitations with regard to catalyst handling, recyclability and the separation of the catalyst from products impede the transfer of advances in homogeneous catalysis to large scale industrial processes. In contrast, many of these limitations can be obviated by use of heterogeneous catalysts, and consequently, the development of such processes is desirable.²⁻⁵

Arylated heterocycles are valuable structural motifs and are present in many natural products and biologically active molecules.⁶ Commonly these are prepared *via* transition metal catalyzed cross-coupling of functionalized heterocycles and suitable coupling partners,⁷ though more sustainable methods are desirable. Recently, significant development of the field of C–H activation has enabled the direct arylation of heterocycles,⁸ circumventing the need for pre-functionalization. Unfortunately though, multicomponent and expensive catalyst systems are often required, and directing groups to control regioselectivity are necessary. Alternatively, many metal catalyzed direct C–H functionalizations exploiting the inherent reactivity of heterocycles are established,^{8c,9} providing viable alternatives, though methods that selectively functionalize the less reactive C–H bonds without recourse to directing groups,¹⁰ or exploiting the steric bulk of ligands,¹¹ substituents¹² or additives¹³ remain underdeveloped.

For valuable benzo[*b*]thiophenes motifs^{14,15} the transition metal catalyzed direct arylation of the C2-position is well established,^{9c} though efficient methods for the more challenging direct C3-arylation remain more elusive with only three reports to date (Scheme 1a-c).¹⁶⁻¹⁸ Of the three reports, two provide a single example, and other limitations include expensive catalyst systems, large excesses of reagents or coupling partners, limited availabil-

ity of coupling partners, and super-stoichiometric use of oxidants. Limitations in the C2/C3 selectivity of these methods are also apparent.

In the course of investigating novel reactivity of heterogeneous catalysts¹⁹ we have established an operationally simple, *complete-ly* selective C3-arylation of unsubstituted benzo[*b*]thiophenes, addressing many of the limitations of existing methods. Herein we describe the coupling of benzo[*b*]thiophenes with inexpensive and widely available aryl chlorides, using Pd/C with CuCl as co-catalyst (Scheme 1d). The reaction is ligand free, scalable, insensitive to air and moisture, and represents reactivity for Pd/C beyond its typical application in cross-coupling and hydrogenation reactions.^{4,20} Furthermore, our investigations suggest the active catalytic species is likely to be heterogeneous in nature.

We initially found that reaction of benzo[b]thiophene 1 and chlorobenzene 2 in the presence of Pd/C (K-0219 from Heraeus) gave 3-phenylbenzo[b]thiophene 3 in 18% isolated yield with excellent C3/C2 selectivity (Table 1, entry 1). Solvent screening and standard optimization of catalyst loading, temperature, ratio of reagents etc. gave a maximum yield of 60% (entry 5), though complete C2/C3 selectivity was maintained. In an attempt to increase the yield of reaction further, we proposed a number of additives to enhance reactivity. Pivalic acid, a common additive in C-H activation chemistry,⁸ proved detrimental to reactivity (entry 2), as did silver salts which were proposed to facilitate the potential formation of cationic palladium species as proposed by Itami and Studer (entries 3 and 4).^{16,17} Pleasingly, in attempts to facilitate reaction by Lewis acid activation of 1, we established a number of zinc, iron, gold and copper salts enhanced reactivity, with CuCl proving most efficacious. A catalyst loading of 9.4 mol% was determined to be optimal. Interestingly, CuBr gave a similar result whereas CuI retarded the reaction completely (entries 10 and 11). Control experiments demonstrated that base is essential for reactivity, copper did not catalyze the reaction in the absence of palladium, and no reaction was observed in the absence of copper and palladium (entries 12-14). We also assessed numerous Pd/C sources and observed significant variation in activity, though 'standard' Pd/C from Heraeus and Sigma-Aldrich both showed excellent reactivity. See SI for all additional experimental data.

In addition to Pd/C, the reactivity of several alternative heterogeneous catalysts under identical conditions was also investigated. Several supported palladium systems were investigated, as well as other transition metals. In all instances no appreciable reactivity was observed, highlighting the unique catalytic activity of Pd/C with respect to this transformation (see SI).





Table 1. Optimization of Reaction Conditions

$\begin{array}{c c} & & & \\ \hline \\ & & \\ & \\ & \\ & \\ & \\ & \\ &$			
Entry	Additive (mol%)	Yield ^a	C3/C2 ^b
1	-	30%	>98:2
2	PivOH (10)	22%	>99:1
3	$Ag_2CO_3(5)$	n. r.	-
4	$AgSbF_{6}(5)$	traces	>99:1
5 ^{<i>c</i>}	-	60%	>99:1
6 ^{<i>c</i>}	$\operatorname{AuCl}_{3}(5)$	64%	>99:1
7^c	CuCl (5)	78%	>99:1
8 ^c	CuCl (10)	88%	>99:1
9 ^c	CuCl (15)	69%	>99:1
10^{c}	CuBr (5)	77%	>99:1
11^{c}	CuI (5)	n. r.	-
$12^{c,d}$	CuCl (10)	n. r.	-
$13^{c,d}$	-	n. r.	-
14 ^{<i>c</i>,<i>e</i>}	CuCl (10)	n. r.	-

General conditions: **1** (0.300 mmol), **2** (1.0 equiv.), Cs₂CO₃ (1.1 equiv.), Pd/C (9.4 mol%), 1,4-dioxane (1.5 ml), 150 °C, 48 h. ^{*a*} Yields determined by GC of crude reaction mixture; mesitylene as internal standard. ^{*b*} C3/C2 ratios determined by GC-MS of crude reaction mixture. ^{*c*} **1** (0.300 mmol), **2** (2.0 equiv.), 1,4-dioxane (1.5 ml). ^{*d*} W/o Pd/C. ^{*e*} W/o base.

With optimized conditions established we explored the scope of the reaction (Table 2). Electron-rich and electron-neutral aryl chlorides gave reaction products in typically excellent yield. Sterically encumbered aryl chlorides also proceeded in excellent yield, though extended reaction times were required; the reaction of 2-chloro-*m*-xylene is of particular note. Reaction with electrondeficient aryl halides including nitro- and ester-substituted systems proceeds, though the rate of reaction was significantly reduced.²¹ Interestingly, electron deficient systems bearing only fluoride heteroatoms proceeded in high yield and acceptable reaction time, indicating that the electron deficient nature of the aryl chloride alone is not significantly detrimental to reactivity. Furthermore, C5-substituted benzo[b]thiophenes were amenable to the reaction conditions and were coupled in very high yield and pleasingly, reaction of 1 with 1,4-dichlorobenzene gave the *bis*-benzo[b]thiophene adduct 15 in 41% yield. The scalability of the reaction has also been demonstrated, with reaction to give 3 undertaken on a 3 mmol scale. Unfortunately, efficient recycling of the catalyst has not been established.

Table 2. Reaction Scope



General conditions: Benzo[*b*]thiophene (0.300 mmol), ArCl (2.0 equiv.) Pd/C (9.4 mol%), Cs_2CO_3 (1.1 equiv.), 1,4-dioxane (1.5 ml), 48-72 h. Yields given are isolated. C3/C2 ratios determined by GC-MS of crude reaction mixture. ^{*a*} 3 mmol scale, reaction time 72 h, GC yield. ^{*b*} Product isolated as mixture with corresponding homo-coupled ArCl. Yield given refers to denoted compound. ^{*c*} Reaction time 96 h. ^{*d*} ArCl (0.300 mmol), **1** (3.0 equiv.).

The application of this method to other heterocycles has provided limited success to date. Only thiophenes have demonstrated any notable reactivity, though poor C3/C2 selectivity is observed. Pleasingly though, we have demonstrated the selective nature of this reaction in an intermolecular competition experiment. Using the standard reaction conditions, a 1:1:1 mixture of benzo[*b*]thiophene **1**, 2-*n*-butylthiophene and pyridine showed almost exclusive reactivity for benzo[*b*]thiophene **1**, suggesting this method is appropriate for application to more complex substrates (Scheme 2).

60





The hetero- or homogeneous nature of the active catalyst in many reactions is often highly contentious, and specifically with regard to Pd/C the active species is often thought to be homogeneous following leaching of palladium into solution.²² Consequently, we have undertaken significant effort to determine the nature of the active species in our system. Considering both the high temperature of the reaction and the presence of aryl chlorides – which in given examples are known to leach palladium from carbon supports *via* an oxidative addition process²³ – it is highly notable that we have demonstrated that the active catalyst is very likely to be heterogeneous in nature.

Although no current methods evaluating the homo/heterogeneous nature of an active catalyst can be considered completely definitive,²² the results of numerous experimental tests and literature precedent for the conclusions drawn has led us to propose that the active catalytic species is highly likely to be heterogeneous in nature. Key experiments include: i) Two 3-phase tests,²² whereby one reagent (the aryl chloride or benzo[b]thiophene) is immobilized on a solid support, showed that active homogeneous catalytic species are not generated under these reaction conditions. ii) Standard Hg(0) poisoning and hot ² iii) The filtration tests are positive for heterogeneous catalysis.²² introduction of PPh₃ (10 mol%), an established poison for heterogeneous catalysts results in significant inhibition of reactivity. Complete inhibition of the catalyst is typically expected with substoichiometric quantities of PPh₃ (with respect to the catalyst) due to irreversible binding of the phosphine to the surface, though in this instance we propose that the high reaction temperature results in dynamic rather than irreversible binding of the phosphine, and thus trace reaction is observed. iv) Numerous comparable homogeneous Pd catalysts showed no reactivity. Reactivity of PdCl₂(PPh₃)₂ and Pd(PPh₃)₄ was observed, though resulted exclusively in C2-arylation. v) An analysis of the reaction mixture by X-ray fluorescence spectroscopy (XRF) after filtration showed palladium present at a concentration of <4 ppm, corresponding to 0.01% of the initial palladium loading. Additional experiments, clarification of results, and further circumstantial evidence are provided in the supporting information.

The results of our investigation strongly suggest a heterogeneously catalyzed reaction. The 3-phase tests, and experiments demonstrating that site-selectivity is reversed when employing active homogeneous Pd(0) are particularly compelling. This change in selectivity highlights the unique reactivity possible when employing a heterogeneous catalyst. It should be noted a short induction period was observed when investigating the kinetic progress of the reaction. We propose this could be accounted for by the initial reduction/modification of the catalyst surface, to generate the active catalyst. Alternatively, leaching of palladium into solution could account for the induction period, though should this be the case, our results suggest likely agglomeration of leached palladium, and formation of a new heterogeneous active species rather than generation of a homogeneous active species.^{22a}

In addition to our empirical investigation into the physical nature of the active catalyst, we have also evaluated possible mechanistic pathways. A number of viable homogeneous mechanistic pathways have been considered, though none is satisfactory (Figure 1, see SI for associated experimental data): i) Direct C–H

activation/palladation at C3 to give 18 followed by reductive elimination. As described by Hartwig, homogeneous C-H activation mechanisms are highly amenable to investigation using kinetic isotope studies.²⁴ In this instance, a 'homogenous type' C-H activation mechanism is directly refuted by the absence of a KIE in the appropriate kinetic and competition experiments proposed by Hartwig. Furthermore, palladation at the more acidic C2position would be expected for typical C-H activation mechanisms.⁸ ii) Palladation at C2 generating cationic species 19, followed by aryl-migration and elimination of palladium as proposed by Itami.¹⁶ This mechanism was deemed improbable as unlike in the report of Itami, when silver salts were employed to promote formation of cationic palladium species via sequestering of halides, retardation of reaction was observed. Furthermore, no reaction of 3-methylbenzo[b]thiophene 22 under identical reaction conditions was observed. We suggest it is likely arylation at C2 would be expected in this instance via the reductive elimination of 19b, as the migratory pathway is shut down. We appreciate subtle steric effects may also account for the lack of reactivity. iii) A Heck type mechanism. This was ruled out due to the mechanistic requirement for syn β -hydride elimination, thus requiring C-C bond rotation of cyclic intermediate 20 which is not possible. Alternatively, base catalyzed elimination of the palladium driven by re-aromatization may be considered, though the selectivity observed in the hypothetical migratory insertion to benzo[b]thiophene is contrary to that expected.²⁵ iv) Electrophilic palladation at C3 to give 21, followed by reductive elimination and re-aromatization. Reaction of 2-methylbenzo[b]thiophene 23 and 3-methylbenzo[b]thiophene 22 under the standard reaction conditons gave 11% and 0% yield, respectively. Although electrophilic arylation in a manner analogous to indole would give preferential reactivity at C3, the complete inhibition of reactivity with 22, rather than a shift in selectivity is unlikely; additionally, the relatively small and electronically neutral substitution at C2 as present in 22, would not be expected to inhibit reactivity to such an extent using simple such simple catalytic systems.²⁶ As a result of the discussed mechanistic considerations, we propose that traditional mechanisms associated with homogeneous catalysis are not *directly* applicable in this instance, and consequently that the supported nature of the catalyst may have a direct impact on the regioselectivity of the reaction.





In summary, we have described the first heterogeneously catalyzed C–H arylation of benzo[b]thiophenes, demonstrating novel reactivity mediated by a simple palladium catalyst. The reaction appears intrinsically selective for the typically less reactive C3position, requiring no additional ligands or directing groups. Cheap and widely available catalysts and coupling partners are employed, and the reaction is both operationally simple, and tolerant of air and water. Our mechanistic investigation suggests typical homogeneous reaction mechanisms are unlikely to be in operation. Further studies relating to the interaction of the substrate and catalyst, the mechanism of reaction, and the role of CuCl are ongoing within our laboratory.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures and full spectroscopic data for all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

glorius@uni-muenster.de

ACKNOWLEDGMENT

This paper is dedicated to Prof. Andreas Pfaltz, a great teacher and pioneer of asymmetric catalysis, on the occasion of his 65th birthday. Generous financial support by the Deutsche Forschungsgemeinschaft (SFB 858) and the Fonds der Chemischen Industrie (D.-T. D. T.) is gratefully acknowledged. We thank Prof. Dr. Uwe Karst and Christoph Wehe for TXRF analyses, Karin Gottschalk for experimental support and Dr. Rainer Kiemel (Heraeus) and Dr. Sébastien Parisel and Dr. Steve Hawker (both Johnson Matthey) for stimulating discussions and for the kind donation of catalyst samples (Johnson Matthey).

REFERENCES

(1) (a) Fine chemicals Through Heterogeneous Catalysis; Sheldon, R. A.; van Bekkum, H., Eds.; Wiley-VCH, Weinheim, **2001**. (b) Asymmetric Catalysis on Industrial Scale, Blaser, H. U.; Schmidt, E., Eds.; Wiley-VCH, Weinheim, 2004.

(2) For illustrative reviews on asymmetric heterogeneous catalysis see: (a) Heitbaum, M.; Glorius, F.; Escher, I. Angew. Chem., Int. Ed. 2006, 45, 4732. 2. (b) Mallat, T.; Orglmeister, E.; Baiker, A. Chem. Rev. 2007, 107, 4863. (c) Trindade, A. F.; Gois, P. M. P.; Afonso, C. A. M. Chem. Rev. 2009, 109, 418. (d) Wang, Z.; Chen, G.; Ding, K. L. Chem. Rev. 2009, 109, 322. For a recent example, see: (e) Gross, E.; Liu, J. H.; Alayoglu, S.; Marcus, M. A.; Fakra, S. C.; Toste, F. D.; Somorjai, G. A. J. Am. Chem. Soc. 2013, 135 3881.

(3) Thayer, A. Chem. Eng. News 2005, 83, 55.

(4) Review on heterogeneous palladium catalyzed couplings: Yin, L.; Liebscher, J. Chem. Rev. 2007, 107, 133.

(5) For selected reviews on the application of nanocatalysts see: (a) Polshettiwar, V.; Luque, R.; Fihri, A.; Zhu, H.; Bouhrara, M.; Basset, J.-M. *Chem. Rev.* **2011**, *111*, 3036. (b) Ranganath, K. V. S.; Glorius, F. *Catal. Sci. Technol.* **2011**, *1*, 13. (c) Schätz, A.; Reiser, O.; Stark, W. J. *Chem. Eur. J.* **2010**, *16*, 8950. (d) Shylesh, S.; Schünemann, V.; Thiel, W. R. *Angew. Chem., Int. Ed.* **2010**, *49*, 3428. (e) Lu, A.-H.; Salabas, E. L.; Schüth, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 1222. (f) Astruc, D.; Lu, F.: Aranzaes, J. R. *Angew. Chem., Int. Ed.* **2005**, *44*, 7852. For a recent example, see: (h) Gross, E.; Liu, J.H-C.; Toste, F. D.; Somorjai, G.A. *Nature Chem.* **2012**, *4*, 947.

(6) (a) Nicolaou, K. C.; Hale, C. R. H.; Nilewski, C.; Ioannidou, H. A. *Chem. Soc. Rev.* **2012**, *41*, 5185. (b) McGlacken, G. P.; Bateman, L. M. *Chem. Soc. Rev.* **2009**, *38*, 2447. (c) *Bioactive Heterocy-clic Compound Classes: Pharmaceuticals and Agrochemicals*; Lamberth, C.; Dinges, J., Eds.; Wiley-VCH, New York, **2012**.

(7) For reviews on transition metal catalyzed cross-coupling of heterocycles see: (a) Jana, R.; Pathak, T. P.; Sigman, M. S. *Chem. Rev.* **2011**, *111*, 1417. (b) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359.

(8) For recent reviews on C-H activation see: (a) Beccalli, S.; Broggini, G.; Martinelli, M.; Sottocornola, S. Chem. Rev. 2007, 107, 5318. (b) Li, B.-J.; Yang, S.-D.; Shi, Z.-J. Synlett 2008, 7, 949. (c) Ackermann, L.; Vicente, R.; Kapdi, A. R. Angew. Chem., Int. Ed. 2009, 48, 9792. (d) Chen, X.; Engle, K. M.; Wang, D.-H.; Yu, J.-Q. Angew. Chem., Int. Ed. 2009, 48, 5094. (e) Lautens, M.; Thansandote, P. Chem. Eur. J. 2009, 15, 5874. (f) Kitamura, T. Eur. J. Org. Chem. 2009, 1111. (g) Giri, R.; Shi, B.-F.; Engle, K. M.; Maugel, N.; Yu, J.-

Q. Chem. Soc. Rev. 2009, 38, 3242. (h) Daugulis, O.; Do, H.-Q.; Shabashov, D. Acc. Chem. Res, 2009, 42, 1074. (i) Jazzar, R.; Hitce, J, Renaudat, A.; Sofack-Kreutzer, J.; Baudoin, O. Chem. Eur. J. 2010, 16, 2654. (j) Lyons, T. W.; Sanford, M. S. Chem. Rev. 2010, 110, 1147. (k) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Chem. Rev. 2010, 110, 624. (1) Satoh, T.; Miura, M. Chem. Eur. J. 2010, 16, 11212. (m) Cho, S. H.; Kim, J. Y.; Kwak, J.; Chang, S. Chem. Soc. Rev. 2011, 40, 5068. (n) Wencel-Delord, J.; Dröge, T.; Liu, F.; Glorius, F. Chem. Soc. Rev. 2011, 40, 4740. (o) Newhouse, T.; Baran, P. S. Angew. Chem., Int. Ed. 2011, 50, 3362. (p) Ackermann, L. Chem. Rev. 2011, 111, 1315. (q) McMurray, L.; O'Hara, F;. Gaunt, M. J. Chem. Soc. Rev. 2011, 40, 1885. (r) Yeung, C. S.; Dong, V. M. Chem Rev. 2011, 111, 1215. (s) White, M. C. Science 2012, 335, 807. (t) Engle, K. M.; Mei, T.-S.; Wasa, M.; Yu, J.-Q. Acc. Chem. Res. 2012, 45, 788. (u) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. Angew. Chem., Int. Ed. 2012, 51, 8960. (v) Neufeldt, S. R.; Sanford, M. S. Acc. Chem. Res. 2012, 45, 936. (w) Kuhl, N.; Hopkinson, M. N.; Wencel-Delord, J.; Glorius, F. Angew. Chem., Int. Ed. 2012, 51, 10236. (x) Wencel-Delord, J.; Glorius, F. Nature Chem. 2013, 5, 369.

(9) For selected reviews on direct arylation of (hetero)aromatics see: (a) Joucla, L.; Djakovitch, L. *Adv. Synth. Catal.* 2009, *351*, 673.
(b) Bellina, F.; Rossi, R. *Tetrahedron* 2009, *65*, 10269. (c) Satoh, T.; Miura, M. *Chem. Lett.* 2007, *36*, 200.

(10) For recent reports see: (a) Larbi, K. S.; Fu, H. Y.; Laidaoui, N.; Beydoun, K.; Miloudi, A.; El Abed, D.; Djabbar, S.; Doucet, H. *ChemCatChem* **2012**, *4*, 815. (b) Schnapperelle, I.; Breitenlechner, S.; Bach, T. *Org. Lett.* **2011**, *13*, 3640. (c) Yanagisawa, S.; Ueda, K.; Sekizawa, H.; Itami, K. J. Am. Chem. Soc. **2009**, *131*, 14622.

(11) Ackermann, L.; Barfüßer, S. Synlett 2009, 20, 808.

(12) Akita, Y.; Itagaki, Y.; Takizawa, S.; Ohta, A. Chem. Pharm. Bull. 1989, 37, 1477.

(13) Lane, B. S.; Brown, M. A.; Sames, D. J. Am. Chem. Soc. 2005, 127, 8050.

(14) For illustrative examples of benzo[b]thiophenes in biologically active molecules see: (a) Zhang, T. Y.; O'Toole, J.; Proctor, C. S. *Sulfur reports* **1999**, *22*, 1. (b) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893.

(15) For illustrative examples of benzo[b]thiophenes in organic materials see: (a) Murphy, A. R.; Fréchet, J. M. J. *Chem. Rev.* 2007, 107, 1066. (b) Wang, C.; Dong, H.; Hu, W.; Liu, Y.; Zhu, D. *Chem. Rev.* 2012, *112*, 2208.

(16) Ueda, K.; Yanagisawa, S.; Yamaguchi, J.; Itami, K. Angew. Chem. Int. Ed. 2010, 49, 8946.

(17) Kirchberg, S.; Tani, S.; Ueda, K.; Yamaguchi, J.; Studer, A.; Itami, K. Angew. Chem. Int. Ed. 2011, 50, 2387.

(18) Funaki, K.; Sato, T.; Oi, S. Org. Lett. 2012, 14, 6186.

(19) (a) Ranganath, K. V. S.; Kloesges, J.; Schäfer, A. H.; Glorius, F. *Angew. Chem., Int. Ed.* **2010**, *49*, 7786. (b) Ranganath, K. V. S.; Schäfer, A. H.; Glorius, F. *ChemCatChem* **2011**, *3*, 1889.

(20) Blaser, H.-U.; Indolese, A.; Schnyder, A.; Steiner, H.; Studer, M. J. Mol. Catal. A: Chem. **2001**, *173*, 3.

(21) Significant amounts of 1 and the respective aryl chloride remained after the given reaction time.

(22) For application and limitations see: (a) Crabtree, R. H. Chem. Rev. 2012, 112, 1536. (b) Widegren, J. A.; Finke, R. G. J. Mol. Catal. A: Chem. 2003, 198, 317 and references within.

(23) (a) Lamblin, M.; Nasser-Hardy, L; Hierso, J-C.; Fouquet, E.; Felpin, F-X; *Adv. Syn. Catal.* **2010**, *352*, 33. (b) Huang, L.; Wong, P. W.; Tan, J.; Ang, T. P.; Wang, Z.; J. Phys. Chem. C **2009**, *113*, 10120.

(24) Simmons, E. M.; Hartwig, J. F.; Angew. Chem. Int. Ed. 2012, 51, 3066.

(25) (a) de Meijere, A.; Meyer, Jr., F. E. Angew. Chem. Int. Ed. Engl. **1994**, *33*, 2379. (b) Beletskaya, I. P.; Cheprakov, A. V. Chem. Rev. **2000** *100*, 3009.

(26) See Liégault, B.; Petrov, I.; Gorelsky, S. I.; Fagnou, K. J. Org. Chem., 2010, 75, 1047 and references within.

Journal of the American Chemical Society

