## Asymmetric Catalysis

## Enantioselective Synthesis of Tertiary and Quaternary Stereogenic Centers: Copper/Phosphoramidite-Catalyzed Allylic Alkylation with Organolithium Reagents\*\*

Martín Fañanás-Mastral, Manuel Pérez, Pieter H. Bos, Alena Rudolph, Syuzanna R. Harutyunyan,\* and Ben L. Feringa\*

Copper-catalyzed asymmetric allylic alkylation (AAA) has become a very powerful tool for the enantioselective construction of optically active tertiary carbon stereocenters.<sup>[1]</sup> Recently, this C–C bond-forming reaction has also been successfully applied to the challenging enantioselective synthesis of all-carbon quaternary stereogenic centers in acyclic systems.<sup>[2]</sup> Organozinc,<sup>[3]</sup> aluminium,<sup>[4]</sup> and Grignard<sup>[5]</sup> reagents have been shown to be highly effective in the AAA reaction. However, the cheap and readily available organolithium reagents have not been successfully used in the catalytic asymmetric C–C bond formation of quaternary centers until now.

Organolithium reagents are among the most versatile and widely used reagents in organic synthesis.<sup>[6]</sup> However, the nature of these extremely reactive reagents has hampered its use in enantioselective catalytic C–C bond-formation processes, and successful applications pertain to the use of stoichiometric amounts of chiral ligands<sup>[7]</sup> or to catalytic asymmetric deprotonations<sup>[8]</sup> and additions to imines,<sup>[9]</sup> often with high catalyst loadings and modest enantioselectivities.

Recently, we reported for the first time a general and highly enantioselective catalytic method for the direct addition of organolithium reagents through a copper-catalyzed asymmetric allylic alkylation.<sup>[10]</sup> By using a catalyst comprising CuBr·SMe<sub>2</sub> and TaniaPhos as a chiral ligand, and by selecting a proper combination of dichloromethane and *n*hexane as solvent and co-solvent, we could tame the highly reactive alkyllithium reagents and apply these in a catalytic, highly regio- and enantioselective allylic alkylation of a large variety of allyl bromides (Scheme 1). A limitation found for the CuBr·SMe<sub>2</sub>/TaniaPhos system was that the use of simple allyl chlorides or secondary organolithium reagents in combination with this catalyst led to disappointing selectivities.<sup>[11]</sup> However, we showed in preliminary studies<sup>[10]</sup> that easily

[*] Dr. M. Fañanás-Mastral, Dr. M. Pérez, P. H. Bos, Dr. A. Rudolph,
Dr. S. R. Harutyunyan, Prof. Dr. B. L. Feringa
Stratingh Institute for Chemistry, University of Groningen
Nijenborgh 4, 9747 AG, Groningen (The Netherlands)
E-mail: s.harutyunyan@rug.nl
b.l.feringa@rug.nl
Homepage: http://feringa.fmns.rug.nl/

- [\*\*] We thank the Netherlands Organization for Scientific Research (NWO-CW) and the National Research School Catalysis (NRSC-C) for financial support. M. P. thanks the Xunta de Galicia for an Angeles Alvariño contract and Fondo Social Europeo.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201107840.



R = aryl, halogen-substituted aryl, alkyl, carbamate, ester, protected and free alcohols R' = Me, Et, nBu, nHex

**Scheme 1.** Copper/TaniaPhos-catalyzed AAA of allyl bromides with organolithium reagents

accessible phosphoramidites<sup>[12]</sup> might be suitable ligands for this transformation.

Herein, we report that the use of phosphoramidites as chiral ligands allows the use of both allyl chlorides or bromides in highly enantioselective allylic alkylations in combination with primary as well as secondary organolithium reagents. We also found that a phosphoramidite-based copper catalyst provides all-carbon quaternary stereogenic centers in an enantioselective manner.

We started our study by screening different copper salts, phosphoramidite ligands, and reaction conditions for the addition of *n*-butyllithium to cinnamyl chloride (**1a**) (Table 1). The use of copper(I) thiophene carboxylate  $(CuTC)^{[13]}$  as the copper salt and the phosphoramidite ligand **L1** (Figure 1) in a 1:2 ratio gave rise to modest

**Table 1:** Screening of ligands and reaction conditions for the coppercatalyzed allylic substitution with *n*BuLi.<sup>[a]</sup>

Ph Cl		<i>n</i> BuLi (1 [Cu] (5 r <b>L</b> ( <i>x</i> mol CH <sub>2</sub> Cl <sub>2</sub> ,	I.2 equiv) nol%) (%) –80 °C	nBu Ph	+ Ph	∕∩Bu 3a
Entry	Cu salt	L	[Cu]/L	Addition time [h]	2 a/3 a <sup>[b]</sup>	e.r. <sup>[c]</sup> ( <b>2 a</b> )
1	CuTC	LI	1:2	2	58:42	67:33
2	CuTC	L1	1:1	2	71:29	86:14
3	CuBr·SMe <sub>2</sub>	L1	1:1	2	70:30	91:9
4	CuBr·SMe <sub>2</sub>	L1	1:1	5	91:9	98:2
5	CuBr·SMe <sub>2</sub>	L2	1:1	5	38:62	66:34
6	CuBr·SMe <sub>2</sub>	L3	1:1	5	49:51	86:14
7	CuBr·SMe <sub>2</sub>	L4	1:1	5	82:18	20:80

[a] Reactions performed on a 0.2 mmol scale. *n*BuLi was diluted with *n*-hexane and added dropwise. Full conversion was reached in all cases. [b]  $S_N 2'/S_N 2$  ratio determined by GC and <sup>1</sup>H NMR analysis. [c] Determined by GC analysis using a chiral stationary phase.

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



Figure 1. Phosphoramidite ligands.

selectivity (entry 1). A drastic improvement in the regio- and enantioselectivity was observed when a 1:1 copper/ligand ratio was used (entry 2). By changing the copper salt to CuBr·SMe<sub>2</sub> we observed an enhancement of the enantioselectivity, thus obtaining **2a** with a 91:9 e.r. (entry 3). A longer addition time of 5 hours, using CuBr·SMe<sub>2</sub> as the preferred copper salt, afforded the desired product **2a** with an excellent  $S_N2'/S_N2$  ratio of 91:9 and 98:2 e.r. (entry 4). The use of different phosphoramidite ligands (Figure 1) did not improve these results (entries 5–7).

Once having established the optimized reaction conditions (Table 1, entry 4), we examined the scope of the copper/ phosphoramidite-catalyzed addition of primary organolithium reagents to both allyl chlorides and bromides (Table 2). Excellent yields and enantiomeric ratios were obtained in nearly all cases.

**Table 2:** Copper/phosphoramidite-catalyzed allylic substitution of allyl chlorides and bromides with primary organolithium reagents.<sup>[a]</sup>

Б	$\sim$	R'LI (1.2 ec CuBr•SMe <sub>2</sub> L1 (5.5 mo	quiv) 2 (5 mol% I%)	) →	R'	
R	1: X = 0 4: X = 1	^ СН <sub>2</sub> СІ <sub>2</sub> , - СІ Вг	-80 °C	R´	2	3
Entry	ν X	1/4 (R)	R′	2/3 <sup>[b]</sup>	$Yield \ [\%]^{[c]}$	e.r. <sup>[d]</sup> (2)
1	Cl	<b>1a</b> (Ph)	nBu	91:9	90	98:2 ( <b>2</b> a)
2	Br	<b>4a</b> (Ph)	nВu	87:13	92	96:4 ( <b>2a</b> )
3	Cl	<b>1a</b> (Ph)	nHex	80:20	90	98:2 ( <b>2b</b> )
4	Br	<b>4a</b> (Ph)	nHex	89:11	91	99:1 ( <b>2b</b> )
5	Cl	<b>1a</b> (Ph)	Et <sup>[e]</sup>	85:15	90 <sup>[f]</sup>	96:4 ( <b>2c</b> )
6	Br	<b>4a</b> (Ph)	Et <sup>[e]</sup>	88:12	95 <sup>[f]</sup>	95:5 ( <b>2c</b> )
7	Cl	<b>1b</b> (1-naphthyl)	nBu	50:50	93	96:4 ( <b>2 d</b> )
8	Br	<b>4b</b> (1-naphthyl)	nВu	85:15	91	95:5 ( <b>2 d</b> )
9	Cl	1c (4-ClC <sub>6</sub> H <sub>4</sub> )	nВu	90:10	89	98:2 ( <b>2e</b> )
10	Br	<b>4c</b> , 4-Cl-C <sub>6</sub> H <sub>4</sub>	nBu	85:15	88	97:3 ( <b>2e</b> )
11	Cl	1c (4-ClC <sub>6</sub> H <sub>4</sub> )	Et <sup>[e]</sup>	89:11	72	98:2 ( <b>2 f</b> )
12	Cl	1c (4-ClC <sub>6</sub> H <sub>4</sub> )	nHex	80:20	87	>99:1 ( <b>2 g</b> )
13	Cl	1d (Cy)	nBu	90:10	91	88:12 ( <b>2 h</b> )
14	Br	<b>4d</b> (Me)	nHex	90:10	<b>99</b> <sup>[f]</sup>	91:9 ( <b>2 i</b> )

[a] Reactions performed on a 0.2 mmol scale. The corresponding organolithium reagent was diluted with *n*-hexane and added over 5 h. [b]  $S_N2'/S_N2$  ratios determined by GC and <sup>1</sup>H NMR analysis. [c] Yield of isolated product. [d] Determined by GC analysis using a chiral stationary phase. [e] EtLi was diluted with toluene. [f] Conversion.

The phosphoramidite/copper-catalyzed AAA proved to be also very efficient with allyl bromides, thus providing a complementary catalytic procedure for the earlier TaniaPhosbased method.<sup>[10]</sup> The addition of *n*BuLi to cinnamyl bromide (4a) afforded the product 2a with comparable regio- and enantioselectivity as observed for 1a (Table 2, entry 2). Different primary alkyl lithium reagents gave also very good enantioselectivities both with the chloride 1a and bromide 4a (entries 3-6). When the allyl chloride 1b bearing the more sterically demanding 1-naphthyl substituent was used, the reaction still gave a good level of enantioselectivity but the  $S_N 2'/S_N 2$  products were obtained in a 1:1 ratio (entry 7). However, when the analogous allyl bromide 4b was used the corresponding product 2d was obtained with high enantioselectivity and good regioselectivity (entry 8). Remarkably, the catalytic system also tolerates the presence of aromatic halides. Both p-chloro-substituted cinnamyl chloride (1c) and bromide (4c) could be used without any lithium-halogen exchange and with excellent enantioselectivity (97:3 to > 99:1 e.r.; entries 9–12). Alkyl-substituted allyl halides could also be used for this transformation. Both substrates bearing cyclic (R = Cy; entry 13) and acyclic (R =Me; entry 14) aliphatic substituents afforded the corresponding products 2h and 2i with good selectivity but with slightly lower e.r. values.

The use of the optimal ligand **L1** for the addition of primary organolithium reagents in the reaction of a secondary organolithium reagent, such as *sec*-BuLi, to **1a** afforded the desired compound **5a** as a mixture of diastereoisomers with good regioselectivity ( $S_N2'/S_N2$  91:9) but with disappointing enantioselectivity (Table 3, entry 1). The phosphoramidite **L3** 

Table 3: Optimization for the allylic substitution with sec-BuLi.<sup>[a]</sup>

~ ~		sec-BuLi (1.2 equiv) CuBr·SMe <sub>2</sub> (5 mol%) L (5.5 mol%)	ww.	~	
Ph	X	CH <sub>2</sub> Cl <sub>2</sub> , -80 °C	Ph M	* Ph	
<b>1a</b> : X = Cl <b>4a</b> : X = Br			5a/5a'		6a
Entry	Х	L	5/6 <sup>[b,c]</sup>	e	.r. <sup>[b]</sup>
				5 a	5 a'
1 <sup>[d]</sup>	Cl	LI	91:9	60:40	66:34
2 <sup>[e]</sup>	Cl	L2	90:10	28:72	26:74
3 <sup>[f]</sup>	Cl	L3	97:3	91:9	91:9
4 <sup>[g]</sup>	Br	L3	79:21	69:31	71:29

[a] Reaction conditions: see Table 2. Full conversion was reached in all cases.
[b] Determined by GC analysis using a chiral stationary phase.
[c] Compound **6a** was obtained as a racemic mixture in all cases.
[d] d.r. = 1.5:1. [e] d.r. = 1.2:1. [f] d.r. = 1:1. [g] d.r. = 1.3:1.

was found to be the most effective ligand for the addition of *sec*-BuLi, thus affording the compound **5a** with a high  $S_N 2'/S_N 2$  ratio (97:3) and 91:9 e.r. for both diastereoisomers (entry 3).<sup>[14]</sup> In this case, the use of the more reactive cinnamyl bromide (**4a**) led to a decrease in both regio- and enantio-selectivity of the reaction (entry 4).

By using the optimized ligand L3, the addition of *sec*-BuLi to different allyl chlorides 1 provided the corresponding products 5 in high yields and with high regioselectivities and

Table 4: Copper/phosphoramidite-catalyzed allylic substitution	on with
secondary organolithium reagents. <sup>[a]</sup>	

R	Cl + Li	CuBr•SMe <sub>2</sub> (5 mol%) L <b>3</b> (5.5 mol CH <sub>2</sub> Cl <sub>2</sub> , –8	%) 0°℃ R		R
1	<i>sec</i> -BuLi (R' = Me) <i>i</i> PrLi (R' = H)		5/5': I 7: R =	R = Me - H	6: R = Me 8: R = H
Entry	1 (R)	R′	5/6 or 7/8 <sup>[b]</sup>	Yield [%] <sup>[c]</sup>	e.r. <sup>[e]</sup> ( <b>5</b> <sup>[d]</sup> or <b>7</b> )
1 <sup>[d]</sup>	<b>1a</b> (Ph)	Me	97:3	80	91:9 ( <b>5 a</b> ) 91:9 ( <b>5 a'</b> )
2 <sup>[d]</sup>	<b>1b</b> (1-naphthyl)	Me	92:8	85	88:12 (5b) 88:12 (5b')
3 <sup>[d]</sup>	1c (4-ClC <sub>6</sub> H <sub>4</sub> )	Me	93:7	96	90:10 ( <b>5 c</b> ) 88:12 ( <b>5 c'</b> )
4	<b>1a</b> (Ph)	Н	90:10	77	96:4 ( <b>7 a</b> )
5	<b>1b</b> (1-naphthyl)	Н	96:4	98	94:6 ( <b>7 b</b> )
6	1c (4-ClC <sub>6</sub> H <sub>4</sub> )	Н	92:8	95	96:4 ( <b>7 c</b> )
7	1 d (Cy)	Н	89:11	83	91:9 ( <b>7 d</b> )

 <sup>[</sup>a]-[c] See Table 2. Full conversion was reached in all cases. [d] d.r. = 1:1.
 [e] Determined by GC or HPLC analysis using a chiral stationary phase.

enantioselectivities (Table 4, entries 1-3). Although the e.r. values are only up to 91:9, it is important to note that these examples represent, to the best of our knowledge, the first catalytic asymmetric transfer of the sec-butyl moiety. Furthermore, the isopropyl moiety has been a challenging secondary alkyl group to transfer by the copper-catalyzed AAA, and generally only modest enantioselectivities are obtained with cinnamyl substrates and other organometallic reagents.<sup>[3b,15]</sup> Remarkably, by using *i*PrLi in combination with CuBr·SMe2 and L3, we could achieve 96:4 e.r. in the allylic alkylation of 1a (entry 4). Similar to the addition of primary organolithium reagents, the addition of iPrLi to the more sterically demanding 1b and the p-chloro-substituted 1c gave rise to high enantioselectivities (entries 5 and 6). The cyclohexyl-substituted allyl chloride 1d could also be used, but with slightly lower regio- and enantioselectivity were obtained (entry 7).

Notably, by using the same catalytic system (CuBr·SMe<sub>2</sub>/ L3) phenyllithium could be used in the allylic substitution of *p*-chloro cinnamylbromide **4c**, thus providing the corresponding diaryl-substituted product with >99:1 e.r. but with low regioselectivity ( $S_N 2'/S_N 2$  40:60).<sup>[10]</sup>

This protocol could also be extended to the first enantioselective synthesis of all-carbon quaternary stereogenic centers employing organolithium reagents. After extensive screening of different phosphoramidite ligands (see the Supporting Information for details), ligand **L4** (Figure 1) turned out to be the best ligand for the addition of *n*BuLi to *E*-trisubstituted allyl bromide **9a** (Table 5). When *n*BuLi was added over a 10 hour period to a solution of **9a** and the catalyst at -80 °C, the reaction gave rise to the product **10a** with 92:8 e.r. and with nearly perfect S<sub>N</sub>2' selectivity (entry 1). The use of the trisubstituted allyl chloride analogue of **9a** did not result in any conversion. By using these optimized reaction conditions we examined the scope of this new method for the enantioselective synthesis of all-carbon quaternary centers (Table 5).

Table 5:	Enantiose	lective c	opper/p	hospl	horamid	ite-cata	lyzed	syntl	nesis
of quate	ernary carbo	on atom	s with o	rgano	olithium	reagent	s. <sup>[a]</sup>		

	R'L Cu L4	i (1.2 equi Br•SMe <sub>2</sub> ( (5.5 mol%	v) 5 mol%) )	R' +	
F	Br C	CH <sub>2</sub> CI <sub>2</sub> , -80	o°C R′	R	~R'
	9			10	11
Entry	<b>9</b> (R)	R′	10/11 <sup>[b]</sup>	Yield [%] <sup>[c]</sup>	e.r. <sup>[d]</sup> ( <b>10</b> )
1	<b>9a</b> (Ph)	nBu	98:2	92	92:8 ( <b>10</b> a)
2	<b>9a</b> (Ph)	nHex	92:8	92	86:14 ( <b>10b</b> )
3	9b (4-ClC <sub>6</sub> H <sub>4</sub> )	<i>n</i> Bu	97:3	72	90:10 ( <b>10 c</b> )
4	9b (4-ClC <sub>6</sub> H <sub>4</sub> )	nHex	94:6	90	91:9 ( <b>10d</b> )
5	<b>9c</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	nВu	94:6	91	88:12 ( <b>10e</b> )
5	<b>9c</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	nHex	91:9	74	89:11 ( <b>10 f</b> )
7	<b>9d</b> (Cy)	nВu	98:2	77	88:12 ( <b>10g</b> )
8	<b>9e</b> (2-MeC <sub>6</sub> H <sub>4</sub> )	<i>n</i> Bu	92:8	92	53:47 (10h)
Э	<b>9e</b> (2-MeC <sub>6</sub> H <sub>4</sub> )	nHex	91:9	90	53:47 ( <b>10i</b> )
10	<b>9 f</b> (2-MeOC <sub>6</sub> H <sub>4</sub> )	nВu	98:2	89	95:5 ( <b>10j</b> )
11	<b>9 f</b> (2-MeOC <sub>6</sub> H <sub>4</sub> )	nHex	95:5	90	95:5 ( <b>10k</b> )
12	<b>9g</b> (2-BrC <sub>6</sub> H <sub>4</sub> )	nВu	>98:2	93	92:8 ( <b>10</b> )
13	<b>9g</b> (2-BrC <sub>6</sub> H <sub>4</sub> )	nHex	>98:2	93	91:9 ( <b>10</b> m)

[a]–[c] See Table 2. Addition time: 10 h. Full conversion was reached in all cases. [d] Determined by GC or HPLC analysis using a chiral stationary phase.

This new protocol was found to be very efficient with primary alkyl organolithium reagents, such as nBuLi and n-HexLi, which gave excellent regioselectivity and good enantioselectivities ranging from 86:14 to 92:8 e.r. when the phenyl derivative 9a (entries 1 and 2) and p-chloro- and pmethyl-substituted allyl bromides 9b,c were used (entries 3-6). An alkyl-substituted allyl bromide, such as the cyclohexyl derivative 9d, could also be used in this transformation to afford the desired product 10g with excellent regioselectivity and good enantioselectivity (entry 7). An intriguing and synthetically highly significant observation was made when ortho-substituted cinnamyl bromides were used. Reactions with the *o*-methyl-substituted bromide **9e** still showed very good regioselectivity but the corresponding products 10h,i were obtained as a nearly racemic mixture (entries 8 and 9). In sharp contrast, when the *o*-methoxy-substituted bromide 9 f was used the desired products 10 j,k were obtained with excellent stereoselectivity (95:5 e.r.) and with almost total  $S_N 2'$  selectivity (entries 10 and 11). A similar trend in the regio- and enantioselectivity was observed when another substrate bearing a coordinating functionality in the ortho position, such as o-bromo-substituted cinnamyl bromide (9g), was used (entries 12 and 13). These results suggest that a possible coordination between the methoxy or bromide substituent atom and the copper complex can be a key factor to afford high enantioselectivity when ortho-substituted cinnamyl bromides are used. It is also important to note, as described above for other aromatic halides, that there was no evidence of the common lithium-halogen exchange even in the case of o-bromo-substituted cinnamyl bromide 9g.

In summary, we have developed efficient catalytic and highly enantioselective methodology for the asymmetric alkylation of both allyl chlorides and bromides with organolithium reagents using monodentate phosphoramidites as chiral ligands. These protocols include efficient asymmetric addition of both primary and secondary organolithium reagents. Furthermore, for the first time, the enantioselective synthesis of quaternary carbon stereogenic centers using these extremely reactive organometallic reagents is presented.

## **Experimental Section**

Typical procedure: A Schlenk tube equipped with septum and stirring bar was charged with CuBr·SMe2 (0.01 mmol, 2.06 mg, 5 mol%) and the appropriate ligand (0.012 mmol, 6 mol%). Dry dichloromethane (2 mL) was added and the solution was stirred under nitrogen at room temperature for 15 min. Then, the allyl halide (0.2 mmol) was added and the resulting solution was cooled to -80 °C. In a separate Schlenk tube, the corresponding organolithium reagent (0.24 mmol, 1.2 equiv) was diluted with n-hexane (combined volume of 1 mL) under nitrogen and added dropwise to the reaction mixture over 5 h (over 10 h for quaternary centers) using a syringe pump. Once the addition was complete, the mixture was stirred for another 2 h at -80 °C. The reaction was quenched with a saturated aqueous NH<sub>4</sub>Cl solution (2 mL) and the mixture was warmed up to room temperature, diluted with dichloromethane and the layers were separated. The aqueous layer was extracted with dichloromethane  $(3 \times 5 \text{ mL})$  and the combined organic layers were dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and the solvent was evaporated in vacuo. The crude reaction mixture was purified by flash chromatography on silica gel using different mixtures of *n*-pentane/Et<sub>2</sub>O as the eluent.

Received: November 7, 2011 Revised: December 9, 2011 Published online: January 19, 2012

**Keywords:** allylic compounds · copper · ligand effects · organolithium reagents · synthetic methods

- [1] For reviews, see: a) A. S. E. Karlström, J. E. Bäckvall in Modern Organocopper Chemistry (Ed.: N. Krause), Wiley-VCH, Weinheim, 2002; b) H. Yorimitsu, K. Oshima, Angew. Chem. 2005, 117, 4509-4513; Angew. Chem. Int. Ed. 2005, 44, 4435-4439; c) S. Harutyunyan, T. den Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, Chem. Rev. 2008, 108, 2824-2852; d) A. Alexakis, J. E. Bäckvall, N. Krause, O. Pàmies, M. Diéguez, Chem. Rev. 2008, 108, 2796-2823; For recent examples, see: e) K. B. Selim, Y. Matsumoto, K. Yamada, K. Tomioka, Angew. Chem. 2009, 121, 8889-8891; Angew. Chem. Int. Ed. 2009, 48, 8733-8735; f) K. Akiyama, F. Gao, A. H. Hoveyda, Angew. Chem. 2010, 122, 429-433; Angew. Chem. Int. Ed. 2010, 49, 419-423; g) J. F. Teichert, S. Zhang, A. W. van Zijl, J. W. Slaa, A. J. Minnaard, B. L. Feringa, Org. Lett. 2010, 12, 4658-4660; h) J.-B. Langlois, A. Alexakis, Angew. Chem. 2011, 123, 1917-1921; Angew. Chem. Int. Ed. 2011, 50, 1877-1881; i) M. Fañanás-Mastral, B. ter Horst, A. J. Minnaard, B. L. Feringa, Chem. Commun. 2011, 47, 5843-5845; j) P. Tosatti, A. J. Campbell, D. House, A. Nelson, S. P. Marsden, J. Org. Chem. 2011, 76, 5495-5501.
- For a review, see: a) J. Prakash, I. Marek, *Chem. Commun.* 2011, 47, 4593-4623; For a review on the formation of quaternary centers through metal-catalyzed asymmetric conjugate addition, see: b) C. Hawner, A. Alexakis, *Chem. Commun.* 2010, 46, 7295-7306.

- [3] a) N. Harrington-Frost, H. Leuser, M. I. Calaza, F. F. Kneisel, P. Knochel, Org. Lett. 2003, 5, 2111–2114; b) H. Leuser, S. Perrone, F. Liron, F. F. Kneisel, P. Knochel, Angew. Chem. 2005, 117, 4703–4707; Angew. Chem. Int. Ed. 2005, 44, 4627–4631; c) M. A. Kacprzynski, A. H. Hoveyda, J. Am. Chem. Soc. 2004, 126, 10676–10681; d) J. J. Van Veldhuizen, J. E. Campbell, R. E. Giudici, A. H. Hoveyda, J. Am. Chem. Soc. 2005, 127, 6877–6882; e) M. A. Kacprzynski, T. L. May, S. A. Kazane, A. H. Hoveyda, Angew. Chem. 2007, 119, 4638–4642; Angew. Chem. Int. Ed. 2007, 46, 4554–4558.
- [4] a) F. Gao, K. P. McGrath, Y. Lee, A. H. Hoveyda, J. Am. Chem. Soc. 2010, 132, 14315-14320; b) F. Gao, Y. Lee, K. Mandai, A. H. Hoveyda, Angew. Chem. 2010, 122, 8548-8552; Angew. Chem. Int. Ed. 2010, 49, 8370-8374.
- [5] a) B. Breit, P. Demel, C. Studte, Angew. Chem. 2004, 116, 3874; Angew. Chem. Int. Ed. 2004, 43, 3785-3789; b) Y. Lee, A. H. Hoveyda, J. Am. Chem. Soc. 2006, 128, 15604-15605; c) O. Jackowski, A. Alexakis, Angew. Chem. 2010, 122, 3418-3422; Angew. Chem. Int. Ed. 2010, 49, 3346-3350.
- [6] The Chemistry of Organolithium Compounds (Eds.: Z. Rappoport, I. Marek), Wiley-VCH, Weinheim, 2004.
- [7] a) E. J. Corey, R. Naef, F. J. Hannon, J. Am. Chem. Soc. 1986, 108, 7114–7116; b) S. E. Denmark, C. M. Stiff, J. Org. Chem. 2000, 65, 5875–5878; c) M. R. Luderer, W. F. Bailey, M. R. Luderer, J. D. Fair, R. J. Dancer, M. B. Sommer, Tetrahedron: Asymmetry 2009, 20, 981–998.
- [8] a) J. L. Bilke, S. P. Moore, P. O'Brien, J. Gilday, Org. Lett. 2009, 11, 1935–1938; b) T. K. Beng, R. E. Gawley, J. Am. Chem. Soc. 2010, 132, 12216–12217.
- [9] a) K. Tomioka, I. Inoue, I. Mitsuru, S. Kenji, K. Koga, *Tetrahedron Lett.* **1991**, *24*, 3095–3098; b) S. E. Denmark, N. Nakajima, O. J.-C. Nicaise, *J. Am. Chem. Soc.* **1994**, *116*, 8797– 8798; c) A. Alexakis, F. Amiot, *Tetrahedron: Asymmetry* **2002**, *13*, 2117–2122.
- [10] M. Pérez, M. Fañanás-Mastral, P. H. Bos, A. Rudolph, S. R. Harutyunyan, B. L. Feringa, *Nat. Chem.* 2011, *3*, 377–381.
- [11] The CuBr·SMe<sub>2</sub>/TaniaPhos-catalyzed addition of *n*BuLi to cinnamyl chloride gave rise to an  $S_N2'/S_N2$  ratio of 10:90 with 55:45 e.r. for the  $S_N2'$  product. The addition of *sec*-BuLi to cinnamyl bromide in combination with this catalyst gave a 13:42:43 mixture of  $S_N2'/S_N2$ /homocoupling products, with the  $S_N2'$  product being racemic.
- [12] J. F. Teichert, B. L. Feringa, Angew. Chem. 2010, 122, 2538– 2582; Angew. Chem. Int. Ed. 2010, 49, 2486–2528.
- [13] CuTC has been shown to be an effective copper salt in combination with phosphoramidite ligands in the copper-catalyzed AAA with Grignard reagents. For some examples, see: a) K. Tissot-Croset, D. Polet, A. Alexakis, Angew. Chem. 2004, 116, 2480-2482; Angew. Chem. Int. Ed. 2004, 43, 2426-2428; b) M. Fañanás-Mastral, B. L. Feringa, J. Am. Chem. Soc. 2010, 132, 13152-13153; c) J.-B. Langlois, A. Alexakis, Adv. Synth. Catal. 2010, 352, 447-457.
- [14] It is important to note that the configuration of the newly formed stereocenter is controlled by the catalyst while the presence of two diastereoisomers is the result of using a racemic alkyllithium reagent.
- [15] a) A. Alexakis, K. Croset, Org. Lett. 2002, 4, 4147–4149;
  b) A. W. van Zijl, L. A. Arnold, A. J. Minnaard, B. L. Feringa, Adv. Synth. Catal. 2004, 346, 413–420; c) W. Lölsberg, S. Ye, H.-G. Schmalz, Adv. Synth. Catal. 2010, 352, 2023–2031.