Bioorganic & Medicinal Chemistry Letters 22 (2012) 3152-3156

Contents lists available at SciVerse ScienceDirect



Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

The facile synthesis of multifunctional PAMAM dendrimer conjugates through copper-free click chemistry

Baohua Huang^{*}, Jolanta F. Kukowska-Latallo, Shengzhuang Tang, Hong Zong, Kali B. Johnson, Ankur Desai, Chris L. Gordon, Pascale R. Leroueil, James R. Baker Jr.^{*}

Michigan Nanotechnology Institute for Medicine and Biological Sciences, University of Michigan, 9220 MSRB III, Box 0648, Ann Arbor, MI 48109, USA

ARTICLE INFO

Article history: Received 31 January 2012 Revised 8 March 2012 Accepted 13 March 2012 Available online 21 March 2012

Keywords: PAMAM dendrimer Copper-free click conjugation Drug delivery

ABSTRACT

The facile conjugation of three azido modified functionalities, namely a therapeutic drug (methotrexate), a targeting moiety (folic acid), and an imaging agent (fluorescein) with a G5 PAMAM dendrimer scaffold with cyclooctyne molecules at the surface through copper-free click chemistry is reported. Mono-, di-, and tri-functional PAMAM dendrimer conjugates can be obtained via combinatorial mixing of different azido modified functionalities simultaneously or sequentially with the dendrimer platform. Preliminary flow cytometry results indicate that the folic acid targeted nanoparticles are efficiently binding with KB cells.

© 2012 Elsevier Ltd. All rights reserved.



Figure 1. Structures of PAMAM dendrimer platform, γ -azido modified MTX, and azido modified fluorescein (numbers of functionalities are rounded to the nearest integers).

amine groups on the dendrimer surface were neutralized by acetylation (Fig. 1, compound 1). Variable equivalents (5, 10, and 18 equiv) of γ -azido modified methotrexate (γ -N₃-MTX) were reacted with the dendrimer platform efficiently with >90% of the modified drug conjugated. Here we report the facile conjugation of multiple functional groups with the PAMAM dendrimer platform via copper-free click chemistry. Three types of azido modified functionalities, namely a therapeutic drug (γ -N₃-MTX, **2**), a targeting group (γ -N₃-folic acid, **7**), and an imaging agent (N₃-FITC, **3**) were reacted with the G5 dendrimer platform simultaneously or sequentially to achieve multifunctional nanodevices.

have been used as a platform to deliver therapeutics and imaging agents, both in vitro¹ and in vivo.² A sequential synthetic strategy based on amide and ester chemistry is often used to conjugate small molecules, such as therapeutic drugs, targeting groups and imaging agents, to generate the functional forms of the dendrimer.³ While sufficient for small-scale needs, the extensive purification & characterization required between each small molecule addition, as well as batch-to-batch reproducibility issues, made it ill-suited for large-scale needs. Cu(I)-catalyzed alkyne azide 1,3-dipolar cycloaddition (CuAAC), widely used for bioconjugation due to its high yields, tolerance toward functional groups, and lack of concurrent side-reactions, was initially an attractive alternative to our amide and ester conjugation chemistry.⁴ However, the copper catalyst proved difficult to remove from the dendrimers⁵ and the cytotoxicity⁶ associated with the residual copper prevented CuAAC from being a viable option. Several metal-free bioorthogonal labeling reactions such as strain-promoted cycloaddition of azides,⁷ which relies on the unusually high ring strain associated with cyclooctyne to react rapidly and selectively with azide groups,⁸ have since been developed.⁹

For several decades, polyamidoamine (PAMAM) dendrimers

Recently, we synthesized a G5 PAMAM platform with cyclooctyne ligand on the surface for copper-free click conjugation with azido modified functionalities.¹⁰ Briefly, the G5 PAMAM dendrimer was first partially (70%) acetylated and then coupled with 20 cyclooctyne ligands through amide bonds. The remaining primary

* Corresponding authors.

E-mail addresses: baohua@med.umich.edu, baohua@umich.edu (B. Huang), jbakerjr@med.umich.edu (J.R. Baker).

⁰⁹⁶⁰⁻⁸⁹⁴X/ $\$ - see front matter @ 2012 Elsevier Ltd. All rights reserved. http://dx.doi.org/10.1016/j.bmcl.2012.03.052

 γ -N₃-MTX (Fig. 1, compound **2**) was synthesized using our published method. Briefly, 4-amino-4-deoxy- N^{10} -methylpteroic acid was first reacted with Glu- α -OtBu using BOP reagent to give α -OtBu-MTX. The γ -carboxylic acid group was then coupled with 3-azido-propylamine in the presence of HATU. After exposure of the α carboxylic acid functional group using trifluoroacetic acid, γ -N₃-MTX was obtained in excellent yield.¹⁰ Azido modified fluorescein (Fig. 1, compound **3**) was synthesized in a straightforward manner using fluorescein isothiocyanate and 3-azido-propylamine (see Supplementary data).

A number of cancer cell lines exhibit high levels of the cell surface folate receptor (FR) while its expression is highly restricted in most normal tissues. The degree of overexpression of the FR on human epithelial cancer cells ranges from 30-fold (ovarian carcinoma) to 1400-fold (endometrial cancer).^{4b} Hence, the FR is a very attractive molecular target for receptor-based tumor imaging and therapy.¹¹ The folic acid structure offers two positions for bioconiugation, the α - and γ -carboxylic acid. Whether or not the binding affinity and in vivo pharmacological behavior is dependent on the site of conjugation is still debated in the literature.¹² Here we synthesized a γ -azido modified folic acid (Scheme 1). Mindt et al. synthesized a γ -azido modified folic acid starting from double-protected pteroic acid.¹³ Starting from commercially available N^{10} -(trifluoroacetyl) mono-protected pteroic acid, we synthesized γ -azido folic acid (5) in five steps with good yield. Originally, we tried to use BocGluOMe as a precursor,¹³ but the final removal of α -methyl ester with NaOH yielded 8-15% racemization (data not shown). In order to avoid the basic conditions that were responsible for the racemization, we replaced BocGluOMe with FmocGluOtBu. Briefly, FmocGluOtBu was coupled with 3-azido-propylamine to obtain compound 4, followed by the exposure of amino group with piperidine to give compound **5**. N^{10} -(trifluoroacetyl) pteroic acid was then coupled with **5** to give a folic acid intermediate with α -*t*Bu ester and N^{10} -(trifluoroacetyl) protecting groups (**6**). The removal of these two groups sequentially with ammonium hydroxide and TFA gave compound 7. Compound 7 was obtained by precipitation with diethyl ether and collected by centrifugation (see Supplementary data). We later found that the protection of the primary amino group at the pteridine ring was not necessary.

The copper-free click conjugation of the dendrimer platform with azido modified functionalities was then carried out (Scheme 2). Briefly, the methanol solution of G5 PAMAM dendrimer with cyclooctyne ligand molecules (1) was mixed simultaneously or sequentially with one, two or three of the previously mentioned targeting moiety (γ -N₃-FA, 7), therapeutic drug (γ -N₃-MTX, 2), and imaging agent (N₃-FITC, 3). Because of the structural similarity of the azido modified folic acid and methotrexate, we assumed that

they would have similar reactivity. Therefore, the di-functional conjugate was made by mixing these two compounds simultaneously with the platform. The conjugates containing the structurally dissimilar fluorescein were synthesized in a slightly different manner to ensure that the average number of fluoresceins per conjugate was identical, a critical requirement for the in vitro cell binding studies. In that case, we mixed the N₃-FITC with the dendrimer platform and allowed the click reaction to go to completion. The reaction mixture was then separated into four equal aliquots; one aliquot was purified to give compound **11**. Subsequent addition of γ -N₃-FA, γ -N₃-MTX, and both of these two compounds to three of the aliquots gave conjugates 12, 13, and 14, each containing the same average number of fluorescein molecules (Scheme 2). All of the click reactions were stirred for 24 h at room temperature shielded from light. After removal of the organic solvent, the residues were dissolved in phosphate buffer saline $(1 \times PBS)$ buffer. pH = 7.4) and purified by 10 K centrifugal filters (PBS \times 3. DI water \times 6) and then lyophilized to give compounds 8-14 with good yields. More than 90% of the azido modified functionalities were conjugated through click reaction with the dendrimer platform.

Using the integration of the acetamide proton of the dendrimer platform as an internal reference, the average number of each type of small molecule conjugated was determined using ¹H NMR. The aromatic proton in the pteridine ring of MTX and folic acid had a chemical shift at 8.61 and 8.64 ppm.^{10,13} After the conjugation, the pteridine proton signal of folic acid shifted down field to 8.71–8.73 ppm.¹³ By comparing the integration of these two peaks to the integration of the acetamide proton peak we determined the numbers of MTX and folic acid molecules conjugated to the dendrimer platform. The ¹H NMR spectra of compounds **8**, **9**, and **10** are shown in Figure 2. In these cases, the integrations of the internal reference peaks were set to 276 which represented 96 acetamide groups.¹⁰ The results indicated that the average numbers of MTX and folic acid in compound 10 were 5.95 and 3.06. Compound 8 had 5.93 MTX molecules and compound 9 had 2.79 folic acid molecules per dendrimer molecule (see inserts in Fig. 2), respectively.

The ¹H NMR spectra of compounds **11–14** are shown in Figure 3. The number of fluorescein molecules conjugated was first calculated based on the relative proton integrations of the eight aromatic protons in the fluorescein structure, located between 6.40 and 7.81 ppm, and the internal reference. The result indicated there were 3 dye molecules per dendrimer molecule averagely (Fig. 3c). Because compounds **12**, **13**, and **14** were synthesized by subsequent additions of azido modified MTX and/or folic acid, the average number of fluorescein molecules per conjugate was identical. Using the same method described above, we determined



Scheme 1. The synthesis of a γ -azido modified folic acid.



Scheme 2. The copper-free click conjugations of the dendrimer platform with azido modified functionalities.



Figure 2. ¹H NMR of compounds 10(a), 8(b), and 9(c). Inserts are the enlargements of the spectra (8.40–8.90 ppm) showing the pteridine ring proton signals of folic acid (FA) and methotrexate (MTX).



Figure 3. ¹H NMR of compounds 11(a), 13(b), 12(c), and 14(d).

the average numbers of MTX and folic acid molecules conjugated to the platform. Results indicated that there were 3.53 molecules of folic acid and 6.10 molecules of MTX in compound **13** (Fig. 3b), 6.06 molecules of MTX in compound **12** (Fig. 3c), and 3.37 molecules of folic acid in compound **14** (Fig. 3d), respectively.

The KB human tumor cell line, which overexpresses the folate receptor¹⁴ was purchased from the American Type Tissue Collection (Manassas, VA) and maintained in vitro at 37 °C, 5% CO₂ in folate-deficient RPMI 1640 supplemented with penicillin (100 units/mL), streptomycin (100 μ G/mL), and 10% heat-inactivated fetal bovine serum. The cells were harvested with trypsin–EDTA solution, washed and resuspended in PBS. The cell suspension (10⁶ cells in 0.1 mL) was incubated for 30 min on ice with 0.3 μ M of targeted (G5-FITC-FA) or non-targeted (G5-FITC) conjugate. After incubation the cells were washed once with phosphate-buffered saline (PBS) and re-suspended in PBS for analysis. Flow cytometry analysis was done using Accuri C6 Flow Cytometer (BDBiosciences) and CFlow Plus software. Data were reported as the mean channel fluorescence of the cell population stained with 0.3 μ M of G5-FITC-FA or G5-FITC. Flow cytometry results indicated that the targeted



Figure 4. Flow cytometry analysis of targeted (G5-FITC-FA), non-targeted (G5-FITC), and unstained control (PBS).

nanoparticle binds specifically to KB cells. The non-targeted nanoparticle did not bind to KB cells (Fig. 4). In conclusion, we have successfully synthesized mono-, di-, and tri-functionalized G5 PAMAM dendrimer conjugates with copperfree click conjugation method. An azido modified targeting moiety, a therapeutic drug, and an imaging reagent were mixed with a G5 PAMAM dendrimer nanoplatform simultaneously or sequentially to give mono-, di-, and tri-functional conjugates. The reactions were very efficient and reproducible. Preliminary flow cytometry results indicate that the FA targeted nanoparticles are efficiently binding with KB cells. The in vitro toxicity studies of these materials toward different cancer cell lines are under investigation.

Acknowledgements

This project has been funded in whole or in part with Federal funds from the National Cancer Institute, National Institutes of Health, under award 1 R01 CA119409.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/j.bmcl.2012.03.052.

References and notes

 Thomas, T. P.; Majoros, I. J.; Kotlyar, A.; Kukowska-Latallo, J. F.; Bielinska, A.; Myc, A.; Baker, J. R. J. Med. Chem. 2005, 48, 3729.

- Myc, A.; Kukowska-Latallo, J.; Cao, P.; Swanson, B.; Battista, J.; Dunham, T.; Baker, J. R. Anticancer Drugs 2010, 21, 186.
- (a) Thomas, T. P.; Shukla, R.; Kotlyar, A.; Liang, B.; Ye, J. Y.; Norris, T. B.; Baker, J. R. *Biomacromolecules* **2008**, *9*, 603; (b) Shukla, R.; Thomas, T. P.; Peters, J.; Kotlyar, A.; Myc, A.; Baker, J. R. *Chem. Commun.* **2005**, 5739.
- (a) Meldal, M.; Tornoe, C. W. *Chem. Rev.* **2008**, *108*, 2952; (b) Mullen, D. G.; McNerny, D. Q.; Desai, A.; Cheng, X.-m.; DiMaggio, S. C.; Kotlyar, A.; Zhong, Y.; Qin, S.; Kelly, C. V.; Thomas, T. P.; Majoros, I.; Orr, B. G.; Baker, J. R., Jr.; Holl, M. M. B. *Bioconjug. Chem.* **2011**, *22*, 679.
- van Dijk, M.; Rijkers, D. T. S.; Liskamp, R. M. J.; van Nostrum, C. F.; Hennink, W. E. Bioconjug. Chem. 2001, 2009, 20.
- (a) Link, A. J.; Vink, M. K. S.; Tirrell, D. A. J. Am. Chem. Soc. 2004, 126, 10598; (b) Gierlich, J.; Burley, G. A.; Gramlich, P. M. E.; Hammond, D. M.; Carell, T. Org. Lett. 2006, 8, 3639.
- Sletten, E. M.; Nakamura, H.; Jewett, J. C.; Bertozzi, C. R. J. Am. Chem. Soc. 2010, 132, 11799.
- 8. Ornelas, C.; Broichhagen, J.; Weck, M. J. Am. Chem. Soc. 2010, 132, 3923.
- Becer, C. R.; Hoogenboom, R.; Schubert, U. S. Angew. Chem., Int. Ed. 2009, 48, 4900
- Huang, B. H.; Desai, A.; Zong, H.; Tang, S. Z.; Leroueil, P.; Baker, J. R. Tetrahedron Lett. 2011, 52, 1411.
- (a) Kukowska-Latallo, J. F.; Candido, K. A.; Cao, Z. Y.; Nigavekar, S. S.; Majoros, I. J.; Thomas, T. P.; Balogh, L. P.; Khan, M. K.; Baker, J. R. *Cancer Res.* **2005**, *65*, 5317;
 (b) Majoros, I. J.; Myc, A.; Thomas, T.; Mehta, C. B.; Baker, J. R. *Biomacromolecules* **2006**, *7*, 572.
- (a) Ross, T. L.; Honer, M.; Lam, P. Y. H.; Mindt, T. L.; Groehn, V.; Schibli, R.; Schubiger, P. A.; Ametamey, S. M. *Bioconjug. Chem.* **2008**, *19*, 2462; (b) Leamon, C. P.; DePrince, R. B.; Hendren, R. W. J. Drug Target. **1999**, *7*, 157; (c) Ke, C. Y.; Mathias, C. J.; Green, M. A. Nucl. Med. Biol. **2003**, *30*, 811.
- Mindt, T. L.; Muller, C.; Melis, M.; de Jong, M.; Schibli, R. *Bioconjug. Chem.* 2008, 19, 1689.
- 14. Turek, J. J.; Leamon, C. P.; Low, P. S. J. Cell Sci. 1993, 106, 423.