C–H Activation

Highly Selective Trifluoromethylation of 1,3-Disubstituted Arenes through Iridium-Catalyzed Arene Borylation**

Tianfei Liu, Xinxin Shao, Yaming Wu, and Qilong Shen*

Dedicated to Professor Weiyuan Huang on the occasion of his 90th birthday

The trifluoromethyl group has evolved into an indispensable structural motif^[1] for the pharmaceutical industry because of its unique size, electronic properties, and excellent metabolic stability, as evidenced by a large number of trifluoromethylated pharmaceuticals and drug candidates, such as the antidepressant Prozac,^[2] the fungicide Trifloxystrobin,^[3] and the herbicide Fusilade.^[4] Consequently, development of new and efficient methods to introduce the trifluoromethyl group to small molecules are of high interest.^[5]

Toward this end, several copper- and palladium-catalyzed trifluoromethylations of aryl halides or aryl boronic acids have been reported recently.^[6-10] For example, Amii and coworkers reported the first copper-catalyzed trifluoromethylation of electron-poor aryl iodides with the Ruppert–Prakash reagent, CF₃SiEt₃, in the presence of CuI/1,10-phenanthroline (Scheme 1).^[7] Buchwald and co-workers reported a palladium-catalyzed trifluoromethylation of aryl chlorides by use of sterically hindered electron-rich ligands.^[8] Qing and Chu,



Scheme 1. Previous metal-catalyzed approaches to trifluoromethyl arenes.

- [*] T. Liu, X. Shao, Dr. Y. Wu, Prof. Dr. Q. Shen Key Laboratory of Organofluorine Chemistry Shanghai Institute of Organic Chemistry Chinese Academy of Sciences 345 Lingling Road, Shanghai 200032 (China) E-mail: shenql@sioc.ac.cn
- [**] The authors thank Ms. Xiangqing Jia of the China Pharmaceutical University for conducting some experiments. We also thank the Key Program of the National Natural Science Foundation of China (21032006, 20632070), the National Natural Science Foundation of China (21172245), and the National Basic Research Program of China (2010CB126103, 2012CB821602) for financial support.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201106673.

and Buchwald and co-workers independently reported the copper-catalyzed trifluoromethylation of aryl boronic acids under oxidation conditions.^[9,10] While these methods overcome the shortcomings of the classic Swarts reaction^[11] or the [CuCF₃] strategy,^[7,12] which requires harsh reaction conditions, stoichiometric amounts of organometallic reagents, and occurs with limited substrate scope, the trifluoromethyl group introduced by these catalytic methods was typically placed at the position of the C–X (X = halides or boron) bond of the prefunctionalized arenes. Such a prefunctionalization of substrates limits the application of these methodologies for the late-stage modification of drug candidates for structureactivity relationship (SAR) studies. A strategy that could convert the C-H bond of arenes into C-CF₃ bonds is thus highly desirable (Scheme 1). Yu and co-workers recently reported a palladium-catalyzed directed ortho trifluoromethylation of arenes,^[13] but a method for the regioselective trifluoromethylation of arenes at a C-H bond without a directing group is unknown.

Recent work from our laboratory has demonstrated that the copper-catalyzed trifluoromethylation of aryl and vinyl boronic acids with Togni's reagent proceeded under very mild reaction conditions;^[14] Hartwig, Marder, Smith, Maleczka, and their co-workers have independently reported a series of reactions involving the iridium-catalyzed borylation of arenes and subsequent functionalization of the resulting boronic ester.^[15] The regioselectivity of the products from the iridiumcatalyzed borylation of arenes results from steric, rather than electronic control. For example, reactions of 1,3-disubstituted arenes under these conditions generate 3,5-disubstituted arene boronates. We envisioned that if these two catalytic reactions could be combined, a tandem C-H activation/ trifluoromethylation strategy could be realized for a highly meta-selective preparation of trifluoromethyl arenes from readily available 1,3-disubstituted arenes. Herein, we report the development of such a one-pot method for the trifluoromethylation of arenes using the combination of the iridiumcatalyzed borylation of arenes and copper-catalyzed trifluoromethylation of the resulting aryl boronates. The utility of this method is demonstrated by the efficient trifluoromethylation of some drug candidates through this one-pot meta trifluoromethylation. The introduction of a trifluoromethyl group into a molecule of interest at a late stage of a synthesis using mild reaction conditions could accelerate the discovery of a lead compound.

The trifluoromethylation of 4-biphenyl pinacolboronate with Togni's reagent^[16] was chosen at the start of our investigation as a model reaction to identify conditions for

the conversion of pinacolboronate esters into trifluoromethylarenes. We first examined if our previously developed reaction conditions, wherein 5 mol % of CuI and 10 mol % of 1,10-phenanthroline (**L1**) were used as the catalyst, were suitable for the reaction of 4-biphenyl pinacolboronate with Togni's reagent (Table 1). Not surprisingly, the reaction occurred much more slowly than those with aryl boronic acids. Only 80% conversion with a 58% yield (¹⁹F NMR spectroscopy) was observed after 24 hours at 45 °C (Table 1, entry 1). Switching the copper salt from CuI to copper(I)

Table 1: Optimization of the Cul/L1-catalyzed trifluoromethylation of 4-biphenyl pinacolboronate with Togni's reagent.^[a,b]



[a] Reaction conditions: 4-biphenyl pinacolboronate (0.2 mmol), Togni's reagent (0.2 mmol), CuX (10 mol%), ligand (20 mol%) and base (0.4 mmol) in specified solvent (1.0 mL) at 45 °C. [b] Reactions listed in entries 1–13 were conducted in Schlenk tubes with 5 mol% of CuX and 10 mol% of ligand, whereas reactions listed in entries 14–19 were conducted in sealed bombs. [c] Yields were determined by ¹⁹F NMR analysis of the crude reaction mixture with 1-fluoronaphthalene as an internal standard. [d] 10 mol% of the ligand was used.

thiophene-2-carboxylate (CuTc) resulted in a much faster reaction with full conversion after 24 hours (Table 1, entry 4); other copper salts such as $[Cu(OTf)]_2$ ·benzene and $[Cu-(MeCN)_4]PF_6$ did not improve the reaction yield (Table 1, entries 2 and 3). Interestingly, the reaction in the presence of water was much faster, probably because of the faster transmetalation of the aryl pinacolboronate to copper (Table 1, entry 5 versus 7 and entry 6 versus 10). For example, the reaction proceeded to full conversion after 48 hours in the presence of 1.0 equivalent of water when LiOH was used as the base, whereas only 28 % yield was observed in the absence of water (Table 1, entries 6 and 10). Reactions in the presence of other bases such as Li₂CO₃ or NaOH, however, afforded the desired product in less than 45 % yield (Table 1, entries 8-9). Finally, it was found that reactions in CH₂Cl₂ were much faster than those in other solvents and proceeded to full conversion after 4 hours at 45 °C to give the desired product in 97% yield, as determined by ¹⁹F NMR spectroscopy (Table 1, entries 10-16). Other dinitrogen ligands such as 2,9-dimethyl-1,10-phenanthroline (L2), 2,2'-bipyridine (L3), or tetramethylethylenediamine (TMEDA) were tested, but reactions under these conditions formed less than 81% of the trifluoromethyl biphenyl (Table 1, entries 17-19). In the absence of the copper catalyst, less than 5% of the trifluoromethylated product was observed.

With the optimized reaction conditions for the trifluoromethylation of aryl pinacolboronate in hand, we studied a tandem sequence to convert 1,3-disubstituted arenes or heteroarenes into the corresponding trifluoromethylated arenes or heteroarenes. The borylation of 1,3-disubstituted arenes was conducted with 0.7 equivalents of bis(pinacolato)diboron (B₂pin₂) in the presence of 0.25 mol% of [[Ir-(cod)OMe]₂] (cod=1,5-cyclooctadiene) and 0.5 mol% of di-*tert*-butylbipyridine (dtbpy) in THF at 80°C for 24 hours. The resulting arylboronate esters were converted into the trifluoromethylated arenes by evaporation of the volatile materials, dissolution of the residue in CH₂Cl₂, addition of 10 mol% of CuTc, 20 mol% of 1,10-phenanthroline, 1.1 equivalents of Togni's reagent, and 2.0 equivalents of LiOH·H₂O, and then heating at 45°C for 4–8 hours.

A variety of 1,3-disubstituted arenes were subjected to the C-H activation/trifluoromethylation conditions to give 5-trifluoromethyl-1,3-disubstituted the corresponding arenes in good to excellent yields, as summarized in Table 2. Arenes containing ester, protected phenoxy, chloride, and cyano groups were compatible with the reaction conditions and afforded the products in yields of 50-90% (Table 2, entries 1–8). In addition, the reactions of a number of heteroarenes gave the trifluoromethylated product with excellent selectivity and yields. For example, the reaction of a 2,6-disubstituted pyridine produced the corresponding 4-trifluoromethyl pyridine in 90% yield (Table 2, entry 9). Reactions of benzofuran and benzothiophene generated the 2-trifluoromethyl products in 72% and 75% yields, respectively (Table 2, entries 10 and 11). The reaction of Bocprotected indole gave the corresponding 3-trifluoromethylsubstituted product in good yield (Table 2, entry 12).

Late-stage modification of drug candidates is valuable for structure-activity relationship (SAR) studies since the complex target molecules are otherwise more challenging to obtain. We selected several pharmaceutical compounds to illustrate the advantage of late-stage trifluoromethylation compared to conventional synthesis to access complex trifluoromethylated molecules (Scheme 2). Biologically active molecules such as Vitamins B₃ and E, Vitamin E nicotinate, carbohydrates, and steroids are all compatible with the tandem C–H activation/trifluoromethylation procedure. It is worth mentioning that for all trifluoromethylated



Me

 Table 2:
 Trifluoromethylation of arenes by iridium-catalyzed borylation.

 1) 0.25 mol% [{Ir(cod)OMe}2]













[a] Yields of isolated product from the one-pot procedure (1.0 mmol scale). [b] Borylation with 1.0 mol% [$\{Ir(cod)OMe\}_2$] and 2.0 mol% dtbpy.



Scheme 2. Late-stage trifluoromethylation of complex small molecules.

© 2012 Wilev-VCH Verlag GmbH & Co. KGaA. Weinheim

compounds shown in Scheme 2, the trifluoromethylation was regioselective, and no constitutional isomers were observed in most cases. Only in the case of estrone were two isomers isolated in 1:1 ratio of inseparable isomers. The generally high regioselectivity of the sequential iridium-catalyzed C–H borylation and copper-catalyzed trifluoromethylation is an advantage over the other known trifluoromethylation reactions.

In summary, we have demonstrated the first coppercatalyzed trifluoromethylation of arylboronate esters and a sequential iridium-catalyzed C–H activation borylation and copper-catalyzed trifluoromethylation of arenes with a variety of functional groups. The advantage of this tandem procedure was demonstrated by application to a number of biologically active molecules. Mechanistic studies and synthetic applications of these transformations are ongoing in our laboratory.

Experimental Section

 $[{Ir(cod)(OMe)}_2]$ (1.7 mg, 0.25 mol %), dtbpy (1.45 mg, 0.540 mol%), and B₂pin₂ (186 mg, 0.730 mmol) were placed into an oven-dried sealed bomb. The bomb was evacuated and refilled with Ar three times. Under a positive flow of argon, dry THF (2.0 mL) and arene (1.0 mmol) were added. The reaction was stirred at 80 °C and monitored by GC/MS until the disappearance of the arene (24 h). After filtration through a short plug of celite, the volatiles were removed under vacuum and 4.0 mL of CH₂Cl₂ was added. The solution was transferred by a syringe into an oven-dried sealed bomb that contained CuTc (19 mg, 0.10 mmol), 1,10-phenanthroline (36 mg, 0.20 mmol), LiOH·H₂O (83 mg, 2.0 mmol), and Togni's reagent (363 mg, 1.10 mmol) under Ar. The reaction system was quickly degassed through three freeze/pump/thaw cycles and refilled with Ar. The reaction was stirred at 45 °C and monitored by ¹⁹F NMR spectroscopy until the disappearance of Togni's reagent (typically 8 h). Brine (25 mL) and CH_2Cl_2 (10 mL) were added and the organic phase was separated. The aqueous phase was extracted with CH₂Cl₂ $(5 \times 10 \text{ mL})$ and the combined organic phases were dried over anhydrous Na₂SO₄, and concentrated in vacuo. The product was purified by flash chromatography on silica gel with n-pentane, and further purified by Kugelrohr distillation.

Received: September 20, 2011 Published online: November 30, 2011

Keywords: C–H activation · copper · fluorine · homogeneous catalysis · iridium

- a) M. A. McClinton, Tetrahedron 1992, 48, 6555; b) J. H. Clark, D. Wails, T. W. Bastock, Aromatic Fluorination, CRC, Boca Raton, 1996; c) K. Müller, C. Faeh, F. Diederich, Science 2007, 317, 1881; d) J. P. Bégué, D. Bonnet-Delpon, Bioorganic and Medicinal Chemistry of Fluorine, Wiley-Interscience, New Jersey, 2008; e) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, Chem. Soc. Rev. 2008, 37, 320; f) K. L. Kirk, Org. Process Res. Dev. 2008, 12, 305; g) D. J. Burton, Z. Yang, Tetrahedron 1992, 48, 189; h) M. Schlosser, Angew. Chem. 2006, 118, 5558; Angew. Chem. Int. Ed. 2006, 45, 5432; i) S. Roy, B. t. Gregg, G. W. Gribble, V. Le, S. Roy, Tetrahedron 2011, 67, 2161.
- [2] D. T. Wong, K. W. Perry, F. P. Bymaster, Nat. Rev. Drug Discovery 2005, 4, 764.

- [3] P. Margot, F. Huggenberger, J. Amrein, B. Weiss, BCPC Conf-Pests. Dis. 1998, 2, 375.
- [4] G. W. A. Milne, CRC Handbook of Pesticides, CRC, Boca Raton, 1995.
- [5] a) W. R. Dolbier, *Chem. Rev.* **1996**, *96*, 3123; b) J. A. Ma, D. Cahard, *Chem. Rev.* **2008**, *108*, PR1; c) O. A. Tomashenko, V. V. Grushin, *Chem. Rev.* **2011**, *111*, 4475.
- [6] a) R. J. Lundgren, M. Stradiotto, Angew. Chem. 2010, 122, 9510; Angew. Chem. Int. Ed. 2010, 49, 9322; b) T. Knauber, F. G. R. Arikan, L. Gooßen, Chem. Eur. J. 2011, 17, 2689; c) V. V. Grushin, W. J. Marshall, J. Am. Chem. Soc. 2006, 128, 4632; d) V. V. Grushin, W. J. Marshall, J. Am. Chem. Soc. 2006, 128, 12644; e) N. D. Ball, J. W. Kampf, M. S. Sanford, J. Am. Chem. Soc. 2010, 132, 2878; f) J. Xu, D.-F. Luo, B. Xiao, Z.-J. Liu, T.-J. Gong, Y. Fu, L. Liu, Chem. Commun. 2011, 47, 4300; g) C.-P. Zhang, J. Cai, C.-B. Zhou, X.-P. Wang, X. Zheng, Y.-C. Gu, J.-C. Xiao, Chem. Commun. 2011, 47, 9516; h) C.-P. Zhang, Z.-L. Wang, Q.-Y. Chen, C.-T. Zhang, Y.-C. Gu, J.-C. Xiao, Angew. Chem. 2011, 123, 1936; Angew. Chem. Int. Ed. 2011, 50, 1896; i) H. Morimoto, T. Tsubogo, N. D. Litvinas, J. F. Hartwig, Angew. Chem. 2011, 123, 3877; Angew. Chem. Int. Ed. 2011, 50, 3793; j) X.-L. Zheng, G.-S. Liu, Chem. Eur. J. 2011, 15, 2751; k) O. A. Tomashenko, E. C. Escudero-Adán, M. M. Belmonte, V. V. Grushin, Angew. Chem. 2011, 123, 7797; Angew. Chem. Int. Ed. 2011, 50, 7655; l) Y.-M. Li, T. Chen, H.-F. Wang, R. Zhang, K. Jin, X.-N. Wang, C. Y. Duan, Synlett 2011, 1713; K. Araki, M. Inoue, K. Kawada (Jpn. Kokai Tokkyo Koho), JP 2009234921A 20091015. 2009.
- [7] M. Oishi, H. Kondo, H. Amii, Chem. Commun. 2009, 1909.
- [8] E. J. Cho, T. D. Senecal, T. Kinzel, Y. Zhang, D. A. Watson, S. L. Buchwald, *Science* **2010**, *328*, 1679.
- [9] L. Chu, F.-L. Qing, Org. Lett. 2010, 12, 5060; L. Chu, F.-L. Qing, J. Am. Chem. Soc. 2010, 132, 7262.
- [10] T. D. Senecal, A. T. Parsons, S. L. Buchwald, J. Org. Chem. 2011, 76, 1174.

- [11] a) F. Swarts, Bull. Acad. R. Belg. 1892, 24, 309; b) J. H. Simons, C. J. Lewis, J. Am. Chem. Soc. 1938, 60, 492; c) A. L. Henne, M. S. Newman, J. Am. Chem. Soc. 1938, 60, 1697; d) J. Riera, J. Castañer, J. Carilla, A. Robert, Tetrahedron Lett. 1989, 30, 3825; e) J. Castañer, J. Riera, J. Carilla, A. Robert, E. Molins, C. Miravitlles, J. Org. Chem. 1991, 56, 103.
- [12] a) Y. Kobayashi, I. Kumadaki, *Tetrahedron Lett.* 1969, 10, 4095;
 b) V. C. R. McLaughlin, J. Thrower, *Tetrahedron* 1969, 25, 5921;
 c) D. M. Wiemers, D. J. Burton, J. Am. Chem. Soc. 1986, 108, 832;
 d) G. E. Carr, R. D. Chambers, T. F. Holmes, D. G. Parker, J. Chem. Soc. Perkin Trans. 1 1988, 921;
 e) H. Suzuki, Y. Yoshida, A. Osuka, Chem. Lett. 1982, 135;
 f) H. Urata, T. Fuchikami, *Tetrahedron Lett.* 1991, 32, 91;
 g) Q.-Y. Chen, S.-W. Wu, Chem. Commun. 1989, 705;
 h) D.-B. Su, J.-X. Duan, Q.-Y. Chen, *Tetrahedron Lett.* 1991, 32, 7689;
 j) G. G. Dubinina, H. Furutachi, D. A. Vicic, J. Am. Chem. Soc. 2008, 130, 8600;
 k) Z. Weng, R. Lee, W. Jia, Y. Yuan, W. Wang, X. Feng, K. Huang, Organometallics 2011, 30, 3229.
- [13] X. Wang, L. Truesdale, J.-Q. Yu, J. Am. Chem. Soc. 2010, 132, 3648.
- [14] T. Liu, Q. Shen, Org. Lett. 2011, 13, 2342.
- [15] a) I. A. I. Mkhalid, J. H. Barnard, T. B. Marder, J. M. Murphy, J. F. Hartwig, *Chem. Rev.* 2010, *110*, 890; b) H. Y. Chen, S. Schlecht, T. C. Semple, J. F. Hartwig, *Science* 2000, *287*, 1995; c) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi, J. F. Hartwig, *J. Am. Chem. Soc.* 2002, *124*, 390; d) T. M. Boller, J. M. Murphy, M. Hapke, T. Ishiyama, N. Miyaura, J. F. Hartwig, *J. Am. Chem. Soc.* 2005, *127*, 14263; e) J. Y. Cho, C. N. Iverson, M. R. Smith, *J. Am. Chem. Soc.* 2000, *122*, 12868; f) J. Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka, M. R. Smith, *Science* 2002, *295*, 305.
- [16] a) P. Eisenberger, S. Gischig, A. Togni, *Chem. Eur. J.* 2006, *12*, 2579; b) P. Eisenberger, I. Kieltsch, N. Armanino, A. Togni, *Chem. Commun.* 2008, 1575.