

Thioether Synthesis


Copper-Catalyzed Oxidative Trifluoromethylthiolation of Aryl Boronic Acids with TMSCF_3 and Elemental Sulfur**

Chao Chen, Yan Xie, Lingling Chu, Ruo-Wen Wang, Xingang Zhang, and Feng-Ling Qing*

Fluorinated functional groups are key structural units found in various pharmaceuticals and agrochemicals.^[1] Approximately 30% of all agrochemicals and 20% of all pharmaceuticals on the market contain fluorine. Among these substituents, the trifluoromethylthio group ($\text{CF}_3\text{S}-$), especially as an aromatic substituent, plays an important role because of its strong electron-withdrawing effect and high lipophilicity. These characteristics are similar to those of trifluoromethyl (CF_3-) and trifluoromethoxy ($\text{CF}_3\text{O}-$) groups.^[2] Additionally, aryl trifluoromethyl thioethers (CF_3SAr) are also key intermediates in the preparation of trifluoromethyl sulfoxide and sulfone, which are important trifluoromethylation reagents.^[3] Although impressive progress has been made in the trifluoromethylation of arenes in the past several years,^[4–7] only a few methods are available for the synthesis of aryl trifluoromethyl thioethers.^[8,9] Generally, aryl trifluoromethyl thioethers are prepared either by a nucleophilic reaction of trifluoromethylthiolate with aryl halides,^[8] or by a nucleophilic or radical reaction of aryl sulfides and disulfides with a trifluoromethylation reagent.^[9] However, these methods are variously limited by a combination of high temperatures, expensive reagents, and low reactivity with electron-rich aromatic groups. Thus, the development of general, safe, and efficient methods to access aryl trifluoromethyl thioethers is highly desirable. Very recently, Buchwald reported a palladium-catalyzed trifluoromethylthiolation of aryl bromides with CF_3SAg .^[10] This breakthrough for the preparation of ArSCF_3 is highly efficient and compatible with a variety of functional groups. However, from the point view of cost-effectiveness and synthetic convenience, using readily available and inexpensive catalysts and fluorinated reagents, such as copper and (trifluoromethyl)trimethylsilane

(the Ruppert–Prakash reagent, TMSCF_3), to access aryl trifluoromethyl thioethers would be an attractive alternative.

The present study was inspired by our own and Buchwald's recent investigations into the copper mediated oxidative trifluoromethylation of arylboronic acid with TMSCF_3 ,^[11] as well as Karlin's observation of the formation of a stable copper disulfide complex from the reaction of elemental sulfur (S_8) with a Cu^{I} complex.^[12] We hypothesized that a Cu^{I} disulfide complex generated in situ (**II**; Scheme 1) would



Scheme 1. Copper(I)-catalyzed formation of aryl trifluoromethyl thioether from aryl boronic acid, TMSCF_3 , and S_8 .

react with aryl boronic acid to give intermediate **III** (Path A), which would subsequently react with TMSCF_3 , providing the key intermediate complex $\text{L}_n\text{Cu}(\text{CF}_3)(\text{ArS})$ (**V**). Finally, oxidation of complex **V** to $\text{L}_n\text{Cu}^{\text{III}}(\text{CF}_3)(\text{ArS})$,^[13] followed by reductive elimination would lead to the expected aryl trifluoromethyl thioether. Alternatively, intermediate **IV** could be formed by the reaction of TMSCF_3 with complex **II** (Path B). The desired product might still be obtained from oxidation of key intermediate **VI**, generated from complex **IV**.

Herein, we report the first example of the copper-catalyzed oxidative trifluoromethylthiolation of arylboronic acids with TMSCF_3 and elemental sulfur at room temperature. The notable features of this reaction are its high efficiency, excellent functional group compatibility (bromide is also compatible), operational simplicity, inexpensive catalyst, easily accessible starting materials, and mild reaction conditions.

In accordance with our hypothesis, we began this study by reacting phenyl boronic acid **1**, TMSCF_3 , and S_8 in the presence of different copper salts, bases, and oxidants to optimize the reaction conditions. To our delight, when the

[*] C. Chen, Y. Xie, L. Chu, Dr. R.-W. Wang, Prof. Dr. X. Zhang, Prof. Dr. F.-L. Qing
Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences
345 Lingling Lu, Shanghai 200032 (China)
E-mail: flq@mail.sioc.ac.cn


Prof. Dr. F.-L. Qing
College of Chemistry, Chemical Engineering and Biotechnology, Donghua University
2999 North Renmin Lu, Shanghai, 201620 (China)

[**] This work was supported by the National Natural Science Foundation of China (21072028, 20832008) and the National Basic Research Program of China (2012CB21600). The authors thank Prof. Qilong Shen of the Shanghai Institute of Organic Chemistry for helpful discussions.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201108663>.

reaction was conducted with CuI (1.0 equiv) and 1,10-phenanthroline (phen; 1.1 equiv) in the presence of Ag₂CO₃ (2.0 equiv) as oxidant, along with K₃PO₄ (1.0 equiv) and KF (1.0 equiv) as bases, and 4 Å molecular sieves (4 Å M.S.), in DMF at room temperature, phenyl trifluoromethyl thioether **2a** was formed in 33 % yield (Table 1, entry 1). The undesired

Table 1: Optimization of the copper-catalyzed oxidative trifluoromethylthiolation of phenyl boronic acid.^[a]


				
Entry	CuX	Base	Oxidant	Yield [%] ^[b]
1 ^[c]	CuI	K ₃ PO ₄ /KF	Ag ₂ CO ₃	33
2 ^[c,d]	CuI	K ₃ PO ₄ /KF	Ag ₂ CO ₃	0
3 ^[c]	—	K ₃ PO ₄ /KF	Ag ₂ CO ₃	0
4 ^[c]	CuI	K ₃ PO ₄ /KF	—	0
5	CuI	KF	Ag ₂ CO ₃	23
6	CuI	K ₃ PO ₄	Ag ₂ CO ₃	43
7	CuCl	K ₃ PO ₄	Ag ₂ CO ₃	19
8	CuBr	K ₃ PO ₄	Ag ₂ CO ₃	19
9	CuOAc	K ₃ PO ₄	Ag ₂ CO ₃	21
10	(CuOTf) ₂ C ₆ H ₆	K ₃ PO ₄	Ag ₂ CO ₃	18
11	K ₄ [Cu ₂ (CN) ₆]	K ₃ PO ₄	Ag ₂ CO ₃	9
12	CuCN	K ₃ PO ₄	Ag ₂ CO ₃	61
13	CuSCN	K ₃ PO ₄	Ag ₂ CO ₃	72
14 ^[e]	CuSCN	K ₃ PO ₄	Ag ₂ CO ₃	85
15 ^[e,f]	CuSCN	K ₃ PO ₄	Ag ₂ CO ₃	78
16 ^[e,g]	CuSCN	K ₃ PO ₄	Ag ₂ CO ₃	95

[a] Reaction conditions: **1** (0.2 mmol), S₈ (0.6 mmol), TMSCF₃ (0.6 mmol), CuX (0.2 mmol), phen (0.22 mmol), base (0.4 mmol), oxidant (0.4 mmol), 4 Å M.S. (50 mg), DMF (4 mL), 24 h, N₂, RT.
[b] Determined by ¹⁹F NMR. [c] 1.0 equiv of K₃PO₄ and 1.0 equiv of KF.
[d] Without phen. [e] With 5.0 equiv of TMSCF₃ and 3.0 equiv of K₃PO₄.
[f] With 10 mol % of CuSCN and 10 mol % phen. [g] With 10 mol % of CuSCN and 20 mol % phen.

products phenyl disulfide (**3**) and cyclic trimeric phenyl boronic acid anhydride (**4**) were observed in the reaction mixture. Furthermore, reactions without CuI, phen, or Ag₂CO₃ failed to afford desired product **2a**, thus showing the pivotal role of these reagents in the reaction (entries 2–4). Interestingly, when K₃PO₄ was used as both base and initiator for TMSCF₃, the yield increased to 43 % (entry 6). The formation of **4** indicated that the intermediate **III** might be generated slowly. Likewise, the formation of **3** arose from the homocoupling of copper complex **III**; an indication that the reaction of **III** with TMSCF₃ to form **V** was even slower. Therefore, if a suitable copper salt was used to accelerate step 1 (the formation of copper complexes **III** from intermediate **II**) and step 2 (formation of **V** from **III**), and a proper oxidant was employed to facilitate the reductive elimination of **V** (Scheme 1), the formation of byproducts **3** and **4** would be inhibited and thus lead to improved yield of desired product **2a**. Accordingly, different copper salts and oxidants were examined to improve the reaction efficiency. After many attempts, CuSCN and Ag₂CO₃ were found to be the best choices, providing **2a** in 72 % yield along with a small amount of phenyl disulfide **3** (Table 1, entry 13). To further inhibit the

formation of **3**, 5.0 equivalents of TMSCF₃ was used, further improving the yield to 85 % (Table 1, entry 14). Notably, a catalytic amount of CuSCN (10 mol %) and phen (10 mol %) still furnished **2a** in a comparable yield (78 %). Further increasing the loading of phen to 20 mol % provided the optimum yield of **2a**, 95 % (entry 16).

With the optimum reaction conditions (Table 1, entry 16) determined, the substrate scope of the reaction was then investigated (Scheme 2). The mild reaction conditions




Scheme 2. Scope of the Copper(I)-catalyzed oxidative trifluoromethylthiolation of aryl boronic acids. Reaction conditions: **1** (0.2 mmol), S₈ (0.6 mmol), TMSCF₃ (1.0 mmol), CuSCN (0.02 mmol), Phen (0.04 mmol), K₃PO₄ (0.6 mmol), Ag₂CO₃ (0.4 mmol), 4 Å M.S. (50 mg), DMF (3 mL), 24 h, N₂, RT. Yields shown are of isolated products.

allowed for the trifluoromethylthiolation of aryl boronic acids containing a range of functional groups, including ester, unprotected amide, ketone, nitrile, and sulfonyl groups. Notably, even for the substrates bearing bromide or vinyl group, which are reactive in the presence of a Pd⁰ catalyst, good yields were still obtained, thus providing opportunities for further transformation.

To investigate the mechanism of the oxidative trifluoromethylthiolation, several experiments were performed. First, the reaction of 3,4,5-trimethoxyphenyl boronic acid (**1h**) with S₈, CuSCN, and phen in the presence of K₃PO₄ in [D₇]DMF at room temperature was monitored by ¹H NMR spectroscopy (see the Supporting Information). A new species corresponding to intermediate **III** was observed. GC/MS analysis of the

reaction mixture showed a peak at $m/z = 199$, which was assigned to $\text{Ph}(\text{OMe})_3\text{S}^-$, suggesting that the formation of intermediate **III** from boronic acid **1**, S_8 , and Cu^{I} is a reasonable proposal. In contrast, neither $\text{CF}_3\text{S-Cu}$ nor $\text{CF}_3\text{-Cu}$ was observed when a mixture of TMSCF_3 , CuSCN , phen, S_8 , and K_3PO_4 in DMF was stirred at room temperature (see the Supporting Information). Furthermore, the reaction of copper 4-methoxyphenyl thiolate^[14] with TMSCF_3 in the presence of Ag_2CO_3 and K_3PO_4 at room temperature proceeded smoothly to give **2q** as the only fluorinated product in 59% yield (determined by ^{19}F NMR; Eq. (1) of Scheme 3). However, only trace **2q** was detected in the



Scheme 3. Synthesis of trifluoromethyl 4-methoxyphenyl thioether.

reaction of copper trifluoromethyl thiolate (CuCF_3)^[15] with 4-methoxyphenyl boronic acid (Eq. (2) of Scheme 3). Based on these results, we propose that path A (Scheme 1) is the likely pathway for the copper-catalyzed oxidative trifluoromethylthiolation of aryl boronic acids with TMSCF_3 and S_8 .^[16]

In summary, we have developed the first copper(I)-catalyzed oxidative trifluoromethylthiolation of arylboronic acid using TMSCF_3 and S_8 at room temperature. This reaction provides an efficient and convenient method for the preparation of aryl trifluoromethyl thioethers. An organocopper disulfide complex was proposed as the key intermediate in the catalytic cycle.

Experimental Section

General procedure for oxidative trifluoromethylthiolation: In a glove box, 4 Å powdered molecular sieves (50 mg) and K_3PO_4 (128 mg, 0.6 mmol, 3.0 equiv) were added to a test tube equipped with a magnetic stir bar. The vessel was sealed with a septum and flame-dried under vacuum. The tube was cooled to room temperature and backfilled with argon. Then CuSCN (3 mg, 0.02 mmol, 0.1 equiv), 1,10-phenanthroline (8 mg, 0.04 mmol, 0.2 equiv), S_8 (20 mg, 0.6 mmol, 3.0 equiv), aryl boronic acid **1** (0.2 mmol, 1.0 equiv), and Ag_2CO_3 (110 mg, 0.4 mmol, 2.0 equiv) were quickly added under a N_2 atmosphere. The tube was then evacuated and backfilled with argon gas. Freshly distilled DMF (5 mL) and TMSCF_3 (150 μL , 1.0 mmol, 5.0 equiv) were then added to the reaction tube by syringe, which was then placed under a balloon of N_2 and stirred vigorously for 24 h. Fluorobenzene (56 μL , 0.6 mmol) was added as an internal standard, and the yield of the crude reaction was measured by ^{19}F NMR before workup. The reaction solution was filtered through Celite on silica and the filter cake was washed with diethyl ether. The filtrate was then washed with brine and concentrated. The residue was purified by

silica gel column chromatography with hexane to provide pure aryl trifluoromethyl thioether.

Received: December 8, 2011

Published online: January 27, 2012

Keywords: aryl boronic acids · copper catalysis · oxidative coupling · sulfur · trifluoromethylthiolation

- [1] a) T. Yamazaki, T. Taguchi, I. Ojima in *Fluorine in Medicinal Chemistry and Chemical Biology* (Ed.: I. Ojima), Wiley-Blackwell, Chichester, **2009**; b) P. Jeschke, *ChemBioChem* **2004**, *5*, 570; c) K. Muller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881; d) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320; e) W. K. Hagmann, *J. Med. Chem.* **2008**, *51*, 4359; f) T. Furuya, A. S. Kamlet, T. Ritter, *Nature* **2011**, *473*, 470.
- [2] a) S. Manteau, S. Pazenok, J. P. Vors, F. R. Leroux, *J. Fluorine Chem.* **2010**, *131*, 140; b) F. Leroux, P. Jeschke, M. Schlosser, *Chem. Rev.* **2005**, *105*, 827; c) C. Hansch, A. Leo, R. W. Taft, *Chem. Rev.* **1991**, *91*, 165; d) A. Leo, C. Hansch, D. Elkins, *Chem. Rev.* **1971**, *71*, 525.
- [3] a) L. Xu, J. Cheng, M. L. Trudell, *J. Org. Chem.* **2003**, *68*, 5388; b) S.-Y. Tang, P. Zhong, Q.-L. Lin, *J. Fluorine Chem.* **2007**, *128*, 636; c) G. K. S. Prakash, J. Hu, G. A. Olah, *Org. Lett.* **2003**, *5*, 3253; d) Y. Zhao, J. Zhu, C. Ni, J. Hu, *Synthesis* **2010**, 1899.
- [4] For conventional examples of arene trifluoromethylation reactions, see: a) C. Wakselman, M. Tordeux, *J. Chem. Soc. Chem. Commun.* **1987**, 1701; b) B. R. Langlois, E. Laurent, M. Roidot, *Tetrahedron Lett.* **1991**, *32*, 7525; c) Y. Macé, B. Raymondeau, C. Pradet, J. C. Blazewski, E. Magnier, *Eur. J. Org. Chem.* **2009**, 1390; d) M. S. Wiehn, E. V. Vinogradova, A. Togni, *J. Fluorine Chem.* **2010**, *131*, 951; e) Y. Ji, T. Bruecki, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond, P. S. Baran, *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 14411.
- [5] For recent examples of palladium-based arene trifluoromethylation reactions, see: a) V. V. Grushin, W. J. Marshall, *J. Am. Chem. Soc.* **2006**, *128*, 12644; b) N. D. Ball, J. W. Kampf, M. S. Sanford, *J. Am. Chem. Soc.* **2010**, *132*, 2878; c) Y. Ye, N. D. Ball, J. W. Kampf, M. S. Sanford, *J. Am. Chem. Soc.* **2010**, *132*, 14682; d) E. J. Cho, T. D. Senecal, T. Kinzel, Y. Zhang, D. A. Watson, S. L. Buchwald, *Science* **2010**, *328*, 1679; e) X. Wang, L. Truesdale, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, *132*, 3648; f) B. S. Samant, G. W. Kabalka, *Chem. Commun.* **2011**, *47*, 7236; g) N. D. Ball, J. B. Gary, Y. Ye, M. S. Sanford, *J. Am. Chem. Soc.* **2011**, *133*, 7577; h) X. Mu, S. Chen, X. Zhen, G. Liu, *Chem. Eur. J.* **2011**, *17*, 6039; i) R. N. Loy, M. S. Sanford, *Org. Lett.* **2011**, *13*, 2548; j) E. J. Cho, S. L. Buchwald, *Org. Lett.* **2011**, *13*, 6552; k) Y. Zhao, J. Hu, *Angew. Chem.* **2012**, *124*, 1057; *Angew. Chem. Int. Ed.* **2012**, *51*, 1033.
- [6] For recent examples of copper-based arene trifluoromethylation reactions, see: a) G. G. Dubinina, H. Furutachi, D. A. Vicić, *J. Am. Chem. Soc.* **2008**, *130*, 8600; b) M. Oishi, H. Kondo, H. Amii, *Chem. Commun.* **2009**, 1909; c) K. A. McReynolds, R. S. Lewis, L. K. G. Ackerman, G. G. Dubinina, W. W. Brennessel, D. A. Vicić, *J. Fluorine Chem.* **2010**, *131*, 1108; d) R. Shimizu, H. Egami, T. Nagi, J. Chae, Y. Hamashima, M. Sodeoka, *Tetrahedron Lett.* **2010**, *51*, 5947; e) C.-P. Zhang, Z.-L. Wang, Q.-Y. Chen, C.-T. Zhang, Y.-C. Gu, J.-C. Xiao, *Angew. Chem.* **2011**, *123*, 1936; *Angew. Chem. Int. Ed.* **2011**, *50*, 1896; f) H. Morimoto, T. Tsubogo, N. D. Litvinas, J. F. Hartwig, *Angew. Chem.* **2011**, *123*, 3877; *Angew. Chem. Int. Ed.* **2011**, *50*, 3793; g) O. A. Tomashenko, E. C. Escudero-Adan, M. M. Belmonte, V. V. Grushin, *Angew. Chem.* **2011**, *123*, 7797; *Angew. Chem. Int. Ed.* **2011**, *50*, 7655; h) T. Knauber, F. Arikan, G.-V. Rösenthaller, L. J. Goßen, *Chem. Eur. J.* **2011**, *17*, 2689; i) H. Kondo, M. Oishi, K. Fujikawa, H. Amii, *Adv. Synth. Catal.*

- 2011, 353, 1247; j) Y. Li, T. Chen, H. Wang, R. Zhang, K. Jin, X. Wang, C. Duan, *Synlett* **2011**, 1713; k) I. Popov, S. Lindeman, O. Daugulis, *J. Am. Chem. Soc.* **2011**, 133, 9286; l) J. Xu, D.-F. Luo, B. Xiao, Z.-J. Liu, T.-J. Gong, Y. Fu, L. Liu, *Chem. Commun.* **2011**, 47, 4300; m) T. Liu, Q. Shen, *Org. Lett.* **2011**, 13, 2342; n) C. P. Zhang, J. Cai, C.-B. Zhou, X. P. Wang, X. Zheng, Y. C. Gu, J. C. Xiao, *Chem. Commun.* **2011**, 47, 9516; o) H. Kawai, T. Furukawa, Y. Nomura, E. Tokunaga, N. Shibata, *Org. Lett.* **2011**, 13, 3596; p) M. M. Kremlev, A. I. Mushta, W. Tyrre, Y. L. Yagupolskii, *J. Fluorine Chem.* **2012**, 133, 67; q) N. D. Litvinas, P. S. Fier, J. F. Hartwig, *Angew. Chem.* **2012**, 124, 551; *Angew. Chem. Int. Ed.* **2012**, 51, 536; r) T. Liu, X. Shao, Y. Wu, Q. Shen, *Angew. Chem.* **2012**, 124, 555; *Angew. Chem. Int. Ed.* **2012**, 51, 540.
- [7] For recent examples of silver-mediated arene trifluoromethylation and trifluoromethoxylation reactions, see: a) Z. Weng, R. Lee, W. Jia, Y. Yuan, W. Wang, X. Feng, K.-W. Huang, *Organometallics* **2011**, 30, 3229; b) Y. Ye, S. H. Lee, M. S. Sanford, *Org. Lett.* **2011**, 13, 5464; c) C. Huang, T. Liang, S. Harada, E. Lee, T. Ritter, *J. Am. Chem. Soc.* **2011**, 133, 13308.
- [8] a) L. M. Yagupolskii, N. V. Kondratenko, V. P. Sabur, *Synthesis* **1975**, 721; b) D. V. Remy, K. E. Rittle, C. A. Hunt, M. B. Freedman, *J. Org. Chem.* **1976**, 41, 1644; c) Q. Y. Chen, J. X. Duan, *J. Chem. Soc. Chem. Commun.* **1993**, 918; d) D. J. Adams, A. Goddard, J. H. Clark, D. J. Macquarrie, *Chem. Commun.* **2000**, 987; e) D. J. Adams, J. H. Clark, *J. Org. Chem.* **2000**, 65, 1456; f) W. Tyrre, D. Naumann, B. Hoge, Y. L. Yagupolskii, *J. Fluorine Chem.* **2003**, 119, 101.
- [9] a) V. N. Boiko, G. M. Shchupak, L. M. Yagupolskii, *J. Org. Chem. USSR* **1977**, 13, 972; b) C. Wakselman, M. Tordeux, *J. Org. Chem.* **1985**, 50, 4047; c) C. Wakselman, M. Tordeux, J.-L. Clavel, B. Langlois, *J. Chem. Soc. Chem. Commun.* **1991**, 993; d) T. Umemoto, S. Ishihara, *J. Am. Chem. Soc.* **1993**, 115, 2156; e) T. Billard, B. R. Langlois, *Tetrahedron Lett.* **1996**, 37, 6865; f) B. Quiclet-Sire, R. N. Saicic, S. Z. Zars, *Tetrahedron Lett.* **1996**, 37, 9057; g) T. Billard, N. Roques, B. R. Langlois, *J. Org. Chem.* **1999**, 64, 3813; h) N. Roques, *J. Fluorine Chem.* **2001**, 107, 311; i) G. Blond, T. Billard, B. R. Langlois, *Tetrahedron Lett.* **2001**, 42, 2473; j) C. Pooput, M. Medebielle, W. R. Dolbier, *Org. Lett.* **2004**, 6, 301; k) C. Pooput, W. R. Dolbier, M. Medebielle, *J. Org. Chem.* **2006**, 71, 3564; l) I. Kietlsch, P. Eisenberger, A. Togni, *Angew. Chem.* **2007**, 119, 768; *Angew. Chem. Int. Ed.* **2007**, 46, 754; m) A. Harsányi, E. Dorko, A. Csapo, T. Bako, C. Peltz, J. Rabai, *J. Fluorine Chem.* **2011**, 132, 1241.
- [10] G. Teverovskiy, D. S. Surry, S. L. Buchwald, *Angew. Chem.* **2011**, 123, 7450; *Angew. Chem. Int. Ed.* **2011**, 50, 7312.
- [11] a) L. Chu, F. L. Qing, *Org. Lett.* **2010**, 12, 5060; b) T. D. Senecal, A. T. Parsons, S. L. Buchwald, *J. Org. Chem.* **2011**, 76, 1174.
- [12] M. E. Helton, P. Chen, P. P. Paul, Z. TyeKlar, R. D. Sommer, L. N. Zakharov, A. L. Rheingold, E. I. Solomon, K. D. Karlin, *J. Am. Chem. Soc.* **2003**, 125, 1160.
- [13] a) X. Ribas, D. A. Jackson, B. Donnadieu, J. Mahia, T. Parella, R. Xifra, B. Hedman, K. O. Hodgson, A. Llobet, T. D. P. Stack, *Angew. Chem.* **2002**, 114, 3117; *Angew. Chem. Int. Ed.* **2002**, 41, 2991; b) R. Xifra, X. Ribas, A. Llobet, A. Poater, M. Duran, M. Solà, T. D. P. Stack, J. Benet-Buchholz, B. Donnadieu, J. Mahía, T. Parella, *Chem. Eur. J.* **2005**, 11, 5146; c) L. M. Huffman, S. S. Stahl, *J. Am. Chem. Soc.* **2008**, 130, 9196; d) A. E. King, T. C. Brunold, S. S. Stahl, *J. Am. Chem. Soc.* **2009**, 131, 5044; e) A. E. King, L. M. Huffman, A. Casitas, M. Costas, X. Ribas, S. S. Stahl, *J. Am. Chem. Soc.* **2010**, 132, 12068; f) A. Casitas, A. E. King, T. Parella, M. Costas, S. S. Stahl, X. Ribas, *Chem. Sci.* **2010**, 1, 326; g) X. Ribas, C. Calle, A. Poater, A. Casitas, L. Gómez, R. Xifra, T. Parella, J. Benet-Buchholz, A. Schweiger, G. Mitrikas, M. Solà, A. Llobet, T. D. P. Stack, *J. Am. Chem. Soc.* **2010**, 132, 12299; h) R. Giri, J. F. Hartwig, *J. Am. Chem. Soc.* **2010**, 132, 15860.
- [14] G. H. Posner, D. J. Brunelle, L. Sinoway, *Synthesis* **1974**, 662.
- [15] CuSCF₃ was purchased from TCI.
- [16] The formation of trifluoromethylthiolated arenes by the reaction of AgSCF₃ with aryl boronic acids was also ruled out; see the Supporting Information.