

Activated Nitriles in Heterocyclic Synthesis: Synthesis of Several New Coumarin Derivatives

Sadek Abdou, Sherif Mahmoud Fahmy, Mahmoud M. Khader,
and Mohamed Hilmy Elnagdi*

Department of Chemistry, Faculty of Science, Cairo University, Giza,
A. R. Egypt

(Received 30 October 1981. Accepted 25 November 1981)

Several new benzopyran and benzopyranopyridine derivatives were synthesized via condensation of 2-amino-1,1,3-tricyanopropene (**1**) and diethyl 2-amino-1-cyanopropene-1,3-dicarboxylate (**2**) with salicylaldehyde and treatment of the resulted products with active methylene reagents. The spectroscopic data of the synthesized compounds are reported and discussed.

(Keywords: Benzopyrano[2,3-c]pyridines; Enamino nitriles; Iminocoumarins)

Aktiviert Nitrile in der Heterocyclen-Synthese: Darstellung einiger neuer Cumarinderivate

Einige neue Benzopyran- und Benzopyranopyridin-derivate wurden durch Kondensation von 2-Amino-1,1,3-tricyanopropen (**1**) und 2-Amino-1-cyanopropen-1,3-dicarbonensäuredimethylester (**2**) mit Salicylaldehyd und Behandlung der Reaktionsprodukte mit aktiven Methylenreagenzien dargestellt. Die spektroskopischen Daten der neuen Verbindungen werden diskutiert.

Introduction

As a part of our program directed to the development of new procedures for the synthesis of heterocyclic derivatives from readily accessible activated nitrile derivatives¹⁻³ we have investigated the reaction of salicylaldehyde with 2-amino-1,1,3-tricyano-propene (**1**)⁴ and diethyl 2-amino-1-cyanopropene-1,3-dicarboxylate (**2**)⁵.

Results and Discussion


*Junek*⁶ has reported that salicylaldehyde reacts with **1** to yield the benzopyrano[3,4-*c*]pyridine derivative **3**. Substituted salicylaldehyde have been reported⁶ to afford the iminocoumarin derivatives **4**. In our laboratory, however, salicylaldehyde reacted with **1** to yield a product of the molecular formula $C_{13}H_{10}O_2N_4$. The IR spectrum of this product revealed absorption peaks for NH_2 , CN and CO groups. The presence of phenolic OH was indicated by the development of a deep green colour on adding $FeCl_3$ to a suspension of the reaction product in ethanol. Structure **5** was thus suggested for the reaction product. The formation of **5** in this reaction is assumed to proceed *via* the sequence demonstrated in Scheme 1. Two possible geometrical isomers for **5** were considered. The reaction product was considered to exist mainly in the **5a** form based on its ready conversion into the benzopyrano[2,3-*c*]pyridine derivative **6**, via loss of ammonia, on treatment with ethanolic hydrochloric acid.

In contrast to the behaviour of **1**, compound **2** reacted with salicylaldehyde to yield the coumarin derivative **7**. Although geometrical isomers are possible for **7**, structure **7** was established for this product based on 1H NMR which revealed the coumarin H-4 proton at $\delta 6.22$ ppm—upfield shifted by $\delta 2$ ppm with regard to its expected position. This upfield shift can only be rationalized in terms of shielding by the anisotropy of the cyano group. Compound **7** could be readily converted into several—otherwise difficultly accessible—coumarin derivatives (Scheme 2). Thus, **7** reacted readily with hydrazine hydrate and with phenylhydrazine to yield the 3-coumarinyl aminomethylene-pyrazolone derivatives **8a, b**. The formation of **8a, b** from **7** is assumed to proceed *via* the addition of the reagent to the cyano group and cyclization *via* elimination of ethanol. This is in contrast to the reported addition of hydrazines to the double bond in enamino esters and enamino nitriles under similar conditions⁷. The inactivity of the double bond in **7** toward hydrazines is rationalized by the decrement of its reactivity by conjugation with the coumarin ring.

Compound **7** readily reacted with active methylene heterocyclic derivatives to yield 1:1 adducts. Thus, with the 2-pyrazolin-5-one derivatives **9a, b** and with the 2-isooxazolin-5-one derivative **9c**, the 4-azolyl-substituted dihydrocoumarins **10a-c** were formed in good yields. The reactivity of the double bond in **7** in *Michael* reactions is analogous to the activity of the double bond in coumarins and in benzopyrans in similar reactions^{8,9}.

7 condensed with malononitrile to yield a product of the molecular formula $C_{16}H_8O_3N_4$. The IR spectrum of this product revealed ab-


Scheme 1



sorption peaks at 3400, 1720, and 2220, 2210 cm^{-1} for amino, CO and two CN groups. Structure **11** and **12** were considered for the reaction product and a mechanism could be suggested to account for the formation of both products. Structure **11** was established for the reaction product based on ^1H NMR which revealed a multiplet for two nonaromatic protons at δ 4 ppm.

Similarly, **7** reacted with benzoylacetonitrile to yield the pyridinyl-coumarin derivative **13** (or possible tautomers). The formation of **13** is assumed to proceed *via* initial addition to the double bond. In contrast, ethyl cyanoacetate reacted with **7** to yield the pyridine derivative **14** (cf. Scheme 2).

Scheme 2



The behaviour of **1** and **2** towards the action of other aldehydes is now being investigated. The work offers a new route for the synthesis of pyridine derivatives and will be the subject of a subsequent communication.

Experimental

Melting points are uncorrected. IR spectra were recorded (KBr) on a Pye Unicam SP-1000. ¹H NMR spectra were obtained on an EM-360 MHz spectrometer in *DMSO* using *TMS* as internal standard and chemical shifts are expressed as δ ppm. Analytical data were performed by the microanalytical unit, Cairo University.

Reactions of **1** and **2** with Salicylaldehyde

To a solution of each of **1** and **2** (0.01 mol) in ethanol (30 ml), salicylaldehyde (0.01 mol) and one drop of piperidine were added. The reaction mixture was

Table 1. *Synthetic data for compounds 5 a, 6, 7, 8 a, b, 10 a-c, 11, 13 and 14*

Comp.	colour	m. p. °C	Yield %	Mol. Formula*
5 a	canary yellow	>300	96	C ₁₃ H ₁₀ O ₂ N ₄
6	yellow	277	70	C ₁₃ H ₇ O ₂ N ₃
7	yellow	235	85	C ₁₅ H ₁₂ O ₄ N ₂
8 a	orange	>280	65	C ₁₃ H ₁₀ O ₃ N ₄
8 b	brown	205	60	C ₁₉ H ₁₄ O ₃ N ₄
10 a	pale yellow	268	65	C ₁₉ H ₁₆ O ₃ N ₄
10 b	orange	278	60	C ₂₅ H ₂₀ O ₃ N ₄
10 c	red	240	70	C ₂₄ H ₁₇ O ₆ N ₃
11	pale yellow	>300	85	C ₁₆ H ₈ O ₃ N ₄
13	canary yellow	246	50	C ₂₄ H ₁₈ O ₆ N ₂
14	orange	217	70	C ₂₀ H ₁₆ O ₆ N ₂

* All compounds gave satisfactory elemental analyses (C, H, N).

refluxed for 5 min in case of **1** and for 30 min in case of **2**. The solid product so formed was collected by filtration and crystallized from *DMF*/H₂O (cf. Tables 1 and 2 for data of **5 a** and **7**).

Reaction of 5 a with Ethanolic Hydrochloric Acid

A mixture of **5 a** (1 g), conc. HCl (10 ml) and ethanol (30 ml) was boiled under reflux for 2 h. The reaction mixture was then cooled and neutralized with sodium carbonate solution. The solid thus formed was filtered off and washed with water. Crystallization from *DMF*/H₂O gave **6** (cf. Tables 1 and 2).

Reactions of 7 with Hydrazines

A solution of **7** (0.01 mol) in *DMF* (20 ml) was treated with hydrazine hydrate or phenylhydrazine (0.01 mol). The reaction mixture was boiled under reflux for 2 h, then evaporated in vacuo. The remaining product was triturated with ethanol, collected by filtration and crystallized from *DMF*/H₂O to give **8 a** or **b** (cf. Tables 1 and 2).

Reactions of 7 with Active Methylene Heterocyclic Derivatives

A solution of **7** (0.01 mol) and each of the active methylene heterocyclic derivatives **9 a-c** (0.01 mol) in *DMF* (30 ml) was boiled under reflux for 3 h. The solvent was then evaporated in vacuo and the residue triturated with ethanol. Crystallization of the residue from *DMF*/H₂O gave **10 a-c** (Tables 1 and 2).

Table 2. IR and ^1H NMR data

Comp.	IR; cm^{-1} ; selected bands	^1H NMR; δ ppm
5a*	3490 (OH); 3390, 3370, 3250 and 3100 (NH_2 groups); 2290 (CH); 2210 (conjugated CN); 1635 (ring CO) and 1605 (C=N)	
6	3430 ~ 3300 (chelated NH_2); 2210 (CN); 1660 ~ 1630 (ring CO and NH_2 deformation)	5.6 (br, 2 H, NH_2) and 7.0 ~ 8.0 (m, 5 H, aromatic protons)
7	3420, 3300 (NH_2); 3050 (aromatic CH); 2980 ~ 2960 (CH_2 and CH_3); 2210 (CN); 1710 (conjugated ester CO); 1685 (ring CO) and 1630 (NH_2 deformation)	1.16 (t, 3 H, CH_3); 6.22 (s, 1 H, benzopyran H-4); 4.06 (q, 2 H, CH_2); 7.2 ~ 7.85 (m, 4 H, aromatic protons) and 8.8 ~ 9.0 (m, br, 2 H, NH_2)
8a*	3400 ~ 2400 (chelated NH) and 1680 ~ 1630 (chelated CO groups)	
8b*	3480 ~ 2300 (chelated NH) and 1670 ~ 1630 (ring CO)	
10a*	3400 ~ 2300 (chelated NH); 2220 (conjugated CN); 1750 ~ 1730 (ring and ester CO) and 1630 (C=C)	
10b*	3500 ~ 3200 (NH); 2220 (CN); 1740 (ester CO); 1680 (ring CO) and 1620 (C=C)	
10c	3500, 3200 (NH_2); 2210 (CN); 1740 (ester CO); 1670 (ring CO) and 1620 (C=C)	1.16 (t, 3 H, CH_3); 3.26 (m, 2 H, benzopyran H-3 and H-4); 4.16 (q, 2 H, CH_2) and 7.25 ~ 9.10 (m, 12 H, aromatic, NH_2 and OH protons)
11	3480, 3380, 3270 (NH_2); 2225, 2205 (CN bands); 1710, 1670 (ring CO groups); 1645 (NH_2 deformation) and 1610 (C=C)	4.0 (m, 2 H, benzopyran H-3 and H-4); 7.2-8.16 (m, 6 H, aromatic and NH_2 protons)
13	3320 ~ 3300 (chelated OH); 2240 (CN) and 1730, 1720, 1710 (CO groups)	1.16 (t, 3 H, CH_3); 3.2 (s, 1 H, CH); 4.05 (q, 2 H, CH_2); 5.4 (d, 1 H, benzopyran H-4); 6.8 ~ 7.9 (m, 4 H, aromatic and pyridine H-3) and 10.0 (s, br, 1 H, OH)
14*	3440, 3300 (NH group); 2220 (CN); 1715, 1690 (CO groups) and 1640 ~ 1610 (C=C)	

* Compound is insoluble in most known ^1H NMR solvents.

Reactions of 7 with Activated Nitriles

A solution of **7** (0.0 mol) and each of malononitrile, benzoylacetonitrile or ethyl cyanoacetate (0.01 mol) in *DMF* (20 ml) was treated with piperidine (1 ml) and the whole heated to boiling for 2 h. The solvent was evaporated in vacuo and the residue triturated with ethanol then crystallized from *DMF*/ H_2O to give compounds **11**, **13** and **14** respectively (cf. Tables 1 and 2).

References

- ¹ Abdou S., Fahmy S. M., Sadek K. U., Elnagdi M. H., *Heterocycles* **16**, 2177 (1981).
- ² Elmoghayar M. R. H., Ibraheim M. K. A., El-Ghandour A. H., Elnagdi M. H., *Synthesis* **1981**, 635.
- ³ Elnagdi M. H., Elmoghayar M. R. H., Hammam A. H., Khallaf S. A., *J. Heterocyclic Chem.* **16**, 1541 (1979).
- ⁴ Prepared after the procedure described by Taylor E. C., Hartke K. S., *J. Amer. Chem. Soc.* **81**, 2452 (1959).
- ⁵ Prepared after the procedure described by Junek H., Frosch F., *Z. Naturforsch.* **26b**, 1124 (1971).
- ⁶ Junek H., *Monatsh. Chem.* **94**, 192 (1963).
- ⁷ Elnagdi M. H., *Tetrahedron* **30**, 2791 (1974).
- ⁸ Livingstone R., Watson R. B., *J. Chem. Soc.* **1957**, 1509.
- ⁹ Shimizu T., Hoyashi Y., Yamada K., Shio T. N., Teramara K., *Bull. Chem. Soc. Jpn.* **54**, 217 (1981).