doi:10.1246/cl.2012.151
Published on the web January 28, 2012

Ru/Ag-Catalyzed Oxidative Alkenylation of Benzamides and Phenylazoles through Regioselective C–H Bond Cleavage

Yuto Hashimoto,[1] Timon Ortloff,[2] Koji Hirano,[1] Tetsuya Satoh,*[1] Carsten Bolm,[2] and Masahiro Miura*[1]
[1]*Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita, Osaka 565-0871*
[2]*Institute of Organic Chemistry, RWTH Aachen University, Landoltweg 1, D-52056 Aachen, Germany*

(Received November 17, 2011; CL-111115; E-mail: satoh@chem.eng.osaka-u.ac.jp, miura@chem.eng.osaka-u.ac.jp)

Oxidative alkenylation of *N*,*N*-disubstituted benzamides with alkenes accompanied by regioselective C–H bond cleavage proceeds efficiently in the presence of a Ru/Ag catalyst system together with a copper oxidant. The procedure is also applicable to phenylazoles.

Direct C–C coupling reactions through regioselective C–H bond cleavage have attracted much attention as atom- and step-economic tools for precise synthesis, and various catalytic processes involving different modes to activate the ubiquitously available bond have been developed.[1] Among the most promising strategies is the chelate-assisted version with the aid of directing groups. Particularly, the transition-metal-catalyzed *ortho*-alkenylation of aromatic substrates through oxidative coupling with readily available alkenes[2] is a useful method to selectively construct π-conjugated alkenylarene frameworks, which can be widely seen in organic materials.[3] As early examples, we demonstrated that 2-phenylphenols,[4] *N*-(arylsulfonyl)-2-phenylanilines,[5] and benzoic acids[5,6] undergo directed alkenylation under Pd- or Rh-catalysis. Since then, a number of related oxidative alkenylation reactions have been disclosed by us[7] and others.[8] Compared to well-developed Pd- and Rh-catalyzed reactions, the ruthenium-catalyzed version has been less explored. Recently, we succeeded in finding that heteroarene carboxylic acids[9] and phenylazoles[10] undergo directed alkenylation under Ru-catalysis. Although the Ru-catalyzed alkenylation[11,12] seems to be attractive due to relatively lower catalyst cost compared to Pd and Rh, the effective substrates have so far been limited. Consequently, we have investigated retuning our Ru-catalyst system to improve the activity and to expand its applicability and found that addition of $AgSbF_6$ as a cocatalyst together with $[Ru(p\text{-cymene})Cl_2]_2$ enables *N*,*N*-disubstituted benzamides[13] to undergo the regioselective alkenylation smoothly. Actually, the reaction did not proceed at all without the cocatalyst. The catalyst system $[Ru(p\text{-cymene})Cl_2]_2$/$AgSbF_6$ was also found to be effective for the alkenylation of phenylazoles. These new findings are described herein.

In an initial attempt, *N*,*N*-dimethylbenzamide (**1a**) (0.25 mmol) was treated with butyl acrylate (**2a**) (0.5 mmol) in the presence of $[Ru(p\text{-cymene})Cl_2]_2$ (0.013 mmol) and $Cu(OAc)_2 \cdot H_2O$ (0.5 mmol) as catalyst and oxidant, respectively, in *t*-AmOH(*tert*-pentyl alcohol) at 100 °C for 4 h under N_2. Under such conditions similar to those previously employed for the reactions of 1-phenylpyrazoles,[10a] the starting materials were recovered almost completely (Entry 1 in Table 1). Addition of $AgSbF_6$ (0.05 mmol) promoted the reaction dramatically to quantitatively afford the corresponding alkenylated product, *N*,*N*-dimethyl-2-[(*E*)-(*n*-butoxycarbonyl)ethenyl]benzamide (**3a**) (Entry 2).[14] The reactions of **1a** with ethyl (**2b**) and cyclohexyl (**2c**) acrylates proceeded similarly to give the corresponding alkenylated products **3b** and **3c** in good yields (Entries 3 and 4). Styrene (**2d**) also underwent the reaction with **1a** smoothly to give a stilbene derivative **3d** (Entry 5). Besides **1a**, *N*-benzoylpiperidine (**1b**) and -pyrrolidine (**1c**) also reacted with **2a** to afford alkenylated products, **3e** and **3f**, respectively (Entries 6 and 7). Note that *ortho*-alkenylbenzamide derivatives are of interest due to their photochemical reactivity as well as applicabilities as photosensitizers and chiral auxiliaries.[15]

Table 1. Reaction of benzamides **1** with alkenes **2**[a]



1 **2** **3**

Entry	1	2	Product, Yield/%
1[b]	**1a**	**2a**: $R^2 = CO_2(n\text{-Bu})$	**3a**: $R^2 = CO_2(n\text{-Bu})$, 0
2		**2a**: $R^2 = CO_2(n\text{-Bu})$	**3a**: $R^2 = CO_2(n\text{-Bu})$, 98
3		**2b**: $R^2 = CO_2Et$	**3b**: $R^2 = CO_2Et$, 92
4		**2c**: $R^2 = CO_2Cy$	**3c**: $R^2 = CO_2Cy$, 82
5		**2d**: $R^2 = Ph$	**3d**: $R^2 = Ph$, 62
6	**1b**	**2a**	**3e**, 82
7	**1c**	**2a**	**3f**, 75

[a]Reaction conditions: **1** (0.25 mmol), **2** (0.5 mmol), $[Ru(p\text{-cymene})Cl_2]_2$ (0.013 mmol), $AgSbF_6$ (0.05 mmol), $Cu(OAc)_2 \cdot H_2O$ (0.5 mmol), in *t*-AmOH (3 mL) at 100 °C for 4 h under N_2. [b]Without $AgSbF_6$.

In the presence of the $[Ru(p\text{-cymene})Cl_2]_2/AgSbF_6$ catalyst system, the alkenylation of phenylazoles was also found to be promoted. Previously, we reported that the reaction of 2-phenylbenzothiazole (**4**) with **2a** using $[Ru(p\text{-cymene})Cl_2]_2$ gave the corresponding alkenylated product **5** in a low yield (ca. 20%).[10a] The reaction efficiency was considerably improved by the addition of $AgSbF_6$ cocatalyst to produce **5** in 61% yield (Scheme 1).



Scheme 1.

Table 2. Reaction of phenylazoles **6** with alkenes **2**[a]



Entry	6	2	Product, Yield/%
			
1	**6a**	**2a**: R = CO_2(*n*-Bu)	**7a**: R = CO_2(*n*-Bu), 57
2[b]		**2a**: R = CO_2(*n*-Bu)	**7a**: R = CO_2(*n*-Bu), 8[c]
3		**2b**: R = CO_2Et	**7b**: R = CO_2Et, 70
4		**2c**: R = CO_2Cy	**7c**: R = CO_2Cy, 61
5		**2d**: R = Ph	**7d**: R = Ph, 71[d]
			
6	**6b**	**2a**	**7e**, 82

[a]Reaction conditions: **6** (0.25 mmol), **2** (0.5 mmol), [Ru(*p*-cymene)Cl_2]$_2$ (0.013 mmol), $AgSbF_6$ (0.05 mmol), Cu(OAc)$_2$•H_2O (0.5 mmol), in *t*-AmOH (3 mL) at 100 °C for 4 h under N_2. [b]Without $AgSbF_6$. [c]GC yield. [d]*E*/*Z* = 9:1.

With the result in hand, we reexamined the alkenylation of phenylazoles using [Ru(*p*-cymene)Cl_2]$_2$/$AgSbF_6$. As a result, 1-methyl-2-phenylimidazole (**6a**) was found to undergo the alkenylation with **2a** to produce **7a** selectively (Entry 1 in Table 2). In the absence of $AgSbF_6$, only a small amount of **7a** was formed (Entry 2). Other alkenes **2b**–**2d** also reacted with **6a** to afford the corresponding products **7b**–**7d** in 61–71% yields (Entries 3–5).[16] The reaction of 1-methyl-2-phenylbenzimidazole (**6b**) with **2a** also proceeded efficiently to give **7e** in 82% yield (Entry 6).

In summary, we have demonstrated that benzamides and phenylazoles efficiently undergo regioselective alkenylation in the presence of a Ru/Ag catalyst system and a copper oxidant.[17,18] The present catalyst system and related ones are expected to be applicable to other oxidative coupling reactions.[19] Work is underway toward further development of the catalysis.

This work was partly supported by Grants-in-Aid from MEXT and JSPS, Japan, Deutsche Forschungsgemeinschaft (IRTG 1628), and the Fonds der Chemischen Industrie, Germany. T. O. acknowledges DFG for financial support.

References and Notes

1 For selected reviews, see: a) O. Baudoin, *Chem. Soc. Rev.* **2011**, *40*, 4902. b) J. Wencel-Delord, T. Dröge, F. Liu, F. Glorius, *Chem. Soc. Rev.* **2011**, *40*, 4740. c) Y. Kuninobu, K. Takai, *Chem. Rev.* **2011**, *111*, 1938. d) L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315. e) C. S. Yeung, V. M. Dong, *Chem. Rev.* **2011**, *111*, 1215. f) T. Satoh, M. Miura, *Chem.—Eur. J.* **2010**, *16*, 11212. g) T. Satoh, M. Miura, *Synthesis* **2010**, 3395. h) D. Lapointe, K. Fagnou, *Chem. Lett.* **2010**, *39*, 1118. i) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147. j) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624. k) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Commun.* **2010**, *46*, 677. l) L. Ackermann, R. Vicente, A. R. Kapdi, *Angew. Chem., Int. Ed.* **2009**, *48*, 9792. m) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem., Int. Ed.* **2009**, *48*, 5094. n) A. V. Vasil'ev, *Russ. J. Org. Chem.* **2009**, *45*, 1. o) G. P. McGlacken, L. M. Bateman, *Chem. Soc. Rev.* **2009**, *38*, 2447. p) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013. q) E. M. Ferreira, H. Zhang, B. M. Stoltz, *Tetrahedron* **2008**, *64*, 5987. r) Y. J. Park, J.-W. Park, C.-H. Jun, *Acc. Chem. Res.* **2008**, *41*, 222. s) E. M. Beccalli, G. Broggini, M. Martinelli, S. Sottocornola, *Chem. Rev.* **2007**, *107*, 5318. t) C. I. Herrerías, X. Yao, Z. Li, C.-J. Li, *Chem. Rev.* **2007**, *107*, 2546. u) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174. v) K. Godula, D. Sames, *Science* **2006**, *312*, 67. w) F. Kakiuchi, N. Chatani, *Adv. Synth. Catal.* **2003**, *345*, 1077. x) V. Ritleng, C. Sirlin, M. Pfeffer, *Chem. Rev.* **2002**, *102*, 1731. y) F. Kakiuchi, S. Murai, *Acc. Chem. Res.* **2002**, *35*, 826. z) G. Dyker, *Angew. Chem., Int. Ed.* **1999**, *38*, 1698.

2 For nondirected Fujiwara–Moritani type reactions, see, Pd: a) Y. Fujiwara, I. Moritani, S. Danno, R. Asano, S. Teranishi, *J. Am. Chem. Soc.* **1969**, *91*, 7166. b) I. Moritani, Y. Fujiwara, *Tetrahedron Lett.* **1967**, *8*, 1119. Rh: c) T. Matsumoto, H. Yoshida, *Chem. Lett.* **2000**, 1064. d) K. Sasaki, T. Sakakura, Y. Tokunaga, K. Wada, M. Tanaka, *Chem. Lett.* **1988**, 685. Ru: e) H. Weissman, X. Song, D. Milstein, *J. Am. Chem. Soc.* **2001**, *123*, 337. For reviews, see: f) J. Le Bras, J. Muzart, *Chem. Rev.* **2011**, *111*, 1170. g) S. H. Cho, J. Y. Kim, J. Kwak, S. Chang, *Chem. Soc. Rev.* **2011**, *40*, 5068. h) B. Karimi, H. Behzadnia, D. Elhamifar, P. F. Akhavan, F. K. Esfahani, A. Zamani, *Synthesis* **2010**, 1399. i) C. Jia, T. Kitamura, Y. Fujiwara, *Acc. Chem. Res.* **2001**, *34*, 633.

3 For examples, see: a) A. C. Grimsdale, K. L. Chan, R. E. Martin, P. G. Jokisz, A. B. Holmes, *Chem. Rev.* **2009**, *109*, 897. b) A. Kraft, A. C. Grimsdale, A. B. Holmes, *Angew. Chem., Int. Ed.* **1998**, *37*, 402. c) S. R. Marder, B. Kippelen, A. K.-Y. Jen, N. Peyghambarian, *Nature* **1997**, *388*, 845.

4 M. Miura, T. Tsuda, T. Satoh, M. Nomura, *Chem. Lett.* **1997**, 1103.

5 M. Miura, T. Tsuda, T. Satoh, S. Pivsa-Art, M. Nomura, *J. Org. Chem.* **1998**, *63*, 5211.

6 a) S. Mochida, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2011**, *76*, 3024. b) S. Mochida, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2010**, *12*, 5776. c) T. Satoh, K. Ueura, M. Miura, *Pure Appl. Chem.* **2008**, *80*, 1127. d) K. Ueura, T. Satoh, M. Miura, *J. Org. Chem.* **2007**, *72*, 5362. e) K. Ueura, T. Satoh, M. Miura, *Org. Lett.* **2007**, *9*, 1407.

7 a) N. Umeda, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2009**, *74*, 7094. b) S. Mochida, K. Hirano, T. Satoh, M.

Miura, *J. Org. Chem.* **2009**, *74*, 6295. c) A. Maehara, H. Tsurugi, T. Satoh, M. Miura, *Org. Lett.* **2008**, *10*, 1159.

8 For recent examples, see: a) C. Huang, B. Chattopadhyay, V. Gevorgyan, *J. Am. Chem. Soc.* **2011**, *133*, 12406. b) T.-J. Gong, B. Xiao, Z.-J. Liu, J. Wan, J. Xu, D.-F. Luo, Y. Fu, L. Liu, *Org. Lett.* **2011**, *13*, 3235. c) M. Yu, Z. Liang, Y. Wang, Y. Zhang, *J. Org. Chem.* **2011**, *76*, 4987. d) N. Guimond, S. I. Gorelsky, K. Fagnou, *J. Am. Chem. Soc.* **2011**, *133*, 6449. e) S. H. Park, J. Y. Kim, S. Chang, *Org. Lett.* **2011**, *13*, 2372. f) K. J. Stowers, K. C. Fortner, M. S. Sanford, *J. Am. Chem. Soc.* **2011**, *133*, 6541. g) F. Wang, G. Song, Z. Du, X. Li, *J. Org. Chem.* **2011**, *76*, 2926. h) B. S. Kim, S. Y. Lee, S. W. Youn, *Chem.—Asian J.* **2011**, *6*, 1952. i) A. García-Rubia, M. Á. Fernández-Ibáñez, R. G. Arrayás, J. C. Carretero, *Chem.—Eur. J.* **2011**, *17*, 3567. j) S. Rakshit, C. Grohmann, T. Besset, F. Glorius, *J. Am. Chem. Soc.* **2011**, *133*, 2350. k) C. Zhu, J. R. Falck, *Org. Lett.* **2011**, *13*, 1214. l) A. S. Tsai, M. Brasse, R. G. Bergman, J. A. Ellman, *Org. Lett.* **2011**, *13*, 540. m) F. W. Patureau, T. Besset, F. Glorius, *Angew. Chem., Int. Ed.* **2011**, *50*, 1064. n) K. M. Engle, D.-H. Wang, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, *132*, 14137. o) K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem., Int. Ed.* **2010**, *49*, 6169. p) F. Wang, G. Song, X. Li, *Org. Lett.* **2010**, *12*, 5430. q) J. Chen, G. Song, C.-L. Pan, X. Li, *Org. Lett.* **2010**, *12*, 5426. r) B. S. Kim, C. Jang, D. J. Lee, S. W. Youn, *Chem.—Asian J.* **2010**, *5*, 2336. s) F. W. Patureau, F. Glorius, *J. Am. Chem. Soc.* **2010**, *132*, 9982. t) T. Nishikata, B. H. Lipshutz, *Org. Lett.* **2010**, *12*, 1972. u) B.-F. Shi, Y.-H. Zhang, J. K. Lam, D.-H. Wang, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, *132*, 460. v) D.-H. Wang, K. M. Engle, B.-F. Shi, J.-Q. Yu, *Science* **2010**, *327*, 315. w) W. Rauf, A. L. Thompson, J. M. Brown, *Chem. Commun.* **2009**, 3874. x) G. Cai, Y. Fu, Y. Li, X. Wan, Z. Shi, *J. Am. Chem. Soc.* **2007**, *129*, 7666. y) C. Amatore, C. Cammoun, A. Jutand, *Adv. Synth. Catal.* **2007**, *349*, 292. z) M. D. K. Boele, G. P. F. van Strijdonck, A. H. M. de Vries, P. C. J. Kamer, J. G. de Vries, P. W. N. M. van Leeuwen, *J. Am. Chem. Soc.* **2002**, *124*, 1586.

9 a) T. Ueyama, S. Mochida, T. Fukutani, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2011**, *13*, 706. After our report, a related reaction of benzoic acids was reported: b) L. Ackermann, J. Pospech, *Org. Lett.* **2011**, *13*, 4153.

10 a) Y. Hashimoto, T. Ueyama, T. Fukutani, K. Hirano, T. Satoh, M. Miura, *Chem. Lett.* **2011**, *40*, 1165. A similar report appeared independently: b) P. B. Arockiam, C. Fischmeister, C. Bruneau, P. H. Dixneuf, *Green Chem.* **2011**, *13*, 3075.

11 For Ru-catalyzed directed alkenylation using vinyl halides, vinyl boronates, and vinyl and alkyl acetates, see: a) S. Ueno, T. Kochi, N. Chatani, F. Kakiuchi, *Org. Lett.* **2009**, *11*, 855. b) Y. Matsuura, M. Tamura, T. Kochi, M. Sato, N. Chatani, F. Kakiuchi, *J. Am. Chem. Soc.* **2007**, *129*, 9858. c) S. Ueno, N. Chatani, F. Kakiuchi, *J. Org. Chem.* **2007**, *72*, 3600. d) S. Oi, Y. Tanaka, Y. Inoue, *Organometallics* **2006**, *25*, 4773. e) S. Oi, E. Aizawa, Y. Ogino, Y. Inoue, *J. Org. Chem.* **2005**, *70*, 3113. f) S. Oi, Y. Ogino, S. Fukita, Y. Inoue, *Org. Lett.* **2002**, *4*, 1783. g) S. Oi, S. Fukita, N. Hirata, N. Watanuki, S. Miyano, Y. Inoue, *Org. Lett.* **2001**, *3*, 2579.

12 Ru-Catalyzed coupling of arylamides with alkenes without any oxidants has been reported. In the reactions, mixtures of oxidative- and nonoxidative coupling products were formed: K.-H. Kwon, D. W. Lee, C. S. Yi, *Organometallics* **2010**, *29*, 5748.

13 For the Rh-catalyzed alkenylation of these amides, see reference 8m.

14 Use of $AgSbF_6$ under Rh catalysis, see for example: a) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess, K. Fagnou, *J. Am. Chem. Soc.* **2008**, *130*, 16474. Similar Ru/Ag catalyst systems were employed in the nonoxidative cyclization of arene–ene substrates: b) S. W. Youn, S. J. Pastine, D. Sames, *Org. Lett.* **2004**, *6*, 581.

15 a) C. B. Nielsen, J. Arnbjerg, M. Johnsen, M. Jørgensen, P. R. Ogilby, *J. Org. Chem.* **2009**, *74*, 9094. b) R. Rios, C. Jimeno, P. J. Carroll, P. J. Walsh, *J. Am. Chem. Soc.* **2002**, *124*, 10272. c) F. D. Lewis, S. G. Kultgen, *J. Photochem. Photobiol., A* **1998**, *112*, 159.

16 In the case using **2d**, a minor amount of (*Z*)-**7d** was also formed. However, the reason why the *Z*-isomer formation was observed only in this case is obscure.

17 Through the present study, dialkenylation products could not be detected.

18 Supporting Information is available electronically on the CSJ-Journal Web site, http://www.csj.jp/journals/chem-lett/index.html.

19 A report of *ortho*-alkenylation of aromatic ketones using a Ru/Ag catalyst appeared during the preparation of this manuscript. K. Padala, M. Jeganmohan, *Org. Lett.* **2011**, *13*, 6144.

