## A one-pot synthesis of (*E*,*E*)-1-arylseleno-substituted 1,5-dien-3-ynes by palladium-catalysed three component cross-coupling reaction Fang Yao<sup>a,b</sup>, Wenyan Hao<sup>a</sup> and Mingzhong Cai<sup>a</sup>\*

<sup>a</sup>Department of Chemistry, Jiangxi Normal University, Nanchang 330022, P. R. China <sup>b</sup>Department of Chemistry and Pharmaceutical Engineering, West Branch of Zhejiang University of Technology, Quzhou 324000, P. R. China

(E,E)-1-Arylseleno-substituted 1,5-dien-3-ynes can be stereoselectively synthesised in one pot, in good yields with high stereoselectivity, by the Pd(PPh<sub>3</sub>)<sub>4</sub>-catalysed Stille coupling of (E)-1-iodo-2-arylselenoethenes with trimethyl-stannyl(trimethylsilyl)ethyne, followed by the cross-coupling with (E)-vinyl iodides in the presence of tris(diethylamino) sulfonium trimethyldifluorosilicate (TASF).

**Keywords:** (*E*)-1-iodo-2-arylselenoethene, Stille coupling, vinyl iodide, 1,5-dien-3-yne, tandem reaction, stereoselective synthesis

Enyne systems have attracted much attention from synthetic organic chemists as they show interesting chemical and biological reactivities.<sup>1-3</sup> Stereodefined conjugated polyenynes containing an internal carbon-carbon triple bond unit are widely distributed in nature and show interesting biological activities.4,5 Recently, dienyne compounds have attracted much interest since they are important building blocks for the formation of natural products in organic synthesis.<sup>6-10</sup> Uenishi and Matsui reported the stereocontrolled synthesis of dienynes by stepwise Suzuki/Sonogashira coupling reactions of 1,1dibromoalk-1-enes.11 Sato and coworkers reported that conjugated divnes underwent selective mono-titanation with a Ti(II) reagent to give 1:1 diyne-titanium alkoxide complexes, which reacted with another acetylene to give stereo-defined dienynes.12 van Otterlo et al. described the synthesis of dienynes from alkenes and diynes using ruthenium-mediated ring-closing metathesis.13 Shi and coworkers14 reported that 2-iodo-4-(phenylchalcogenyl)but-1-enes can be enynylated with alkynes catalysed by Pd(OAc)<sub>2</sub> to give conjugated dienynes. The formation of dienynes containing metal or heteroatom functional groups has also attracted considerable interest in organic synthesis because many useful functional group transformations can be achieved by introduction and removal of metal or heteroatom functions. Alami and Ferri reported the regioselective synthesis of stannylated dienynes by the palladium-catalysed hydrostannylation of enediynes.<sup>15</sup> However, the synthesis of heteroatom-substituted 1,5-dien-3-ynes has received less attention. The tandem reaction has recently been of interest for organic synthesis because it offers a convenient and economical method with which to prepare target organic molecules.<sup>16-19</sup> We now report that (E,E)-1-arylseleno-substituted 1,5-dien-3ynes can be conveniently synthesised in one pot, in good yields with high stereoselectivity, by the Stille coupling of (E)-1iodo-2-arylselenoethenes 1 with trimethylstannyl(trimethylsil yl)ethyne, followed by the cross-coupling with (E)-vinyl

iodides **3** in the presence of tris(diethylamino)sulfonium trimethyldifluorosilicate (TASF).

(E)-1-Iodo-2-arylselenoethenes 1 were easily prepared by the hydrozirconation of arylselenoacetylenes, followed by the reaction with iodine according to a literature procedure.<sup>20</sup> The Stille coupling reaction of (E)-1-iodo-2-arylselenoethenes 1 with trimethylstannyl(trimethylsilyl)ethyne in the presence of  $Pd(PPh_3)_4$  in THF gave stereoselectively (E)-1-arylseleno-4-(trimethylsilyl)but-1-en-3-ynes 2 in high yields (Scheme 1). The (E)-configuration of compounds **2a-c** was confirmed by their <sup>1</sup>H NMR spectra, which showed two doublets in the  $\delta = 7.21 - 5.64$  with a coupling constant of 15.6-16.0 Hz. Hatanaka and Hiyama reported that in the presence of a palladium catalyst and tris(diethylamino)sulfonium trimethyldifluorosilicate (TASF), alkynylsilanes can react with alkenyl halides to give the corresponding coupled products in a stereospecific and chemoselective manner.<sup>21</sup> Considering the fact that both the Stille coupling and the cross-coupling reactions of alkynylsilanes with alkenyl halides were catalysed by palladium complexes, we tried to combine the two reactions, in one pot, to synthesise stereoselectively (E,E)-1arylseleno-substituted 1,5-dien-3-ynes. The basic concept of our reaction is illustrated in Scheme 2.

We found that, if after the Stille coupling of (*E*)-1-iodo-2-arylselenoethenes **1** with trimethylstannyl(trimethylsilyl) ethyne (1.2 equiv) using 5 mol% Pd(PPh<sub>3</sub>)<sub>4</sub> in THF at 50 °C for 3 h, (*E*)-vinyl iodides and TASF (2.4 equiv) were added at -78 °C and the mixture was slowly warmed to ambient temperature and then allowed to react at 50 °C for 2 h, (*E*,*E*)-1-arylseleno-substituted 1,5-dien-3-ynes **4** were obtained in good yields. The experimental results are summarised in Table 1. It was found that the kind of the palladium catalyst is critical for the success of the tandem reaction and Pd(PPh<sub>3</sub>)<sub>4</sub> was the most efficient. PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>- and PdCl<sub>2</sub>(MeCN)<sub>2</sub>-catalysed



Scheme 1

<sup>\*</sup> Correspondent. E-mail: caimzhong@163.com



tandem reactions gave a relatively large amount of protodesilylation products of the intermediates **2**. The use of  $(\eta^3-C_3H_5PdCl)_2$  as the catalyst, which was effective for the TASFmediated reaction of alkynyltrimethylsilanes with alkenyl halides, resulted in extensive homo-coupling of trimethylstannyl(trimethylsilyl)ethynetoproduce 1,4-bis(trimethylsilyl)buta-1,3-diyne.<sup>21</sup> As shown in Table 1, the tandem Stille/crosscoupling reaction of trimethylstannyl(trimethylsilyl)ethyne with a variety of (*E*)-1-iodo-2-arylselenoethenes and (*E*)vinyl iodides proceeded smoothly under mild conditions to afford stereoselectively the corresponding (*E*,*E*)-1-arylselenosubstituted 1,5-dien-3-ynes **4**.

The stereochemistry of compounds **4** was easily established since their <sup>1</sup>H NMR spectra give rise to three or four doublets at  $\delta = 7.15-5.55$  with a coupling constant of 15.6–16.0 Hz. The results summarised in Table 1 show the following salient features of the novel process. The palladium(0) catalyst [Pd(PPh<sub>3</sub>)<sub>4</sub>] can be applied in a single flask to double-coupling reactions based on the reactivity difference between Me<sub>3</sub>Sn and Me<sub>3</sub>Si groups. The tandem reaction proceeds with retention of the double bond geometry of the alkenyl iodides as is often the case with the coupling reaction of organotin and organosilicon compounds.

In summary, we have developed an efficient and stereoselective one-pot method for the synthesis of (E,E)-1-arylselenosubstituted 1,5-dien-3-ynes 4 by the Pd(PPh<sub>3</sub>)<sub>4</sub>-catalysed Stille coupling of (E)-1-iodo-2-arylselenoethenes with trimethylstannyl(trimethylsilyl)ethyne, followed by the cross-coupling with (E)-vinyl iodides in the presence of tris(diethylamino)sulfonium trimethyldifluorosilicate (TASF). The present method has the advantages of readily available starting materials, straightforward and simple procedures, mild reaction conditions, high stereoselectivity and good yields. The procedure should find wide application to the synthesis of a large array of naturally occurring substances having the conjugated dienyne system.

## Experimental

<sup>1</sup>H NMR spectra were recorded on a Bruker AC-P400 (400 MHz) spectrometer with TMS as an internal standard using CDCl<sub>3</sub> as the solvent. <sup>13</sup>C NMR (100 MHz) spectra were recorded on a Bruker AC-P400 (400 MHz) spectrometer using CDCl<sub>3</sub> as the solvent. IR spectra

 
 Table 1
 Synthesis of (*E,E*)-1-arylseleno-substituted 1,5-dien-3-ynes 4<sup>a</sup>

Entry	Ar	R	Product	Yield /% <sup>b</sup>
	5	<u></u>		
1	Ph	<i>n</i> -C₄H <sub>9</sub>	4a	79
2	Ph	CH <sub>3</sub> OCH <sub>2</sub>	4b	82
3	Ph	Ph	4c	85
4	4-CIC <sub>6</sub> H <sub>4</sub>	<i>n</i> -C₄H <sub>9</sub>	4d	80
5	4-CIC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub> OCH <sub>2</sub>	4e	84
6	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	$n - C_6 H_{13}$	4f	80
7	$4-CH_3C_6H_4$	CH <sub>3</sub> OCH <sub>2</sub>	4g	78

<sup>a</sup> The reactions were performed with 1.2 mmol of trimethylstan nyl(trimethylsilyl)ethyne, 1.0 mmol of 1, 0.05 mmol of Pd(PPh<sub>3</sub>)<sub>4</sub>, 1.0 mmol of 3 and 2.4 mmol of TASF in THF (13.5 mL) under Ar.

<sup>b</sup> Isolated yield based on the **1** used.

were determined on an FTS-185 instrument as neat films. Mass spectra were obtained on a Finigan 8239 mass spectrometer. Microanalyses were measured using a Yanaco MT-3 CHN microelemental analyser. All reactions were carried out in pre-dried glassware (150 °C, 4 h and cooled under a stream of dry Ar).

## Synthesis of (E)-1-arylseleno-4-(trimethylsilyl)but-1-en-3-ynes **2a–c;** general procedure

Trimethylstannyl(trimethylsilyl)ethyne (1.2 mmol) was added to a solution of (*E*)-1-iodo-2-arylselenoethene **1** (1.0 mmol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (0.05 mmol) in THF (10 mL) under an argon atmosphere at room temperature. The resulting mixture was then stirred at 50 °C and the reaction was monitored by TLC (SiO<sub>2</sub>) until the compound **1** was consumed (3 h). The reaction mixture was cooled to room temperature, diluted with diethyl ether (50 mL), filtered and then treated with 20% aqueous KF (10 mL) for 30 min before being dried and concentrated. The residue was purified by column chromatography on silica gel, eluting with light petroleum (b.p. 30–60 °C).

(*E*)-*1*-*Phenylseleno-4-trimethylsilylbut-1-en-3-yne* (**2a**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3059, 2958, 2142, 1712, 1578, 1250, 844, 760, 690; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.54–7.51 (m, 2H), 7.34–7.32 (m, 3H), 7.21 (d, *J* = 16.0 Hz, 1H), 5.71 (d, *J* = 16.0 Hz, 1H), 0.16 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  136.4, 134.1, 129.6, 128.3, 127.8, 111.7, 103.3, 95.2, -0.1; MS: *m*/z 279 (M<sup>+</sup>, 12), 250 (55), 78 (44), 73 (100). Anal. Calcd for C<sub>13</sub>H<sub>16</sub>SiSe: C, 55.89; H, 5.77. Found: C, 55.68; H, 5.56%.

(*E*)-1-(4-Methylphenyl)seleno-4-trimethylsilylbut-1-en-3-yne (**2b**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3033, 2961, 2163, 1704, 1563, 1251, 844, 803, 760; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.43–7.41 (m, 2H), 7.19 (d, J = 15.6 Hz, 1H), 7.14 (d, J = 8.0 Hz, 2H), 5.64 (d, J = 15.6 Hz, 1H), 2.35 (s, 3H), 0.16 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  138.8, 137.1, 134.7, 130.5, 123.9, 111.1, 103.6, 95.1, 21.3, 0.1; MS: *m*/z 294 (M<sup>+</sup>, 14), 263 (33), 91 (35), 73 (100). Anal. Calcd for C<sub>14</sub>H<sub>18</sub>SiSe: C, 57.31; H, 6.18%. Found: C, 57.58; H, 6.29%. (*E*)-1-(4-Chlorophenyl)seleno-4-trimethylsilylbut-1-en-3-yne

(*E*)-*1*-(*4*-*Methylphenyl*)*seleno*-*4*-*trimethylsilylbut*-*1*-*en*-*3*-*yne* (**2c**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3031, 2957, 2166, 1706, 1556, 1473, 1254, 842, 824, 760; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.46–7.43 (m, 2H), 7.31–7.28 (m, 2H), 7.15 (d, *J* = 15.6 Hz, 1H), 5.73 (d, *J* = 15.6 Hz, 1H), 0.17 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  135.5, 135.4, 134.9, 129.9, 126.2, 112.6, 103.1, 95.8, 0.0; MS: *m*/z 314 (M<sup>+</sup>, <sup>35</sup>Cl, 32), 191 (100), 189 (56), 156 (50), 73 (33). Anal. Calcd for C<sub>13</sub>H<sub>15</sub>SiSeCl: C, 49.75; H, 4.82. Found: C, 49.48; H, 4.56%.

Synthesis of (E,E)-1-arylseleno-substituted 1,5-dien-3-ynes **4a-g**; general procedure

Trimethylstannyl(trimethylsilyl)ethyne (1.2 mmol) was added to a solution of (*E*)-1-iodo-2-arylselenoethene **1** (1.0 mmol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (0.05 mmol) in THF (10 mL) under an argon atmosphere at room temperature. The resulting mixture was then stirred at 50 °C and the reaction was monitored by TLC (SiO<sub>2</sub>) until the compound **1** was consumed (3 h). (*E*)-vinyl iodide **3** (1.0 mmol) and a solution of TASF (0.69 M, 2.4 mmol) in THF (3.5 mL) were added at -78 °C. The mixture was slowly warmed to room temperature, allowed to react at 50 °C for 2 h, then cooled to room temperature, and finally was quenched with aq. sodium bicarbonate. The mixture was extracted with diethyl ether (2 × 30 mL), the ether solution was treated with 20% aqueous KF (10 mL) for 30 min before being dried and concentrated. The residue was purified by column chromatography on silica gel, eluting with light petroleum (b.p. 30–60 °C).

(*1E,5E*)-*1-Phenylselenodeca-1,5-dien-3-yne* (**4a**): Colourless oil. IR (film): ν (cm<sup>-1</sup>) 3018, 2957, 2927, 2174, 1578, 1477, 1438, 926, 735, 690; <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.533–7.51 (m, 2H), 7.34–7.26 (m, 3H), 7.08 (d, J = 16.0 Hz, 1H), 6.13 (dt, J = 16.0, 7.2 Hz, 1H), 5.89 (d, J = 16.0 Hz, 1H), 5.56 (d, J = 16.0 Hz, 1H), 2.12–2.08 (m, 2H), 1.39–1.30 (m, 4H), 0.89 (t, J = 7.2 Hz, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  145.1, 133.5, 133.2, 129.5, 129.3, 128.0, 113.0, 109.4, 89.4, 86.4, 32.9, 30.8, 22.1, 13.9; MS: m/z 290 (M<sup>+</sup>, 43), 233 (32), 165 (100), 128 (53), 78 (84), 77 (66). Anal. Calcd for C<sub>16</sub>H<sub>18</sub>Se: C, 66.42; H, 6.27. Found: C, 66.19; H, 6.08%.

(*1E*,*5E*)-*1*-*Phenylselenodeca-1*,*5*-*dien-3*-*yne* (**4b**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3054, 2924, 2175, 1578, 1554, 1477, 1438, 1120, 928, 738, 690; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.54–7.52 (m, 2H), 7.35–7.32 (m, 3H), 7.14 (d, *J* = 15.6 Hz, 1H), 6.13 (dt, *J* = 15.6, 5.6 Hz, 1H), 5.86 (d, *J* = 15.6 Hz, 1H), 5.82 (d, *J* = 15.6 Hz, 1H), 3.96 (d, *J* = 5.6 Hz, 2H), 3.34 (s, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  139.1, 134.6, 133.7, 129.6, 129.5, 128.2, 112.1, 111.6, 88.5, 88.2, 72.3, 58.2; MS: *m/z* 278 (M<sup>+</sup>, 57), 165 (100), 152 (51), 121 (45), 78 (87), 77 (99), 63 (65). Anal. Calcd for C<sub>14</sub>H<sub>14</sub>OSe: C, 60.65; H, 5.09. Found: C, 60.40; H, 4.88%.

(*1E*,5*E*)-*1*-Phenylseleno-6-phenylhexa-1,5-dien-3-yne(**4c**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3058, 2957, 2175, 1597, 1548, 1492, 928, 867, 734, 691; 'H NMR (CDCl<sub>3</sub>):  $\delta$  7.55-7.52 (m, 2H), 7.42–7.37 (m, 2H), 7.29–7.25 (m, 6H), 7.12 (d, *J* = 16.0 Hz, 1H), 7.05 (d, *J* = 16.0 Hz, 1H), 6.09 (d, *J* = 16.0 Hz, 1H), 5.86 (d, *J* = 16.0 Hz, 1H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  145.4, 136.5, 133.7, 132.5, 130.3, 129.6, 129.4, 128.2, 126.7, 123.1, 113.3, 109.7, 90.2, 87.1; MS: *m/z* 310 (M<sup>+</sup>, 100), 165 (84), 153 (45), 123 (38), 77 (71). Anal. Calcd for C<sub>18</sub>H<sub>14</sub>Se: C, 69.90; H, 4.56. Found: C, 69.62; H, 4.70%.

(*1E*,5*E*)-*1*-(*4*-Chlorophenyl)selenodeca-*1*,5-dien-3-yne (**4d**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3020, 2957, 2928, 2183, 1558, 1474, 1387, 1090, 926, 814, 730; 'H NMR (CDCl<sub>3</sub>):  $\delta$  7.47–7.43 (m, 2H), 7.31–7.26 (m, 2H), 7.02 (d, *J* = 16.0 Hz, 1H), 6.14 (dt, *J* = 15.6, 7.2 Hz, 1H), 5.89 (d, *J* = 15.6 Hz, 1H), 5.56 (d, *J* = 16.0 Hz, 1H), 2.14–2.09 (m, 2H), 1.41–1.25 (m, 4H), 0.89 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  145.5, 144.5, 134.7, 132.3, 129.7, 126.8, 113.8, 109.3, 89.8, 86.1, 32.9, 30.8, 22.1, 13.9; MS: *m*/z 324 (M<sup>+</sup>, <sup>35</sup>Cl, 84), 232 (39), 165 (100), 152 (35). Anal. Calcd for C<sub>16</sub>H<sub>17</sub>SeCl: C, 59.35; H, 5.29. Found: C, 59.52; H, 5.48%.

(*1E*,5*E*)-*1*-(*4*-Chlorophenyl)seleno-7-methoxyhepta-1,5-dien-3-yne (**4e**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3036, 2958, 2174, 1582, 1548, 1479, 1121, 927, 867, 815, 732; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.42–7.39 (m, 2H), 7.31–7.29 (m, 2H), 7.15 (d, *J* = 15.6 Hz, 1H), 6.14 (dt, *J* = 15.6, 5.6 Hz, 1H), 5.88 (d, *J* = 15.6 Hz, 1H), 5.83 (d, *J* = 15.6 Hz, 1H), 5.98 (d, *J* = 15.6 Hz, 1H), 5.83 (d, *J* = 15.6 Hz, 1H), 3.97 (d, *J* = 5.6 Hz, 2H), 3.35 (s, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  1462, 134.8, 133.9, 129.8, 129.4, 128.5, 112.3, 111.8, 88.7, 88.4, 72.5, 58.3; MS: m/z 312 (M<sup>+</sup>, <sup>35</sup>Cl, 42), 165 (100), 45 (74). Anal. Calcd for C<sub>14</sub>H<sub>13</sub>OSeCl: C, 53.94; H, 4.20. Found: C, 53.67; H, 4.39%.

(1E, 5E)-1-(4-Methylphenyl)seleno-1,5-dodecadien-3-yne (**4f**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 3020, 2956, 2926, 2182, 1645, 1559, 1490, 926, 803; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.41 (d, J = 8.0 Hz, 2H), 7.13 (d, J = 8.0 Hz, 2H), 7.06 (d, J = 16.0 Hz, 1H), 6.11 (dt, J = 16.0, 7.2 Hz, 1H), 5.80 (d, J = 16.0 Hz, 1H), 5.55 (d, J = 16.0 Hz, 1H), 2.34 (s, 3H), 2.14–2.05 (m, 2H), 1.39–1.25 (m, 8H), 0.87 (t, J = 6.8 Hz, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  145.0, 138.3, 134.0, 133.9, 130.3, 124.5, 112.1,

109.5, 89.2, 86.5, 33.2, 31.7, 28.8, 28.7, 22.6, 21.2, 14.1; MS: m/z 332 (M<sup>+</sup>, 81), 247 (59), 181 (64), 165 (100), 91 (91). Anal. Calcd for C<sub>19</sub>H<sub>24</sub>Se: C, 68.86; H, 7.30. Found: C, 68.62; H, 7.09%.

(*1E*,5*E*)-*1*-(4-*Methylphenyl*)*seleno*-7-*methoxyhepta*-1,5-*dien*-3-*yne* (**4g**): Colourless oil. IR (film): v (cm<sup>-1</sup>) 2958, 2180, 1557, 1497, 1119, 927, 867, 805; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.44–7.42 (m, 2H), 7.16–7.14 (m, 2H), 7.11 (d, *J* = 15.6 Hz, 1H), 6.10 (dt, *J* = 15.6, 5.6 Hz, 1H), 5.84 (d, *J* = 15.6 Hz, 1H), 5.80 (d, *J* = 15.6 Hz, 1H), 3.95 (d, *J* = 5.6 Hz, 2H), 3.34 (s, 3H), 2.36 (s, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  139.3, 134.5, 133.9, 129.5, 129.2, 128.5, 112.3, 111.5, 88.6, 88.2, 72.5, 58.3, 21.5; MS: *m*/z 292 (M<sup>+</sup>, 35), 165 (100), 91 (78). Anal. Calcd for C<sub>15</sub>H<sub>16</sub>OSe: C, 61.85; H, 5.54. Found: C, 61.57; H, 5.38%.

We thank the National Natural Science Foundation of China (Project No. 20862008) and the Natural Science Foundation of Jiangxi Province of China (2008GQH0034) for financial support.

Received 11 September 2010; accepted 18 December 2010 Paper 1000348 doi: 10.3184/030823410X12878393498458 Published online: 24 November 2010

## References

- R. Kouno, T. Okauchi, M. Nakamura, J. Ichikawa and T. Minami, J. Org. Chem., 1998, 63, 6239.
- 2 S. Ikeda and Y. Sato, J. Am. Chem. Soc., 1994, 116, 5975.
- 3 H.A. Stefani, R. Cella, F.A. Dorr, C.M.P. Pereira, G. Zeni and M. Gomes, *Tetrahedron Lett.*, 2005, 46, 563.
- 4 E.R.H. Jones and V. Thaller, *The chemistry of carbon-carbon triple bond*, Part II; S. Patai, Ed.; Interscience; New York, 1978, p 621.
- 5 F. Bohlmann, T. Burkhardt and C. Zdero, *Naturally occurring acetylenes*, Academic, New York, 1973.
- 6 F.D. Boyer, I. Hanna and L. Ricard, Org. Lett., 2001, 3, 3095.
- 7 E.M. Codesido, L. Castedo and J.R. Granja, Org. Lett., 2001, 3, 1483.
- 8 B.P. Peppers and S.T. Diver, J. Am. Chem. Soc., 2004, 126, 9524.
- 9 R. Garcia-Fandino, E.M. Codesido, E. Sobarzo-Sanchez, L. Castedo and J.R. Granja, Org. Lett., 2004, 6, 193.
- 10 F.D. Boyer and I. Hanna, Org. Lett., 2007, 9, 2293.
- 11 J. Uenishi and K. Matsui, Tetrahedron Lett., 2001, 42, 4353.
- 12 C. Delas, H, Urabe and F. Sato, Chem. Commun., 2002, 272.
- 13 W.A.L. van Otterlo, E.L. Ngidi, C.B. de Koning and M.A. Fernandes, *Tetrahedron Lett.*, 2004, 45, 659.
- 14 M. Shi, L.-P. Liu and J. Tang, Org. Lett., 2005, 7, 3085.
- 15 F. Frri and M. Alami, Tetrahedron Lett., 1996, 37, 7971.
- 16 G.H. Posner, Chem. Rev., 1986, 86, 831.
- 17 L.F. Tietze and U. Beifuss, Angew. Chem. Int. Ed., 1993, 32, 131.
- 18 L.F. Tietze, Chem. Rev., 1996, 96, 115.
- 19 X. Huang and M. Xia, Org. Lett., 2002, 4, 1331.
- 20 L.-S. Zhu, Z.-Z. Huang and X. Huang, J. Chem. Res., 1996, 112.
- 21 Y. Hatanaka and T. Hiyama, J. Org. Chem., 1988, 53, 918.

Copyright of Journal of Chemical Research is the property of Science Reviews 2000 Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.