

**Figure 1.** Cyclic voltammograms (10 mV/s) of (a) Pt/Nafion,TTF<sup>+</sup> electrode in 1 M KBr; (b) Pt/TFF (~800-Å thickness) electrode in 1 M KBr; (c) Pt/Nafion,Cp<sub>2</sub>FeTMA<sup>+</sup> in 0.2 M Na<sub>2</sub>SO<sub>4</sub>; (d) Pt/Nafion,TTF<sup>+</sup> electrode in 0.08 M FeY<sup>2-</sup>, 1 M KBr; (e) Pt electrode in same solution as (d); (f) Pt/Nafion,Cp<sub>2</sub>FeTMA<sup>+</sup> electrode in 0.1 M FeY<sup>2-</sup>, 0.2 M Na<sub>2</sub>SO<sub>4</sub>. All electrode potentials are given vs. SCE. Current and potentials scales given for (a) also supply to (b), (d), and (e). Potentials are negative to the right and cathodic currents are drawn upward.

covered with Nafion shows currents for FeY<sup>2-</sup> and Fe(CN)<sub>6</sub><sup>3-</sup> at least 50 times smaller than that of a bare Pt electrode of the same area, so that transport of species through the film itself or via pin holes<sup>2a</sup> makes only a small contribution.

At this time the effect observed can best be explained as the formation of an internal TTF structure in the polymer film analogous to that found in TTF<sup>+</sup> salts,<sup>5,7</sup> especially after the polymer electrode has been taken through several cycles. Indeed after repetitive electrochemical cycling, observation of the electrode surface by scanning electron microscopy shows the formation of small (<1-μm diameter) needles of crystalline TTF Br on the surface of the polymer exposed to the solution, with some of these embedded into the Nafion surface. The behavior and formation of these crystals is currently under investigation. However, the enhanced conductive properties and color changes in the Nafion,TTF<sup>+</sup> electrodes are found even after a few cycles before large domains of crystalline TTF Br are apparent. One can also contrast the Nafion,TTF<sup>+</sup> electrodes with TTF films prepared by evaporation in terms of stability. The TTF Br film electrodes<sup>7</sup> are stable under cycling for only ~30 min, while the polymer electrodes can be cycled without change for at least several hours.

Moreover, incorporation of TTF<sup>+</sup> decreases the "dry" resistance of the Nafion film. The Nafion,TTF<sup>+</sup> layer on Pt was cycled in 1 M KBr, removed from solution, blotted dry, and allowed to dry in air for 10 min. A drop of Hg was placed on the polymer layer as an electrical contact. The *i*-*V* curve across the TTF-containing polymer between the Pt and Hg contacts was typically ohmic and independent of *v* and showed a resistance of ~3 kΩ. A Nafion,TTF electrode which had not been cycled showed a resistance of ~4 MΩ. A "dry" Pt/Nafion electrode which does not contain TTF and was subjected to cycling in 1 M KBr shows purely capacitive behavior (*i* proportional to *v* and essentially independent of *V*). While the actual resistivity of the film cannot be computed from these measurements because of uncertainties in the nature of the Hg contact, the results clearly show the dramatic effect of incorporation of TTF<sup>+</sup> into the film.

These Nafion,TTF<sup>+</sup> films may find several applications, e.g., as electrochromic devices and in stabilization of semiconductor electrodes. Experiments with covalently attached polymer layers<sup>9</sup>

or electrodeposited polypyrrole<sup>4b</sup> have shown that irradiated semiconductor electrodes (e.g., n-Si) in photoelectrochemical cells can often be stabilized by such layers. A critical factor in the design of such layers is the rate at which photogenerated charge in the semiconductor can be transported through them to solution species. The Nafion,TTF<sup>+</sup> layers look especially attractive for such an application and preliminary experiments have shown similar enhanced charge transport with these layers on semiconductors.<sup>10</sup>

(10) The support of this research by the National Science Foundation (CHE 8000682) and the Solar Energy Research Institute is gratefully acknowledged.

## Kijanimicin. 1. Structures of the Individual Sugar Components

Alan K. Mallams,\* Mohindar S. Puar, and  
Randall R. Rossman

Research Division, Schering-Plough Corporation  
Bloomfield, New Jersey 07003  
Received January 19, 1981

Kijanimicin,<sup>1</sup> the major component of a complex of antibiotics produced by *Actinomadura kijaniata* nov. sp.,<sup>2</sup> contains five sugars. Acidic hydrolysis of kijanimicin using 0.5 N methanolic hydrogen chloride at 25 °C for 16.5 h afforded methyl 2,6-di-deoxy-4-*O*-methyl-β-L-ribo-hexopyranoside (1)<sup>3</sup> (Chart), mp 76.0 °C, [α]<sub>D</sub> -12.4° (CH<sub>3</sub>OH) and the α anomer (2) as a gum [α]<sub>D</sub> -209.2°. The <sup>1</sup>H NMR (Table I), <sup>13</sup>C NMR (Table II) and mass

(1) Isolated from a culture of a soil sample collected in Kenya and named after the Swahili word "kijani" for green, which is the color of the fermentation broth. Kijanimicin is also referred to as SCH 25663.

(2) J. A. Waitz, A. C. Horan, M. Kalyanpur, B. K. Lee, D. Loebenberg, J. A. Marquez, G. Miller, and M. G. Patel, *J. Antibiotics*, in press.

(3) Unless otherwise stated in this and the following paper, all specific rotations were recorded in chloroform at 26 °C (c 0.3); IR and NMR spectra were recorded in CDCl<sub>3</sub> (reference Me<sub>4</sub>Si). All compounds gave either satisfactory microanalyses or high-resolution mass spectral data.

(9) See, e.g.: Fischer, A. B.; Wrighton, M. S.; Umana, M.; Murray, R. W. *J. Am. Chem. Soc.* 1979, 101, 3442 and references therein.

Table I.  $^1\text{H}$  NMR Data

compd	1	2	3	8	9
H <sub>1</sub>	4.73 dd	4.79 dd	4.78 dd	4.59 dd	4.47 dd
H <sub>2eq</sub>	2.19 ddd	2.21 ddd	2.24 ddd	2.71 dd	2.69 ddd
H <sub>2ax</sub>	1.62 ddd	1.87 ddd	1.81 ddd	1.78 dd	1.54 dd
H <sub>3eq</sub>	4.28 ddd	4.24 ddd	5.76 ddd		
H <sub>4</sub>	2.88 dd	2.87 dd	3.00 dd	4.40 dd	4.44 ddd
H <sub>5ax</sub>	3.77 dq	3.99 dq	3.94 dq	4.22 dq	3.60 dq
6-CH <sub>3</sub>	1.30 d	1.33 d	1.36 d	1.19 d	1.23 d
1-OCH <sub>3</sub>	3.48 s	3.47 s <sup>a</sup>	3.49 s	3.19 s	3.51 s
3-CH <sub>3</sub>				1.52 s	1.59 s
3-OH	2.34 s	1.61 bs			
3-OBz			7.40 m		
4-OCH <sub>3</sub>	3.42 s	3.40 s <sup>a</sup>	3.37 s		
4-NHCOOCH <sub>3</sub>				5.00 d	5.14 d
4-NHCOOCH <sub>3</sub>				3.70 s	3.75 s
J <sub>1,2eq</sub> <sup>b</sup>	2.5	1.5	2.5	1	2.5
J <sub>1,2ax</sub>	9.5	3.5	9	4	10
J <sub>2eq,2ax</sub>	14	14	14	15	15
J <sub>2eq,3eq</sub>	3.5	3	3.5		
J <sub>2ax,3eq</sub>	3	3.5	3		
J <sub>2eq,4eq</sub>					1
J <sub>3eq,4ax</sub>	3	3	3		
J <sub>4,5ax</sub>	9.5	10	9	1	1
J <sub>4eq,4-NH</sub>				10	10
J <sub>5ax,6</sub>	6	6	6.5	6	6

<sup>a</sup> May be interchanged. <sup>b</sup> J is in Hz.Table II.  $^{13}\text{C}$  NMR Data


carbon	1	2	3	4	5	6	7	8	9	10	11
C <sub>1</sub>	99.0	98.4	99.2	98.4	99.0 (d)	105.3	105.5	96.9 (d)	99.9	99.1	100.4
C <sub>2</sub>	36.7	35.2	35.9	35.2	37.7 (t)	42.0	42.7	34.6 (t)	35.8	37.4	40.2
C <sub>3</sub>	64.0	63.5	66.4	67.4	68.0 (d)	67.7	68.7	86.0 (s)	90.7	59.3	57.8
C <sub>4</sub>	82.8	82.4	81.4	72.6	73.2 (d)	90.9	91.1	52.8 (d)	53.9	50.7	53.0
C <sub>5</sub>	68.2	62.4	69.7	64.4	69.6 (d)	71.2	71.1	62.3 (d)	68.9	61.4	68.2
C <sub>6</sub>	18.2	18.0	18.3	17.8	18.1 (q)	18.6	18.9	16.9 (q)	16.9	17.4	17.2
1-OCH <sub>3</sub>	56.5	55.1	56.4	55.1	56.5 (q)	54.8	55.4	55.0 (q)	56.6	55.2	56.4
3-CH <sub>3</sub>								26.4 (q)	25.4	23.1	30.7
3-OCOC <sub>6</sub> H <sub>5</sub> <sup>a</sup>			165.7								
4-OCH <sub>3</sub>	57.3	56.5	57.5								
4-NHCOOCH <sub>3</sub>								157.4 (s)	157.4	157.6	157.6
4-NHCOOCH <sub>3</sub>								52.8 (q)	52.7	52.3	52.3

<sup>a</sup> Five signals were observed for the six aromatic carbons in 3 at 133.6, 133.1, 130.2, 129.8, and 128.5.

spectral data were in agreement with the proposed structures of these novel deoxysugars 1 and 2. Benzoylation of 1 afforded the 3-O-benzoate 3, mp 58–60.5 °C;  $[\alpha]_D -31.4^\circ$ ;  $\nu_{\text{max}}$  1720, 1280  $\text{cm}^{-1}$  (Tables I and II). The acidic hydrolysis also afforded methyl 2,6-dideoxy- $\alpha$ -L-ribo-hexopyranoside (methyl  $\alpha$ -L-digitoxoside) (4),  $[\alpha]_D -170.7^\circ$ , the  $\beta$  anomer 5,  $[\alpha]_D +33.2^\circ$ , methyl 2,6-dideoxy- $\alpha$ -L-ribo-hexofuranoside (6),  $[\alpha]_D -135.8^\circ$ , and the  $\beta$  anomer 7,  $[\alpha]_D +98.4^\circ$ , as oils. The  $^1\text{H}$  NMR data were consistent with data reported for methyl D-digitoxosides,<sup>4</sup> and only the  $^{13}\text{C}$  NMR data (Table II) are reported here. In all of the above digitoxose derivatives the absolute stereochemistry was shown to be L by application of Hudson's Rules of Isorotation. This sugar was reported independently to occur in three polyene antibiotics,<sup>5</sup> as well as in the tetrocarcins<sup>6</sup> and antelermicins,<sup>6,7</sup> which are closely related to kijanimicin. The acidic hydrolysis also afforded O- $\beta$ -D-kijanosyl-(1  $\rightarrow$  17)-kijanolid. From the  $^{13}\text{C}$  NMR data of kijanimicin and the above compound,<sup>8</sup> it was evident that kijanimicin contains a tetrasaccharide unit comprised of one 2,6-dideoxy-4-O-methyl- $\beta$ -L-ribo-hexopyranosyl unit and three  $\alpha$ -L-digitoxosyl units.

Acidic hydrolysis of O- $\beta$ -D-kijanosyl-(1  $\rightarrow$  17)-kijanolid by using 5 N hydrogen chloride in methanol at 65 °C for 3 h afforded

Chart I

(4) A. Zeeck, *Liebigs Ann. Chem.*, 2079 (1975).(5) J. Zielinski, E. Jereczek, P. Sowinski, L. Falkowski, A. Rudowski, and E. Borowski, *J. Antibiot.* 32, 565 (1979).(6) F. Tomita, T. Tamaoki, K. Shirahata, M. Kasai, M. Morimoto, S. Ohkubo, K. Mineura and S. Ishii, *J. Antibiot.* 33, 668 (1980).(7) K. Kobinata, M. Uramoto, T. Mizuno and K. Isono, *J. Antibiot.* 33, 244 (1980).

(8) The structure and physical data are given in the following paper.

the novel nitro sugars methyl  $\alpha$ -D-kijanoside (8) as a viscous gum,  $[\alpha]_D +130.0^\circ$  (CH<sub>3</sub>OH);  $\lambda_{\text{max}}$  (CF<sub>3</sub>CH<sub>2</sub>OH) 199 nm ( $\epsilon$  4968);

Table III.  $\Delta\delta_{\text{H}}$  Values for  $\text{H}_2$  and  $\text{H}_4$ 

H	$\delta_{\text{H}}(15)$	$\delta_{\text{H}}(9)$	stereo-chemistry <sup>a</sup>	$\Delta\delta_{\text{H}}$ (15 $\rightarrow$ 9)	H	$\delta_{\text{H}}(16)$	$\delta_{\text{H}}(17)$	stereo-chemistry <sup>a</sup>	$\Delta\delta_{\text{H}}$ (16 $\rightarrow$ 17)
$\text{H}_{2\text{eq}}$	1.76	2.69	cis	+0.93	$\text{H}_{2\text{eq}}$	2.05	3.02	cis	+0.97
$\text{H}_{2\text{ax}}$	1.38	1.54	trans	+0.16	$\text{H}_{2\text{ax}}$	1.59	1.56	trans	-0.03
$\text{H}_{4\text{eq}}$	3.45	4.44	cis	+0.99	$\text{H}_{4\text{ax}}$	4.64	4.60	trans	-0.04

<sup>a</sup> Stereochemistry of the protons at  $\text{C}_2$  and  $\text{C}_4$  relative to the heterosubstituent at  $\text{C}_3$ .

Table IV.  $\Delta\delta_{\text{C}}$  Values for  $\text{C}_2$  and  $\text{C}_4$ 

car- bon	$\delta_{\text{C}}$ (15) <sup>10</sup>	$\delta_{\text{C}}$ (9)	$\Delta\delta_{\text{C}}$ (15 $\rightarrow$ 9)	$\delta_{\text{C}}$ (18) <sup>12</sup>	$\delta_{\text{C}}$ (17) <sup>12</sup>	$\Delta\delta_{\text{C}}$ (18 $\rightarrow$ 17)
$\text{C}_2$	39.7	35.8	-3.9	43.3	39.4	-3.9
$\text{C}_4$	57.6	53.9	-3.7	77.8	77.4	-0.4

$[\theta]_{234} -326$ ,  $[\theta]_{280} +340$  ( $\text{CF}_3\text{CH}_2\text{OH}$ );  $\nu_{\text{max}}$  3430, 2980, 2950, 2940, 2900, 2840, 1735, 1550, 1508, 1230, 1123, 1055  $\text{cm}^{-1}$ ; and the  $\beta$  anomer (9), mp 180.5-181.5  $^{\circ}\text{C}$ ,  $[\alpha]_{\text{D}} +34.1^{\circ}$  ( $\text{CH}_3\text{OH}$ );  $\lambda_{\text{max}}$  ( $\text{CF}_3\text{CH}_2\text{OH}$ ) 199 nm ( $\epsilon$  4909);  $[\theta]_{234} -962$ ,  $[\theta]_{282} +2308$  ( $\text{CF}_3\text{CH}_2\text{OH}$ );  $\nu_{\text{max}}$  3440, 3000, 2955, 2900, 2860, 1730, 1550, 1515, 1315, 1235, 1065  $\text{cm}^{-1}$ . The  $^1\text{H}$  NMR (Table I) and  $^{13}\text{C}$  NMR data (Table II) were in agreement with the proposed 2,3,4,6-tetra-deoxy-4-(methoxycarbonylamino)-3-C-methyl-3-nitro-D-xylo-hexopyranosyl structures 8 and 9 ( $^4\text{C}_1$  conformation). The chemical shift of the 3-methyl ( $\delta_{\text{C}}$  26.4, 25.4) in 8 and 9 was in agreement with published values<sup>9</sup> for an equatorial C-methyl nitro sugar. Independent proof for the stereochemistry at  $\text{C}_3$  was obtained as follows. Methyl  $\beta$ -L-mycaroside (14) was converted by standard techniques into the 4-methoxycarbonylamino derivative (15). Comparison of the  $^1\text{H}$  NMR data for 15<sup>10</sup> with that of 9 (Table III) revealed marked deshielding of the vicinal cis protons  $\text{H}_{2\text{e}}$  and  $\text{H}_{4\text{e}}$  in going from the 3-axial hydroxy compound 15 to the 3-axial nitro compound 9, while little effect was observed on the vicinal trans proton  $\text{H}_{2\text{a}}$ . This was in excellent agreement with the results obtained in going from 16<sup>11</sup> to 17<sup>11</sup> (Table III). Comparison of the  $^{13}\text{C}$  NMR data for 15<sup>10</sup> with those of 9 (Table IV) revealed pronounced shielding of  $\text{C}_2$  and  $\text{C}_4$ , both of which bear vicinal cis protons to the axial nitro group.<sup>12</sup> These data were in excellent agreement with the observed shielding at  $\text{C}_2$ , which bears a vicinal cis proton, and the complete absence of any marked shielding at  $\text{C}_4$ , which does not bear a vicinal cis proton to the 3-O-acetate, in going from 18<sup>12</sup> to 17<sup>12</sup> (Table IV). Thus kijanose has the xylo configuration.

The EI mass spectra of 8 and 9 revealed no molecular ions, but ions at  $m/e$  184, 172, 156, 140, and 128 were diagnostic for structures 8 and 9. The CIMS gave  $\text{MH}^+$  ions for 8 and 9 at  $m/e$  263. Chemical proof for the presence of the nitro group was obtained by reduction of 8 and 9 by using Raney nickel to give 10 and 11 ( $\text{M}^+$ ,  $m/e$  232),<sup>10</sup> which on acetylation afforded 12 and 13.<sup>10</sup>

The application of Hudson's Rules of Isorotation to 8 and 9,  $[\text{M}]_{\text{D}} - [\text{M}]_{\text{D}} = 340.6^{\circ} - 89.3^{\circ} = 251.3^{\circ}$ , and 10 and 11,  $[\text{M}]_{\text{D}} - [\text{M}]_{\text{D}} = 281.7^{\circ} - (-11.4^{\circ}) = 292.1^{\circ}$ , indicated a D-configuration. Recently L-rubranitrose (19)<sup>13</sup> was claimed to have the L-configuration by comparison of the CD of 19 with that of L-evernitrose.<sup>14</sup> The published rotation of 19 did not agree with an L-sugar.<sup>13</sup> Comparison of the CD data for 9 with that reported for 1-O-acetyl- $\beta$ -L-rubranitrose (19) ( $\text{R} = \text{Ac}$ )<sup>13</sup> clearly indicates that both compounds have the same absolute stereochemistry. The published rotation for 19<sup>13</sup> is also in agreement with a D-config-

uration. We therefore propose that rubranitrose is a D-sugar and that the correct structure is 20. D-Kijanose represents the third naturally occurring nitro sugar to be discovered from an antibiotic.<sup>15</sup> The structure of the tetrasaccharide moiety and the location of the sugars on the aglycon in kijanimicin will be described in the following paper.

(15) Recently a nitro sugar having the same composition as D-kijanose has been isolated from the tetrocarcins,<sup>7</sup> but no structure has yet been published.

## Kijanimicin. 2.<sup>1</sup> Structure and Absolute Stereochemistry of Kijanimicin

Alan K. Mallams,\* Mohindar S. Puar, and  
Randall R. Rossman

Research Division, Schering-Plough Corporation  
Bloomfield, New Jersey 07006

Andrew T. McPhail\*

Paul M. Gross Chemical Laboratory, Duke University  
Durham, North Carolina 27706

Ronald D. Macfarlane

Department of Chemistry, Texas A&M University  
College Station, Texas 77843

Received January 19, 1981

Kijanimicin<sup>1</sup> has been shown to have the novel tetrionic acid structure 1. Kijanimicin (1) was purified by preparative HPLC on silica gel using dichloromethane-methanol-triethylamine (98:1:1) as the eluant, affording a colorless amorphous solid, mp 174.5  $^{\circ}\text{C}$  dec;  $[\alpha]_{\text{D}} -124.2^{\circ}$  ( $\text{CH}_3\text{OH}$ );  $\text{pK}_{\text{a}}$  5.0,  $\lambda_{\text{max}}$  ( $\text{CF}_3\text{C}-\text{H}_2\text{OH}$ ) 200 nm ( $\epsilon$  42832), 241 (8946), 264 (sh) (9697), 274 (9446);  $\lambda_{\text{max}}$  ( $\text{CH}_3\text{OH} + 0.1 \text{ N HCl}$ ) 205 nm ( $\epsilon$  38313), 258 (9881);  $\lambda_{\text{max}}$  ( $\text{CH}_3\text{OH} + 0.1 \text{ N NaOH}$ ) 236 nm ( $\epsilon$  14677), 266 (sh) (12002), 276 nm (12002);  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ ) 3625, 3550, 3480, 3440, 2980, 2940, 2910, 1755, 1730, 1605, 1545, 1510, 1230, 1130, and 1058  $\text{cm}^{-1}$ . The above data suggested the presence of a tetrionic acid moiety and hydroxyl, carbonyl, lactone, carbamate, nitro, and ether functions in the molecule. The  $^1\text{H}$  NMR spectrum of 1 revealed methyl groups at  $\delta_{\text{H}}$  ( $\text{CDCl}_3$ ) (220 MHz) 0.65 (3 H, d,  $J = 5 \text{ Hz}$ , 6- $\text{CH}_3$ ), 1.07 (3 H, d,  $J = 7 \text{ Hz}$ , 8- $\text{CH}_3$ ), 1.18 (3 H, d,  $J = 6 \text{ Hz}$ , 6 $^{\text{E}}$ - $\text{CH}_3$ ), 1.20-1.40 (envelope of  $\text{CH}_3$  signals), 1.60 (3 H, s, 3 $^{\text{E}}$ - $\text{CH}_3$ ), 1.64 (3 H, s, 4- $\text{CH}_3$ ), 3.45 (3 H, s, 4 $^{\text{D}}$ - $\text{OCH}_3$ ) and 3.76 (3 H, s, 4 $^{\text{E}}$ - $\text{NHCOOCH}_3$ ). The  $^{13}\text{C}$  NMR spectrum of 1 (Table I) revealed 67 carbon atoms of which five were anomeric carbons, indicating that 1 contains five sugars.<sup>1</sup> Although a satisfactory analysis (C, H, N) for 1 for  $\text{C}_{67}\text{H}_{100}\text{N}_2\text{O}_{24}$  was obtained, it could not be used to unambiguously establish the molecular composition. An EIMS of 1 gave the highest mass fragment ion at  $m/e$  552 (2.1%). Kijanimicin (1) was therefore converted into 26-O-methylkijanimicin (2) by treatment with diazomethane. The 26-O-methyl group occurred as a singlet at  $\delta_{\text{H}}$  ( $\text{CDCl}_3$ ) (600 MHz) 4.12.<sup>3</sup> The EIMS, CIMS, and FDMS

(9) K. Sato, M. Matsuzawa, K. Ajisaka, and J. Yoshimura, *Bull. Chem. Soc. Jpn.*, **53**, 189 (1980).

(10) Full details will be published in *J. Chem. Soc., Perkin Trans. 1*.

(11) R. S. Jaret, A. K. Mallams, and H. Reimann, *J. Chem. Soc., Perkin Trans. 1*, 1374 (1973).

(12) P. Bartner, D. L. Boxler, R. Brambilla, A. K. Mallams, J. B. Morton, P. Reichert, F. D. Sancilio, H. Suprenant, G. Tomalesky, G. Lukacs, A. Olesker, T. T. Thang, L. Valente and S. Omura, *J. Chem. Soc., Perkin Trans. 1*, 1600 (1979).

(13) S. A. Mizesak, H. Hoeksema, and L. M. Pschigoda, *J. Antibiot.*, **32**, 771 (1979).

(14) A. K. Ganguly, O. Z. Sarre, A. T. McPhail, and K. D. Onan, *J. Chem. Soc., Chem. Commun.*, 313 (1977).

(1) Part 1: A. K. Mallams, M. S. Puar, and R. R. Rossman, *J. Am. Chem. Soc.*, preceding paper in this issue.

(2) R. D. Macfarlane and D. F. Torgerson, *Science (Washington, D.C.)*, **191**, 920 (1976). One of us (R.D.M.) gratefully acknowledges NIH Grant GM-26096.

(3) Full details will be published in *J. Chem. Soc., Perkin Trans. 1*.