# ARTICLE

# Synthesis and elaboration of functionalised carbohydrate-derived spiroketals

Peter A. V. van Hooft,<sup>*a*</sup> Farid El Oualid,<sup>*b*</sup> Herman S. Overkleeft,<sup>*b*</sup> Gijsbert A. van der Marel,<sup>*b*</sup> Jacques H. van Boom \*<sup>*b*</sup> and Michiel A. Leeuwenburgh<sup>*b*</sup>

<sup>a</sup> TNO Prins Maurits Laboratory, P.O. Box 45, 2280 AA Rijswijk, The Netherlands

<sup>b</sup> Leiden Institute of Chemistry, Leiden University, Gorlaeus Laboratories, PO Box 9502, 2300 RA Leiden, The Netherlands. E-mail: j.boom@chem.leidenuniv.nl;
 Fax: +31 71 5274307; Tel: +31 71 5274280

Received 5th February 2004, Accepted 8th March 2004 First published as an Advance Article on the web 5th April 2004

The scope of a stereoselective three-step approach for the synthesis of sugar derived spiroketals is presented. The methodology consists of Grignard addition of vinyl- or allylmagnesium bromide to a carbohydrate lactone, followed by K-10 clay mediated glycosidation with a terminal alkenol and subsequent ring-closing metathesis of the resulting diene. The generality of this procedure is demonstrated by the synthesis of various pyranose- and furanose-derived spiroketals, as well as more advanced tricyclic spiroketal derivatives. It is shown that functionalisation of the double bond in the resulting spiroketals leads to fused polycyclic ethers.

# Introduction

Spiroketals, particularly 1,6-dioxa-spiro[4.5]decanes and 1,7dioxa-spiro[5.5]undecanes, which are common structural features of many natural products,<sup>1</sup> have attracted considerable interest from synthetic organic chemists over the past several decades. An attractive approach towards functionalised spiroketals involves the use of carbohydrates as readily available chiral synthons. For instance, Hanessian and Ugolini<sup>2</sup> reported the acid-catalysed intramolecular glycosidation of a ketose derivative to a 1,7-dioxa-spiro[5.5]undecene. Similarly, [4.4] as well as [4.5] spiroketals were produced by acid-catalysed transketalisation of keto-furanosides.<sup>3</sup> Alternatively, a 1,6-dioxaspiro[4.5]undecane framework was constructed by radical promoted cyclisation of C-glycosides.<sup>4</sup> However, the spiroketalisations described above lead to epimeric mixtures which require additional acid treatment to force the equilibrium to the thermodynamically most stable isomer. Sinaÿ and Haudrechy<sup>5</sup> nicely evaded this drawback by predispositioning the configuration at the future spirocentre through a stereoselective glycosidation of a ketose derivative. The resulting ketoglycoside was successively converted into a spiroketal via radical mediated ring-closure.<sup>6</sup> In line with this approach, we developed a threestep procedure,<sup>7</sup> comprising Grignard addition of vinyl- or allylmagnesium bromide to perbenzylated D-gluconolactone, followed by stereoselective Montmorillonite K-10 clay<sup>8</sup> mediated condensation with different terminal alkenols and subsequent formation of the spiroketal by ring-closing metathesis using Grubbs' catalyst.<sup>9</sup> We present here the scope of this approach in full detail by the transformation of pyranose and furanose derived lactones into [4.4], [4.5], [4.6] and [5.5] spiroketals, as well as the synthesis of more advanced tricyclic compounds.

# **Results and discussion**

10.1039/b401699h

ö

In a preliminary paper<sup>7a</sup> we showed that 1,6-dioxa-(5*S*)spiro[4.5]dec-3-ene  $7^{10,11}$  (see Scheme 1) and 1,7-dioxa-(6*S*)spiro[5.5]undec-9-ene  $8^{11}$  could be assembled with a high degree of stereoselectivity following the three-step reaction sequence mentioned above. In the first instance, fully benzylated gluconolactone  $1^{12}$  was converted into the known<sup>13</sup>  $\alpha$ -epimer of vinylketose 2 by nucleophilic addition of vinylmagnesium bromide. Treatment of the same lactone with allylmagnesium



bromide afforded the known<sup>14</sup> allylketose **3** as a mixture of anomers ( $\alpha$  :  $\beta$  = 4; 1). Next, Montmorillonite K-10 clay mediated glycosidation of donors **2** and **3** with an excess of allyl alcohol afforded the corresponding isomerically pure  $\alpha$ -ketoglycosides **4** and **5**. The last step, comprising RCM of dienes **4** and **5** using Grubbs' catalyst **6** (6 mol%) in toluene at 60 °C, gave the respective spiroketal derivatives **7** and **8** in 65% and 50% overall yield from **1**. The *S*-configuration of the spirocentres in compounds **7** and **8** was unambiguously assigned by NOESY experiments (see Scheme 1).<sup>15</sup> In addition, the NMRdata of saturated spiroketal **9**, obtained by hydrogenation of **7** 



# Table 1 Examples of the K-10/RCM approach towards monosaccharide-derived spiroketals

<sup>*a*</sup> Acceptor (6 equiv.), K-10 (200 mass%), 4 Å sieves, CH<sub>2</sub>Cl<sub>2</sub>, 20 °C. <sup>*b*</sup> **6** (0.06 equiv.), toluene, 60 °C. <sup>*c*</sup> **6** (0.06 equiv.), CH<sub>2</sub>Cl<sub>2</sub>, 20 °C. <sup>*d*</sup> **6** (0.03 equiv.), CH<sub>2</sub>Cl<sub>2</sub>, 20 °C.

under the influence of platinum(iv) oxide, were identical to those reported by the groups of Suárez and Taylor.<sup>4d,16</sup> Glycosidation of ketoses **2** and **3** with 3-buten-1-ol and 4-penten-1-ol and subsequent RCM as described above led to spiroketals **11**,<sup>2,17</sup> **14** and **15** in 60%, 66% and 72% yields, respectively, over the two steps.<sup>7,11</sup>

In the next stage, the influence of the carbohydrate configuration on the efficacy of the three-step methodology was studied. Therefore, the lactones derived from fully benzylated L-fuco- and L-rhamnopyranose<sup>18,19</sup> were subjected to Grignard addition of allylmagnesium bromide and K-10 mediated glycosidation with allyl alcohol (entries 1 and 2, Table 1) as described in Scheme 1. RCM of the resulting dienes 17 and 20 using Grubbs' catalyst (6 mol%) in dichloromethane at ambient temperature afforded the respective spiroketals 18 and 21 (55% and 68%, respectively), the structures of which were fully assigned by NMR-spectroscopy. Continuing our investigation, the acid-labile furanoid 2,3;5,6-di-O-isopropylidene-D-manno-1,4lactone<sup>20</sup> was subjected to the three-step sequence. Reaction of this lactone with vinylmagnesium bromide led to the isolation of the expected vinylketose 22 (entry 3) in 45% as well as a sideproduct. Mass spectrometric and NMR-spectroscopic analysis indicated that the latter compound resulted from double vinyl addition which led to an acyclic bis-adduct.<sup>21</sup> However, subsequent glycosidation of 22 and RCM steps proceeded smoothly to give spiroketal 24 in 64% yield over two steps.

Reaction of the same lactone with allylmagnesium bromide furnished allylketose **25** in excellent yield. Subjection of **25** to K-10 glycosidation with allyl alcohol as well as 3-butene-1-ol (entries 4 and 5 respectively), followed by RCM of the resulting dienes afforded spiroketals **27** and **29** in 47% and 42% yield over two steps.

The generality of the approach is further emphasised by its application in the synthesis of the more complex tricyclic spiroketal 34. Lactone 31 (see Scheme 2) was synthesised from known<sup>22</sup> pyranopyran compound **30** in two steps. Liberation of the acetal function by acid hydrolysis with 3 M sulfuric acid in acetic acid at 80 °C, followed by subsequent oxidation furnished lactone 31 in 50% yield. Execution of the three-step reaction sequence on this lactone, as described for the conversion of 2 into 7, proceeded in 20% overall yield to afford spiroketal 34, the structure of which was ascertained by NOESYexperiments. A different tricyclic constellation, namely 37 (see Scheme 3), was accessible through K-10-mediated glycosidation of vinylketose 2 with sugar derived acceptor 35.23 Remarkably, this glycosidation only proceeded in satisfactory yield when 35 was used in a 20-fold excess. RCM of the resulting disaccharide 36 with Grubbs' catalyst 6 (6 mol%) furnished tricyclic adduct 37 in 64% yield.

Interestingly, yet another opportunity to construct oligocyclic systems presents itself by elaboration of the double bond in the spiroketals, as is demonstrated for spiroketal **8** (see



Scheme 2 Reagents and conditions: i, a) 3 M  $H_2SO_4$ , HOAc, 80 °C, 6 h, b) DMSO, acetic anhydride, 16 h (50%). ii, Vinylmagnesium bromide, Et<sub>2</sub>O, -78 °C (42%). iii, Allyl alcohol (6 equiv.), K-10 clay (200 mass%), 4 Å sieves, CH<sub>2</sub>Cl<sub>2</sub> (50%). iv, **6** (0.06 equiv.), CH<sub>2</sub>Cl<sub>2</sub> (95%).



Scheme 3 Reagents and conditions: i, 35 (20 equiv.), K-10 clay (200 mass%), 4 Å sieves,  $CH_2Cl_2$  (45%). ii, 6 (0.06 equiv.),  $CH_2Cl_2$  (64%).



Scheme 4 Reagents and conditions: i,  $(Ph_3P)_3RhCl$  (cat.), DBU, EtOH, 60 °C (90%). ii, a) DMDO,  $CH_2Cl_2$ , 0 °C, b) allylmagnesium chloride, THF, 0 °C (61%). iii, Allyl bromide, NaH, DMF (93%). iv. catalyst 41 (0.5 mol%),  $CH_2Cl_2$ ,  $\Delta$  (80%).

Scheme 4). Conversion of **8** into enol ether **38** was effected in 90% yield by reaction with Wilkinson's catalyst<sup>24</sup> in the presence of a catalytic amount of DBU in ethanol at 60 °C.<sup>22a</sup> Next, **38** was treated with a freshly prepared solution of 3,3-dimethyldioxirane (DMDO) in acetone at 0 °C,<sup>25</sup> followed by

opening of the crude epoxide with allylmagnesium bromide in THF<sup>26</sup> at 0 °C. The resulting alkenol **39** was isolated in 61% yield. Allylation under standard conditions, followed by RCM using 0.5 mol% Grubbs' catalyst **41**<sup>27</sup> in refluxing dichloromethane afforded tricyclic product **42** in 74% over two steps. The structure of **42** was confirmed by NOESY-experiments, of which the most important signals are depicted in Scheme 4.

# Conclusion

In this paper, the scope of a stereoselective and versatile threestep approach towards the construction of pyranose (D-gluco, L-fuco and L-rhamno) and furanose (D-manno) derived unsaturated spiroketals is presented. The methodology has proven to be effective for the construction of [4.4], [4.5], [5.5], [5.6] and [5.7] spiroketals and also of the more advanced tricvclic compounds 34 and 37. It was demonstrated that the double bond in the newly formed spirosystem is amenable to further elaboration, to furnish fused oxacyclic compounds such as 42. It is of interest to note that the allylic ether 42 can be subjected to the same sequence of reactions, to give entrance to a reiterative procedure.<sup>28</sup> Its broad applicability, as well as the potential to further elaborate the produced spiroketals, make this three-step methodology a valuable tool in the synthesis of natural products containing highly functionalised spiroketal moieties.

# Experimental

# General

Dichloromethane, dimethylsulfoxide (DMSO), ethyl acetate, chloroform, NN-dimethylformamide (DMF), pyridine (p.a. grade, Baker) and tetrahydrofuran (THF, Baker, HPLC grade) were stored over molecular sieves (4 Å). Toluene was boiled under reflux with P2O5 for 3 h, distilled and stored over molecular sieves (4 Å). Anhydrous diethyl ether was freshly destilled from LiAlH<sub>4</sub> prior to use. Sulfuric acid, acetic acid, acetic anhydride (p.a. grade, Baker), ethanol (96%, technical grade) and acetone (Acros, p.a.) were used as received. Vinylmagnesium bromide (1 M in THF, Aldrich), allyl alcohol (Aldrich), platinum(IV) oxide (Acros), allylmagnesium bromide (1 M in Et<sub>2</sub>O, Aldrich), 3-buten-1-ol (Aldrich), chlorotris(triphenylphosphine)rhodium(I) ((PPh<sub>3</sub>)<sub>3</sub>Rh(I)Cl, Aldrich), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, Aldrich), allylmagnesium chloride (2 M in THF, Aldrich), allyl bromide (Aldrich), and sodium hydride (60% dispersion in mineral oil, Acros) were used as received. Montmorillonite K-10 clay (Aldrich) was dried (110 °C, 1 mm Hg, 10 h) before use. Reactions were executed at ambient temperature unless stated otherwise. Drying of organic layers after work-up was effected by addition of MgSO<sub>4</sub>. TLC analysis was conducted on DC-Fertigfolien (Schleicher & Schuell, F1500, LS254) or HPTLC aluminum sheets (Merck, silica gel 60, F254) with detection by UVabsorption (254 nm) and charring with 20% H<sub>2</sub>SO<sub>4</sub>in ethanol. Column chromatography was performed either on Baker silica gel (0.063-0.200 mm) or Merck silica gel 60 (0.040-0.063 mm). Solvents used for column chromatography were of technical grade and distilled before use. 1H- and 13C-spectra were recorded on a Jeol JNM-FX-200 (200 MHz and 50.1 MHz, respectively) or a Bruker DPX-300 (300 MHz and 75.1 MHz, respectively). NMR shifts are reported in ppm ( $\delta$ ) relative to tetramethylsilane. Mass spectrometry was performed on a PE/SCIEX API 165 equipped with an electrospray interface. Optical rotation values were measured at 20 °C on a Propol Automatic Polarimeter at 589 nm and are given in unit of  $10^{-1}$ deg cm<sup>2</sup> g<sup>-1</sup>.

The atom numbering used for the bi- and tricyclic products is given in Fig. 1.



Fig. 1 The atom numbering used for the bi- and tricyclic products.

# General procedure for the Grignard reactions

A solution of the lactone in anhydrous  $Et_2O$  (5 mL mmol<sup>-1</sup>) was cooled to -78 °C under a nitrogen atmosphere. The Grignard reagent (1.0 M solution, 1.2 equiv.) was added dropwise and the mixture was stirred at -78 °C until TLC analysis (33% EtOAc–light petroleum ether) revealed completion of the reaction. The reaction was quenched by addition of sat. NH<sub>4</sub>Cl and extracted with Et<sub>2</sub>O. The organic layer was washed with sat. NH<sub>4</sub>Cl, dried and concentrated. Silica gel chromatography (10 to 30% EtOAc–light petroleum ether) yielded the ketose.

# General procedure for the K-10-mediated condensation reactions

To a flask containing Montmorillonite K-10 clay and powdered molecular sieves 4 Å (both flame-dried under vacuum, 200 mass% each), was added a mixture of the ketose and the acceptor (6 equiv.) in anhydrous  $CH_2Cl_2(10 \text{ mL mmol}^{-1})$  under an argon atmosphere. The resulting suspension was stirred vigorously until TLC analysis (25% EtOAc–light petroleum ether) revealed completion of the reaction. A mixture of methanol–pyridine (10 mL mmol<sup>-1</sup>, 1 : 1) was added and stirring was continued for 10 min. The suspension was filtered over Hyflo<sup>®</sup> and concentrated. Silica gel chromatography (5 to 20% EtOAc–light petroleum ether) of the residue yielded the glycosidation product.

# General procedures for ring-closing metathesis

Method A: A 0.08 M solution of the diene in anhydrous toluene was purged for 15 min with argon. Grubbs' catalyst **6** (0.06 equiv.) was added and the mixture was purged with argon for an additional 15 min, after which the solution was heated to 60 °C. Stirring was continued under an argon atmosphere until TLC analysis (20% EtOAc–light petroleum ether) revealed complete disappearance of the starting material. The mixture was allowed to cool to room temperature and was concentrated. The residue was subjected to silica gel chromatography (5 to 20% EtOAc–light petroleum ether) to give the product.

Method B: A 0.08 M solution of the diene in anhydrous dichloromethane was purged for 15 min with argon. Grubbs' catalyst **6** (0.06 equiv.) was added and the mixture was purged with argon for an additional 15 min. Stirring was continued under an argon atmosphere until TLC analysis (20% EtOAc– light petroleum ether) revealed complete disappearance of the starting material. The mixture was concentrated, and purified by silica gel chromatography (5 to 20% EtOAc–light petroleum ether) to give the product.

# 4,5,6,8-Tetra-O-benzyl-1,2-dideoxy-α-D-gluco-oct-1-eno-3-

**ulopyranose (2).** Prepared from 1 by Grignard addition, according to the general procedure. Colourless syrup (3.82 g, 6.87 mmol, 91%);  $[a]_{\rm D}$  +46.5 (*c* 1 in CHCl<sub>3</sub>);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.14–7.28 (m, 20H, CH<sub>arom</sub>), 5.99 (dd, 1H, *J* 11.0 and 17.5 Hz, H-2), 5.26 and 5.60 (2d, 2H, *J* 10.2 and 16.8 Hz, 2 × H-1), 4.54, 4.63, 4.79 and 4.88 (4d, 8H, CH<sub>2</sub> Bn), 4.05–4.10 (m, 1H, H-7), 4.01 (br t, 1H, *J* 9.5 Hz, H-5), 3.65–3.78 (m, 2H, H-6,8), 3.46 (dd, 1H, *J* 3.7 and 7.3 Hz, H-8), 3.42 (br d, 1H, *J* 9.5 Hz, H-4), 3.20 (br s, 1H, OH);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.6 (C-2), 137.7, 138.1 and 138.6 (C<sub>q</sub> Bn), 127.4–128.2 (CH<sub>arom</sub>), 117.0 (C-1), 96.4 (C-3), 71.4, 78.0, 82.6 and 83.2 (C-4,5,6,7), 73.1, 74.8, 75.5 and 76.4 (CH<sub>2</sub> Bn), 68.6

(C-8); MS (ESI): calc. for  $C_{33}H_{38}O_6$  566.3, found *m*/*z* 589.3 (M + Na<sup>+</sup>).

**5,6,7,9-Tetra-***O***-benzyl-1,2,3-trideoxy-D***-gluco***-non-1-eno-4-ulopyranose (3).** Prepared from 1 by Grignard addition, according to the general procedure. Colourless syrup (3.13 g, 5.40 mmol, 32 %,  $\alpha/\beta$  = 4 : 1);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.29–7.43 (m, 20H, CH<sub>arom</sub>), 5.99–6.03 (m, 1H, H-2), 5.30 (d, 2H, 2x H-1), 4.68, 4.71, 4.81, 4.97 and 5.05 (5d, 8H, CH<sub>2</sub> Bn), 4.18 (br t, 1H, *J* 9.5 Hz, H-6), 4.10–4.17 (m, 1H, H-8), 3.66–3.89 (m, 2H, 2 × H-9), 3.78 (br t, 1H, *J* 10.2 Hz, H-7), 3.57 (br d, 1H, *J* 9.5 Hz, H-5), 2.58 (d, 2H, *J* 6.6 Hz, H-3);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.8, 138.0 and 138.4 (C<sub>q</sub> Bn), 132.1 (C-2), 126.7–128.2 (CH<sub>arom</sub>), 119.8 (C-1), 97.6 (C-4), 71.3, 78.3, 81.2 and 83.5 (C-5,6,7,8), 73.1, 74.7, 75.1 and 75.4 (CH<sub>2</sub> Bn), 68.5 (C-9), 42.6 (C-3); MS (ESI): calc. for C<sub>37</sub>H<sub>40</sub>O<sub>6</sub> 580.3, found *m*/*z* 603.4 (M + Na)<sup>+</sup>.

Allyl (4,5,6,8-tetra-O-benzyl-1,2-dideoxy)-α-D-gluco-oct-1eno-3-ulopyranoside (4). Prepared from 2 by K-10-mediated glycosidation according to the general procedure. Colourless syrup (238 mg, 0.39 mmol, 74%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.34-7.43 (m, 20H, CH<sub>arom</sub>), 5.99-6.12 (m, 2H, J 11.0, 17.5 Hz, H-2 and CH All), 5.69 (dd, 1H, J 1.5, 17.5 Hz, H-1), 5.40 (dd, 1H, J 1.5, 11.0 Hz, H-1), 5.35 (d, 1H, J 6.6 Hz, CH<sub>2</sub> All), 5.23 (d, 1H, J 10.2 Hz, CH<sub>2</sub> All), 4.68, 4.73, 4.78, 4.99, and 5.00 (5d, 8H, CH<sub>2</sub> Bn), 4.29 (br t, 1H, J 8.8 Hz, H-5), 4.06 (t, 2H, J 5.9 Hz, CH<sub>2</sub> All), 3.81-3.94 (m, 4H, H-6,7 and  $2 \times$  H-8), 3.50 (d, 1H, J 9.5 Hz, H-4);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.3, 138.4 and 138.8 (Cq Bn), 135.4 (C-2), 134.8 (CH All), 127.6-128.4 (CH<sub>arom</sub>), 118.7 (C-1), 116.6 (CH<sub>2</sub> All), 99.8 (C-3), 71.9, 78.5, 83.0 and 84.3 (C-4,5,6,7), 73.3, 75.0, 75.6 and 75.8 (CH<sub>2</sub> Bn), 68.8 (C-8), 63.0 (CH<sub>2</sub> All); MS (ESI): calc. for C<sub>39</sub>H<sub>42</sub>O<sub>6</sub> 606.3, found m/z 629.5 (M + Na)<sup>+</sup>.

Allyl (5,6,7,9-tetra-*O*-benzyl-1,2,3-trideoxy)-α-D-gluco-non-1eno-4-ulopyranose (5). Prepared from 3 by K-10-mediated glycosidation according to the general procedure. Colourless syrup (228 mg, 0.37 mmol, 69%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.21– 7.30 (m, 20H, CH<sub>arom</sub>), 5.81–6.02 (m, 2H, H-2 and CH allyl), 5.28 (d, 1H, *J* 17.5 Hz, H-1), 5.05–5.14 (m, 3H, H-1 and CH<sub>2</sub> allyl), 4.52, 4.57, 4.63, 4.71, 4.84 and 4.93 (6d, 8H, CH<sub>2</sub> Bn), 4.14 (br t, 1H, *J* 9.5 Hz, H-6), 4.07 (d, 2H, *J* 5.8, CH<sub>2</sub> All), 3.61– 3.77 (m, 4H, H-7,8 and 2 × H-9), 3.53 (br d, 1H, *J* 9.5 Hz, H-4), 2.63 (d, 2H, *J* 7.3 Hz, 2 × H-3); $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.3 and 138.7 (C<sub>q</sub> Bn), 135.0 (C-2), 133.6 (CH allyl), 127.5– 128.4 (CH<sub>arom</sub>), 118.1 (CH<sub>2</sub> allyl), 116.4 (C-1), 101.9 (C-4), 72.0, 78.8, 81.2 and 83.6 (C-5,6,7,8), 73.3, 74.9, 75.1 and 75.5 (CH<sub>2</sub> Bn), 68.9 (C-9), 61.7 (CH<sub>2</sub>allyl), 42.6 (C-3); MS (ESI): calc. for C<sub>40</sub>H<sub>44</sub>O<sub>6</sub> 620.3, found *m*/*z* 643.4 (M + Na)<sup>+</sup>.

### (5S,7R,8R,9S,10R)-8,9,10-Tris-benzyloxy-7-benzyloxy-

**methyl-1,6-dioxa-spiro[4.5]dec-3-ene** (7). Obtained by RCM of **4**, *via* method A of the general procedure, as a greenish syrup (216 mg, 0.37 mmol, 95%);  $[a]_{\rm D}$  + 67.2 ° (*c* 1 in CH<sub>2</sub>Cl<sub>2</sub>);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.11–7.24 (m, 20H, CH<sub>arom</sub>), 6.19 (d, 1H, *J* 5.8 Hz, H-4), 5.62 (m, 1H, H-3), 4.44, 4.54, 4.59, 4.71, 4.74, 4.84, 4.85 and 4.93 (8d, 8H, CH<sub>2</sub> Bn), 4.04 (br t, 1H, *J* 9.5 Hz, H-9), 3.98 (dd, 1H, *J* 6.6 Hz, H-11), 3.71–3.80 (m, 3H, H-7,8,11), 3.64 (d, 1H, *J* 9.5 Hz, H-10), 2.25 (d, 2H, *J* 11.7 Hz, 2 × H-2);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.0 (C<sub>q</sub> Bn), 138.3 and 138.7 (C<sub>q</sub> Bn), 132.2 (C-4), 127.2–128.2 (C-3 and CH<sub>arom</sub>), 112.1 (C-5), 72.3, 77.8, 81.3 and 83.6 (C-7,8,9,10), 73.3 and 74.7 (CH<sub>2</sub> Bn), 72.2 (C-2), 68.5 (CH<sub>2</sub>OBn); MS (ESI): calc. for C<sub>37</sub>H<sub>38</sub>O<sub>6</sub> 578.3, found *m/z* 601.3 (M + Na)<sup>+</sup>.

(2*R*,3*R*,4*S*,5*R*,6*S*)-3,4,5-Tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.5]undec-9-ene (8). Obtained by RCM of 5 *via* method A of the general procedure as a greenish syrup (634 mg, 1.1 mmol, 90%);  $[a]_{\rm D}$  +34.2 (*c* 0.57 in CH<sub>2</sub>Cl<sub>2</sub>);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.25–7.40 (m, 20H, CH<sub>arom</sub>), 5.77 (m, 2H, H-9,10), 4.60, 4.66, 4.71, 4.78, 4.96 and 5.07 (6d, 8H, CH<sub>2</sub> Bn), 4.30 (br t, 1H, *J* 8.8 Hz, H-4), 4.22 (br s, 2H, 2 × H-8), 3.78–3.89 (m, 4H, H-2,3 and CH<sub>2</sub>OBn), 3.47 (d, 1H, *J* 9.5 Hz, H-5), 2.63 (d, 1H, *J* 17.5 Hz, H-11), 1.88 (d, 1H, *J* 16.8 Hz, H-11);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.7, 138.0 and 138.5 (C<sub>q</sub> Bn), 127.5–128.5 (CH<sub>arom</sub>), 123.6 (C-9), 121.2 (C-10), 97.2 (C-6), 71.6, 78.5, 82.0 and 83.3 (C-2,3,4,5), 73.2, 74.8, 75.5, and 75.7 (CH<sub>2</sub>Bn), 68.5 (CH<sub>2</sub>OBn), 59.8 (C-8), 29.9 (C-11); MS (ESI): calc. for C<sub>38</sub>H<sub>40</sub>O<sub>6</sub> 592.3, found *m*/*z* 615.3 (M + Na)<sup>+</sup>.

### (5S,7R,8R,9S,10R)-8,9,10-Tris-benzyloxy-7-benzyloxy-

methyl-1,6-dioxa-spiro[4.5]decane (9). Compound 7 (47 mg, 81 µmol) was dissolved in anhydrous EtOAc (3 mL) and platinum(IV) oxide (10 mg) was added. The mixture was degassed three times after which hydrogen gas was introduced. The mixture was stirred for 5 h when TLC analysis (25% EtOAc-light petroleum ether) revealed a completed conversion. The mixture was purged with nitrogen gas, filtered over Hyflo® and concentrated. Silica gel chromatography (10% EtOAc-light petroleum ether) of the residue yielded 9 as a colourless syrup (40 mg, 69  $\mu$ mol, 85%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.33–7.20 (m, 20H, CH<sub>arom</sub>), 4.54, 4.58, 4.71, 4.87, 4.93 and 4.99 (6d, 8H, CH<sub>2</sub> Bn), 4.07 (br t, 1H, J 9.5 Hz, H-9), 4.04 (m, 1H, CH<sub>2</sub>OBn), 3.84-3.91 (m, 2H, H-2 and CH<sub>2</sub>OBn), 3.67-3.79 (m, 3H, H-2,7,8), 3.58 (d, 1H, J 9.5 Hz, H-10), 1.86-1.97 (m, 4H, 2  $\times$  H-3 and 2  $\times$  H-4);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.0, 138.1, 138.3 and 138.7 ( $C_q$  Bn), 127.5–128.4 ( $CH_{arom}$ ), 107.4 (C-5), 71.2, 78.5, 80.0 and 84.5 (C-7,8,9,10), 75.5, 75.6 and 76.4 (CH<sub>2</sub> Bn), 68.7 (CH<sub>2</sub>OBn), 68.1 (C-8), 33.4 (C-4), 24.0 (C-3); MS (ESI): calc. for  $C_{37}H_{40}O_6$  580.3, found m/z 603.4  $(M + Na)^{+}$ .

3-Buten-1-yl (4,5,6,8-tetra-O-benzyl-1,2-dideoxy)-a-D-glucooct-1-eno-3-ulopyranoside (10). Prepared from 2 by K-10mediated glycosidation according to the general procedure. Colourless syrup (223 mg, 0.36 mmol, 68%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.22-7.30 (m, 20H, CH<sub>arom</sub>), 5.92 (dd, 1H, J 11.0 and 17.5 Hz, H-2), 5.75 (dd, 1H, J 7.3 and 17.5 Hz, CH Bu), 5.56 (dd, 1H, J 1.5 and 17.5 Hz, H-1), 5.26 (dd, 1H, J 2.2 and 11.0 Hz, H-1), 5.06 (d, 1H, J 15.3 Hz, CH<sub>2</sub> Bu), 4.97 (d, 1H, J 5.1 Hz, CH<sub>2</sub> Bu), 4.54, 4.60, 4.64, 485, and 4.87 (5d, 8H, CH<sub>2</sub> Bn), 4.15 (br t, 1H, J 8.8 Hz, H-5), 3.71 (br t, 1H, J 8.8 Hz, H-6), 3.62-3.81 (m, 2H, CH<sub>2</sub> Bu), 3.42-3.52 (m, 3H, H-7 and 2 × H-8), 3.35 (d, 1H, J 9.5 Hz, H-4), 2.36 (dd, 2H, J 7.3 and 13.9 Hz, CH<sub>2</sub> Bu); δ<sub>C</sub> (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.3, 138.5 and 138.9 (C<sub>a</sub> Bn), 135.6 (C-2), 135.1 (CH Bu), 127.6-128.4 (CH<sub>arom</sub>), 118.5 (C-1), 116.5 (CH<sub>2</sub> Bu), 99.5 (C-3), 71.8, 78.6, 83.0 and 84.5 (C-4,5,6,7), 73.4, 75.0, 75.5 and 75.7 (CH<sub>2</sub> Bn), 68.9 (C-8), 61.2 (CH2 Bu), 34.2 (CH2 Bu); MS (ESI): calc. for  $C_{40}H_{44}O_6$  620.3, found *m*/*z* 643.4 (M + Na)<sup>+</sup>.

(2R,3R,4S,5R,6S)-3,4,5-Tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.5]undec-10-ene (11). Obtained from 10 by RCM according to the general procedure (method A) as a colourless syrup (188 mg, 0.32 mmol, 88%); [a]<sub>D</sub> +40.4 (c 1 in CH<sub>2</sub>Cl<sub>2</sub>);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.06–7.68 (m, 20H, CH<sub>arom</sub>), 6.10 (dd, 1H, J 4.7 and 9.9 Hz, H-10), 5.53 (dd, 1H, J 1.6 and 10.1 Hz, H-11), 4.51, 4.62, 4.69, 4.78, 4.83 and 4.93 (6d, 8H, CH<sub>2</sub> Bn), 4.06 (br t, 1H, J 9.3 Hz, H-4), 3.88 (ddd, 1H, J 1.9, 3.6 and 10.3 Hz, H-2), 3.86 (m, 1H, H-8), 3.75 (dd, 1H, J 3.8 and 10.8 Hz, CH<sub>2</sub>OBn), 3.65 (br t, 1H, J 9.9 and 9.0 Hz, H-3), 3.64 (dd, 1H, J 1.6 and 10.6 Hz, CH<sub>2</sub>OBn), 3.48 (br d, 1H, J 9.6 Hz, H-5), 2.43 (m, 1H, H-9), 1.89 (m, 1H, H-9);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.0, 138.2, 138, 3 and 138.8 (C<sub>q</sub> Bn), 130.3 (C-10), 127.6–128.6 (C-11 and CH<sub>arom</sub>), 95.6 (C-6), 71.0, 78.2, 82.9 and 83.2 (C-2,3,4,5), 73.4, 74.9, 75.3 and 75.7 (CH<sub>2</sub>Bn), 68.8 (CH<sub>2</sub>OBn), 58.1 (C-8), 24.2 (C-9); MS (ESI): calc. for  $C_{38}H_{40}O_6$  592.3, found m/z 615.3 (M + Na)<sup>+</sup>,  $631.4 (M + K)^+$ .

(5,6,7,9-tetra-O-benzyl-1,2,3-trideoxy)-α-D-3-Buten-1-yl gluco-non-1-eno-4-ulopyranose (12). Prepared from 3 by K-10mediated glycosidation according to the general procedure. Colourless syrup (181 mg, 0.29 mmol, 69%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.22-7.31 (m, 20H, CH<sub>arom</sub>), 5.80 (m, 2H, H-2 and CH Bu), 5.01–5.12 (m, 4H,  $2 \times$  H-1 and CH<sub>2</sub> Bu), 4.53, 4.58, 4.65, 4.70, 4.85 and 4.91 (6d, 8H, CH<sub>2</sub> Bn), 4.12 (br t, 1H, J 9.5 Hz, H-6), 3.48–3.78 (m, 7H, H-5,7,8 and 2 × H-9 and CH<sub>2</sub> Bu), 2.64 (m, 2H, 2 × H-3), 2.39 (m, 2H, CH<sub>2</sub> Bu);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.8 (C<sub>q</sub> Bn), 135.0 (C-2), 133.7 (CH Bu), 127.5-128.4 (CH<sub>arom</sub>), 118.0 (CH<sub>2</sub> Bu), 116.7 (C-1), 101.5 (C-4), 72.0, 78.8, 81.3 and 83.4 (C-5,6,7,8), 73.2, 74.8, 75.0 and 75.4 (CH<sub>2</sub> Bn), 68.9 (C-9), 60.0 (CH<sub>2</sub> Bu), 38.0 (C-3), 34.4 (CH<sub>2</sub> Bu); MS (ESI): calc. for  $C_{41}H_{46}O_6$  634.3, found *m*/*z* 657.5 (M  $+ Na)^{+}$ .

(5,6,7,9-tetra-O-benzyl-1,2,3-trideoxy)-α-D-4-Penten-1-yl gluco-non-1-eno-4-ulopyranose (13). Prepared from 3 by K-10mediated glycosidation according to the general procedure. Colourless syrup (638 mg, 0.98 mmol, 80%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.47-7.52 (m, 20H, CH<sub>arom</sub>), 5.94-6.11 (m, 2H, H-2 and CH Pent), 4.76, 4.85, 4.86, 4.95, 5.11, 5.13 and 5.15 (7d, 8H, CH<sub>2</sub> Bn), 4.38 (br t, 1H, J 9.5 Hz, H-6), 3.71-4.01 (m, 7H, H-5,7,8 and 2  $\times$  H-9 and CH<sub>2</sub> Pent), 2.86 (m, 2H, CH<sub>2</sub> Pent), 2.35 (m, 2H, 2 × H-3), 1.94–2.00 (m, 2H, CH<sub>2</sub> pent);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.4, 138.7 and 138.9 (C<sub>q</sub> Bn), 138.3 (CH Pent), 134.0 (C-2), 127.6–128.5 (CH<sub>arom</sub>), 118.0 (C-1), 115.0 (CH, Pent), 101.6 (C-4), 72.1, 78.9, 81.4 and 83.6 (C-5,6,7,8), 73.4, 74.8, 75.1 and 75.5 (CH<sub>2</sub> Bn), 69.0 (C-9), 59.8 (CH<sub>2</sub> Pent), 38.1 (C-3), 30.6 (CH<sub>2</sub> Pent), 29.1 (CH<sub>2</sub> Pent); MS (ESI): calc. for C<sub>42</sub>H<sub>48</sub>O<sub>6</sub> 648.3, found m/z 671.3  $(M + Na)^{+}$ .

(2*R*,3*R*,4*S*,5*R*,6*S*)-3,4,5-Tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.6]dodec-10-ene (14). Obtained from 12 by RCM according to the general procedure (method A) as a colourless syrup (128 mg, 0.21 mmol, 95%);  $[a]_{\rm D}$  +47.4 (*c* 1 in CH<sub>2</sub>Cl<sub>2</sub>);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.19–7.32 (m, 20H, CH<sub>arom</sub>), 5.69 (m, 1H, H-11), 5.52 (m, 1H, H-10), 4.52, 4.55, 4.61, 4.70, 4.86 and 4.92 (6d, 8H, CH<sub>2</sub> Bn), 4.07 (br t, 1H, *J* 9.5 Hz, H-4), 3.69–3.90 (m, 5H, H-2 and 2 × H-8 and CH<sub>2</sub>OBn), 3.63 (br t, 1H, *J* 9.5 Hz, H-3), 3.33 (d, 1H, *J* 9.5 Hz, H-5), 2.75 (d, 2H, *J* 16.1 Hz, H-12), 2.19–2.42 (m, 3H, H-12 and 2 × H-9);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.7 (C<sub>q</sub> Bn), 129.7 (C-11), 127.6– 128.7 (CH<sub>arom</sub>), 124.7 (C-10), 101.4 (C-6), 71.5, 78.7, 83.6 and 84.7 (C-2,3,4,5), 73.3, 74.9, and 75.6 (CH<sub>2</sub> Bn), 68.7 (CH<sub>2</sub>OBn), 61.8 (C-8), 36.3 (C-9), 31.3 (C-12); MS (ESI): calc. for C<sub>39</sub>H<sub>42</sub>O<sub>6</sub> 606.3, found *m*/z 629.4 (M + Na)<sup>+</sup>.

(2R,3R,4S,5R,6S)-3,4,5-Tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.7]tridec-11-ene (15). Obtained from 13 by RCM according to the general procedure (method A) as a greenish syrup (319 mg, 0.51 mmol, 90%); [a]<sub>D</sub>+15.8 (c 1 in CHCl<sub>3</sub>);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.30–7.41 (m, 20H, CH<sub>arom</sub>), 5.89 (m, 2H, H-11,12), 4.68, 4.72, 4.78, 4.86, 5.00, 5.05 and 5.09 (7d, 8H, CH<sub>2</sub> Bn), 4.19 (br t, 1H, J 8.8 Hz, H-4), 4.05 (d, 1H, J 11.7 Hz, CH<sub>2</sub>OBn), 3.59-3.84 (m, 5H, H-2,3 and 2 × H-8 and CH<sub>2</sub>OBn), 3.57 (d, 1H, J 9.5 Hz, H-5), 2.79 (dd, 1H, J 5.8 and 13.9 Hz, H-13), 2.04–2.45 (m, 4H, 2 × H-9 and  $2 \times$  H-10);  $\delta_{C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.8, 137.9, 138.1 and 138.6 (C<sub>g</sub> Bn), 131.2 (C-12), 127.3–128.2 (CH<sub>arom</sub>), 126.3 (C-11), 102.0 (C-6), 72.0, 78.4, 83.5 and 84.7 (C-2,3,4,5), 73.1, 74.8, and 75.4 (CH<sub>2</sub> Bn), 68.4 (CH<sub>2</sub>OBn), 59.2 (C-8), 35.0 (C-13), 30.7 (C-10), 23.7 (C-9); MS (ESI): calc. for C<sub>40</sub>H<sub>44</sub>O<sub>6</sub> 620.3, found m/z 643.4 (M + Na)<sup>+</sup>.

**5,6,7-Tri-***O*-benzyl-1,2,3-trideoxy-α-L-*fuco*-non-1-eno-4ulopyranose (16). Prepared from 5,6,7-tri-*O*-benzyl-L-fuco-1,5lactone <sup>16</sup> by Grignard addition, according to the general procedure. Colourless syrup (1.42 g, 2.99 mmol, 87%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.22–7.39 (m, 15H, CH<sub>arom</sub>), 5.82–5.94 (m, 1H, H-2), 5.16 (dd, 2H, J 4.4 and 5.8 Hz, 2 × H-1), 4.73 (m, 3H, CH<sub>2</sub> Bn), 4.70, 4.98 and 4.99 (3d, each 1H, CH<sub>2</sub> Bn), 3.78–4.05 (m, 3H, H-5,6,8), 3.66 (d, 1H, J 1.5 Hz, H-7), 2.75 (s, 1H, OH), 2.35–2.55 (m, 2H, 2 × H-3), 1.14 (d, 3H, J 6.6 Hz, CH<sub>3</sub>);  $\delta_{\rm c}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.4 (C<sub>q</sub> Bn), 132.9 (C-1), 127.5–128.9 (CH<sub>arom</sub>), 119.0 (C-2), 98.0 (C-4), 77.5, 78.0 and 81.4 (C-5,6,7), 72.5, 74.4 and 75.3 (CH<sub>2</sub> Bn), 67.1 (C-8), 43.2 (C-3), 16.9 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>30</sub>H<sub>34</sub>O<sub>5</sub> 474.2, found *m*/*z* 497.2 (M + Na)<sup>+</sup>.

Allyl (5,6,7-tri-O-benzyl-1,2,3-trideoxy)-a-L-fuco-non-1-eno-4-ulopyranose (17). Prepared from 16 by K-10-mediated glycosidation according to the general procedure. Colourless syrup (1.07 g, 2.08 mmol, 70%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.26– 7.44 (m, 15H, CH<sub>arom</sub>), 5.84–6.00 (m, 2H, H-2 and CH allyl), 5.26 and 5.34 (2d, 2H, J 1.5 and 16.8 Hz, CH<sub>2</sub> allyl), 5.12 and 5.17 (2d, 2H, J 1.5 and 10.2 Hz, 2 × H-1), 4.71, 4.74 and 5.00 (3d, each 1H, CH<sub>2</sub> Bn), 4.78 and 5.04 (m, 3H, CH<sub>2</sub> Bn), 3.99-4.18 (m, 3H, H-5,6,8), 3.80 (dd, 2H, J 6.6 and 13.1 Hz, CH<sub>2</sub> allyl), 3.71 (d, 1H, J 2.2 Hz, H-7), 2.66 (m, 2H, H-3), 1.20 (d, 3H, J 5.8 Hz, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.9 and 139.2 (C<sub>a</sub> Bn), 135.5 (C-1), 134.2 (CH<sub>2</sub> allyl), 127.0-128.8 (CH<sub>arom</sub>), 117.5 (C-2), 116.2 (CH allyl), 102.1 (C-4), 78.0, 78.2 and 81.2 (C-5,6,7), 72.7, 74.5 and 75.1 (CH<sub>2</sub> Bn), 67.5 (C-8), 61.6 (CH<sub>2</sub> allyl), 39.0 (C-3), 16.9 (CH<sub>3</sub>); MS (ESI): calc. for  $C_{33}H_{38}O_5$  514.3, found *m*/*z* 537.4 (M + Na)<sup>+</sup>.

(2*S*,3*R*,4*R*,5*S*,6*R*)-3,4,5-Tris-benzyloxy-2-methyl-1,7-dioxaspiro[5.5]undec-9-ene (18). Obtained from 17 by RCM according to the general procedure (method B) as a greenish syrup (0.90 g, 1.85 mmol, 90%);  $[a]_D$  –51.4 (*c* 1 in CHCl<sub>3</sub>);  $\delta_H$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.30–7.24 (m, 15H, CH<sub>arom</sub>), 5.66 (s, 2H, H-9,10), 4.71, 4.99 and 5.00 (3d, each 1H, CH<sub>2</sub> Bn), 4.77 (s, 3H, CH<sub>2</sub> Bn), 4.04–4.19 (m, 3H, H-2,4,5), 3.80 (dd, 2H, *J* 10.2 and 15.3 Hz, H-8), 3.69 (d, 1H, *J* 1.5 Hz, H-3), 2.57 (m, 1H, H-11), 1.82 (m, 1H, H-11), 1.16 (d, 3H, *J* 6.6 Hz, CH<sub>3</sub>);  $\delta_C$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.2, 138.7 and 138.8 (C<sub>q</sub> Bn), 127.6–128.7 (CH<sub>arom</sub>), 121.7 and 123.8 (C-9,10), 97.6 (C-6), 78.1, 78.5 and 80.9 (C-3,4,5), 72.7, 74.6 and 75.9 (CH<sub>2</sub> Bn), 67.1 (C-2), 60.3 (C-8), 30.4 (C-11), 16.9 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>31</sub>H<sub>34</sub>O<sub>5</sub> 486.2, found *m*/z 509.3 (M + Na)<sup>+</sup>.

5,6,7-Tri-O-benzyl-1,2,3-trideoxy-α-L-rhamno-non-1-eno-4-

**ulopyranose (19).** Prepared from 5,6,7-tri-*O*-benzyl-L-rhamno-1,5-lactone<sup>18</sup> by Grignard addition, according to the general procedure. Colourless syrup (1.49 g, 3.14 mmol, 90%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.51–7.59 (m, CH<sub>arom</sub>), 6.00–6.12 (m, 1H, H-2), 4.88–5.45 (m, 8H, 2 × H-1 and CH<sub>2</sub> Bn), 4.38 (d, 1H, *J* 9.5 Hz, H-6), 3.89–4.20 (m, 3H, H-5,7,8), 3.05 (s, 1H, OH), 2.93 (dd, 1H, *J* 5.1 and 13.9 Hz, H-3), 2.54 (dd, 1H, *J* 9.5 and 13.2 Hz, H-3), 1.56 (d, 3H, *J* 5.8 Hz, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.8, 138.9 (C<sub>q</sub> Bn), 132.7 (C-1), 127.7–128.7 (CH<sub>arom</sub>), 120.6 (C-2), 98.0 (C-4), 68.8, 77.7, 80.7 and 81.9 (C-5,6,7,8), 72.7, 74.8 and 75.3 (CH<sub>2</sub> Bn), 42.8 (C-3), 18.3 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>30</sub>H<sub>34</sub>O<sub>5</sub> 474.2, found *m*/*z* 497.3 (M + Na)<sup>+</sup>.

Allyl (5,6,7-tri-*O*-benzyl-1,2,3-trideoxy)-α-L-*rhamno*-non-1eno-4-ulopyranose (20). Prepared from 19 by K-10-mediated glycosidation according to the general procedure. Colourless syrup (1.43 g, 2.78 mmol, 89%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.50–7.65 (m, 15H, CH<sub>arom</sub>), 5.83–6.20 (m, 2H, H-2 and CH allyl), 5.49 (d, 1H, *J* 16.8 Hz, H-1), 5.17 (d, 1H, CH<sub>2</sub> Bn), 4.90– 5.39 (m, 7H, H-1, CH<sub>2</sub> allyl and Bn), 4.79 (d, 1H, CH<sub>2</sub> Bn), 4.39 (d, 1H, *J* 8.8 Hz, H-6), 4.20 (br s, 2H, 2 × H-10), 4.015 (d, 1H, *J* 2.9 Hz, H-5), 3.83–4.00 (m, 2H, H-7,8), 3.07 (dd, 1H, *J* 7.3 and 15.3 Hz, H-3), 2.76 (dd, 1H, *J* 5.8 and 15.3 Hz, H-3), 1.57 (d, 3H, *J* 5.1 Hz, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.7, 138.8 and 139.0 (C<sub>q</sub> Bn), 134.4 (CH<sub>2</sub> allyl), 132.7 (C-1), 127.6–128.5 (CH<sub>arom</sub>), 118.6 (C-2), 116.3 (CH allyl), 102.4 (C-4), 69.4, 76.4, 80.4 and 82.0 (C-5,6,7,8), 72.6, 74.9, 75.1 and 75.4 (CH<sub>2</sub> Bn), 61.0 (CH<sub>2</sub> allyl), 36.4 (C-3), 18.2 (CH<sub>3</sub>); MS (ESI): calc. for  $C_{33}H_{38}O_5$  514.3, found *m*/*z* 537.4 (M + Na)<sup>+</sup>.

(2*S*,3*S*,4*R*,5*R*,6*R*)-3,4,5-Tris-benzyloxy-2-methyl-1,7-dioxaspiro[5.5]undec-9-ene (21). Obtained from 20 by RCM according to the general procedure (method B) as a greenish syrup (1.35 g, 2.78 mmol, 85%);  $[a]_D - 27.2$  (*c* 1 in CHCl<sub>3</sub>);  $\delta_H$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.31–7.43 (m, 15H, CH<sub>arom</sub>), 5.68 (s, 2H, H-9,10), 4.66. 4.68, 4.80, 4.94 and 5.02 (5d, 6H, CH<sub>2</sub> Bn), 4.00– 4.12 (m, 3H, H-2,3,4), 3.79 (d, 1H, *J* 2.2 Hz, H-5), 3.66 (dd, 2H, *J* 7.3 and 16.1 Hz, H-8), 2.25 (m, 1H, H-11), 2.01 (m, 1H, H-11), 1.35 (d, 3H, *J* 5.0 Hz, CH<sub>3</sub>);  $\delta_C$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.3, 137.6 and 137.8 (C<sub>q</sub> Bn), 126.9–130.0 (CH<sub>arom</sub>), 121.6 and 123.9 (C-9,10), 97.4 (C-6), 78.7, 80.3 and 81.5 (C-3,4,5), 72.6, 74.9 and 75.2 (CH<sub>2</sub> Bn), 68.9 (C-2), 59.6 (C-8), 32.2 (C-11), 17.9 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>31</sub>H<sub>34</sub>O<sub>5</sub> 486.2, found *m*/*z* 509.3 (M + Na)<sup>+</sup>.

**1,2-Dideoxy-4,5;7,8-di-***O***-isopropylidene-α-D***-manno***-oct-1-eno-3-ulofuranose (22).** Prepared from 2,3;5,6-di-*O***-**isopropylidene-D-manno-1,4-lactone<sup>19</sup> by Grignard addition, according to the general procedure. Colourless syrup (1.34 g, 4.7 mmol, 45%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.89 (dd, 1H, *J* 10.9 and 17.5 Hz, H-2), 5.41 (dd, 1H, *J* 1.5 and 17.5 Hz, H-1), 5.20 (dd, 1H, *J* 1.5 and 10.9 Hz, H-1), 4.72 (dd, 1H, *J* 2.9 and 5.1 Hz, H-5), 4.34 (d, 1H, *J* 5.8 Hz, H-4), 4.25 (m, 1H, H-7), 4.02 (dd, 1H, *J* 3.6 and 7.3 Hz, H-8), 3.86–3.97 (m, 3H, H-6,8 and OH), 1.17, 1.23, 1.31 and 1.32 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 136.2 (C-2), 117.0 (C-1), 112.4 (C-3), 104.1 and 108.7 (C<sub>q</sub>), 73.1, 78.9, 80.1 and 86.5 (C-4,5,6,7), 66.2 (C-8), 24.1, 25.0, 25.6 and 26.5 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>14</sub>H<sub>22</sub>O<sub>6</sub> 286.1, found *m*/z 309.1 (M + Na)<sup>+</sup>.

Allyl (1,2-dideoxy-4,5;7,8-di-*O*-isopropylidene)-α-D-mannooct-1-eno-3-ulofuranose (23). Prepared from 22 by K-10mediated glycosidation according to the general procedure. Colourless syrup (0.73 g, 2.24 mmol, 66%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.52–5.72 (m, 2H, H-2 and CH allyl), 5.31 (dd, 1H, J.2.2 and 21.9 Hz, H-1), 5.16 (dd, 1H, J 2.2 and 16.1 Hz, H-1), 4.98 (br t, 2H, CH<sub>2</sub> allyl), 4.67 (dd, 1H, J 3.6 and 5.8 Hz, H-5), 4.37 (d, 1H, J 5.8 Hz, H-4), 4.25 (dd, 1H, J 6.6 and 13.1 Hz, H-8), 3.82–3.99 (m, 3H, H-6,7,8), 3.64 (m, 2H, CH<sub>2</sub> allyl), 1.14, 1.21, 1.26 and 1.28 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 134.4 (CH allyl), 132.6 (C-2), 118.8 (C-1), 115.9 (CH<sub>2</sub> allyl), 112.3 (C-3), 107.8 and 108.7 (C<sub>q</sub>), 72.9, 79.1, 79.9 and 86.3 (C-4,5,6,7), 66.6 (C-8), 62.4 (CH<sub>2</sub> allyl), 24.3, 25.0, 25.6 and 26.5 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>17</sub>H<sub>26</sub>O<sub>6</sub> 326.2, found *m*/z 349.0 (M + Na)<sup>+</sup>.

# (2*R*,3*S*,4*S*,5*S*)-3,4-Dihydroxy-3,4-*O*-isopropylidene-2-({(*R*)-1,2-dihydroxy-1,2-*O*-isopropylidene}ethyl)-1,6-dioxa-spiro-

[4.4]non-8-ene (24). Obtained from 23 by RCM according to the general procedure (method B) as a greenish syrup (0.63 g, 2.11 mmol, 95%);  $[a]_{\rm D}$  –10.4 (*c* 1 in CHCl<sub>3</sub>);  $\delta_{\rm H}$  (600 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 6.17 (d, 1H, *J* 6.0 Hz, H-9), 5.73 (m, 1H, H-8), 4.76 (dd, 1H, *J* 4.1 and 5.4 Hz, H-3), 4.58 (d, 1H, *J* 14.0 Hz, H-7), 4.48 (d, 1H, *J* 13.9 Hz, H-7), 4.46 (d, 1H, *J* 5.9 Hz, H-4), 4.23 (m, 1H, CH Et), 3.95 (t, 1H, *J* 8.7 Hz, CH<sub>2</sub> Et), 3.86–3.90 (m, 2H, H-2 and CH<sub>2</sub> Et), 1.21, 1.24, 1.32 and 1.38 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 132.6 (C-9), 124.5 (C-8), 119.2 (C-5), 108.8 and 112.4 (C<sub>q</sub>), 72.9, 79.9 and 85.7 (C-2,3,4 and CH Et), 74.6 (C-7), 66.8 (CH<sub>2</sub> Et), 24.3, 25.0, 25.7 and 26.7 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>15</sub>H<sub>22</sub>O<sub>6</sub> 298.1, found *m*/z 321.0 (M + Na)<sup>+</sup>, 337.2 (M + K)<sup>+</sup>.

**1,2,3-Trideoxy-5,6;8,9-di-***O***-isopropylidene-α-D***-manno***-non-1-eno-4-ulofuranose (25).** Prepared from 2,3;5,6-di-*O*-isopropyl-idene-D-manno-1,4-lactone<sup>19</sup> by Grignard addition, according

to the general procedure. Colourless syrup (1.65 g, 5.50 mmol, 95%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.75–5.96 (m, 1H, H-2), 5.23 (dd, 2H, *J* 10.2 and 17.5 Hz, 2 × H-1), 4.84 (dd, 1H, *J* 3.7 and 4.8 Hz, H-6), 4.49 (d, 1H, *J* 6.6 Hz, H-5), 4.37 (m, 2H, H-8,9), 3.97–4.20 (m, 3H, H-7,9 and OH), 2.52 (d, 2H, *J* 6.6 Hz, 2 × H-3), 1.34, 1.38, 1.45 and 1.49 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 123.4 (C-2), 119.0 (C-1), 112.2 (C-4), 105.1 and 108.7 (C<sub>q</sub>), 73.0, 78.6, 79.9 and 84.8 (C-5,6,7,8), 66.3 (C-9), 39.6 (C-3), 24.1, 24.9, 25.5 and 26.5 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>15</sub>H<sub>24</sub>O<sub>6</sub> 300.2, found *m*/*z* 323.2 (M + Na)<sup>+</sup>.

Allyl (1,2,3-trideoxy-5,6;8,9-di-O-isopropylidene)-a-D-mannonon-1-eno-4-ulofuranose (26). Prepared from 25 by K-10mediated glycosidation according to the general procedure. Colourless syrup (0.60 g, 1.76 mmol, 54%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.64–5.84 (m, 2H, H-2 and CH allyl), 5.19 (m, 1H, H-1), 4.97-5.10 (m, 3H, H-1 and CH<sub>2</sub> allyl), 4.71 (dd, 1H, J 4.0 and 5.8 Hz, H-6), 4.38 (d, 1H, J 5.8 Hz, H-5), 4.24-4.30 (m, 1H, H-9), 3.94–4.03 (dd, 1H, J 6.2 and 8.4 Hz, H-8), 3.79– 3.90 (m, 3H, H-9 and CH<sub>2</sub> allyl), 3.72 (dd, 1H, J 3.7 and 7.7 Hz, H-7), 2.51 (d, 2H, J 7.7 Hz, 2 × H-3), 1.23, 1.27, 1.34 and 1.38 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 134.3 (CH allyl), 132.1 (C-2), 117.6 (C-1), 116.1 (CH<sub>2</sub> allyl), 112.2 (C-4), 108.7 and 108.9 (Cq), 72.8, 79.2, 79.8 and 84.6 (C-5,6,7,8), 66.7 (C-9), 61.3 (CH<sub>2</sub> allyl), 34.2 (C-3), 24.3, 25.0, 25.7 and 26.6 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>18</sub>H<sub>28</sub>O<sub>6</sub> 340.2, found *m/z* 363.2  $(M + Na)^{+}$ .

(2*R*,3*S*,4*S*,5*S*)-3,4-Dihydroxy-3,4-*O*-isopropylidene-2-({(*R*)-1,2-dihydroxy-1,2-*O*-isopropylidene}ethyl)-1,6-dioxa-spiro[4.5]-dec-8-ene (27). Obtained from 26 by RCM according to the general procedure (method A) as a colourless syrup (0.48 g, 1.54 mmol, 87%);  $[a]_D$ +58.2 (*c* 1 in CHCl<sub>3</sub>);  $\delta_H$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.72 (br s, 2H, H-8,9), 4.82 (dd, 1H, *J* 3.6 and 5.8 Hz, H-3), 4.43 (d, 1H, *J* 5.8 Hz, H-4), 4.35–4.40 (m, 1H, CH<sub>2</sub> Et), 3.95–4.18 (m, 4H, 2 × H-7 and CH and CH<sub>2</sub> Et), 3.85 (dd, 1H, *J* 3.7 and 8.0 Hz, H-2), 2.50 (dd, 1H, *J* 1.5 and 19 Hz, H-10), 2.12 (dd, 1H, *J* 2.2 and 17.5 Hz, H-10), 1.33, 1.36, 1.44 and 1.46 (4s, each 3H, CH<sub>3</sub>);  $\delta_C$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 124.2 (C-8), 121.6 (C-9), 108.9 and 112.1 (C<sub>q</sub>), 104.6 (C-5), 72.8, 79.3, 79.8 and 85.5 (C-2,3,4 and CH Et), 67.0 (CH<sub>2</sub> Et), 59.8 (C-7), 28.6 (C-10), 24.4, 24.9, 25.7 and 26.6 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>16</sub>H<sub>24</sub>O<sub>6</sub> 312.2, found *m*/z 335.2 (M + Na)<sup>+</sup>.

3-Buten-1yl (1,2,3-trideoxy-5,6;8,9-di-O-isopropylidene)-α-Dmanno-non-1-eno-4-ulofuranose (28). Prepared from 25 by K-10-mediated glycosidation according to the general procedure. Colourless syrup (0.30 g, 0.85 mmol, 42%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.68-5.81 (m, 2H, H-2 and CH Bu), 5.00-5.19 (m, 4H, 2 × H-1 and CH<sub>2</sub> Bu), 4.79 (dd, 1H, J 4.4 and 5.8 Hz, H-6), 4.43 (d, 1H, J 5.8 Hz, H-5), 4.32–4.39 (m, 1H, H-8), 4.10 (dd, 1H, J 6.6 and 8.8 Hz, H-9), 3.97 (dd, 1H, J 4.4 and 8.8 Hz, H-9), 3.80 (dd, 1H, J 3.7 and 8.0 Hz, H-7), 3.47 (br t, 2H, J 6.6 Hz, CH<sub>2</sub> Bu), 2.56–2.61 (m, 2H, 2 × H-3), 2.50 (dd, 2H, J 6.6 and 13.1 Hz, CH<sub>2</sub> Bu), 1.32, 1.37, 1.45 and 1.47 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 135.3 (CH Bu), 132.3 (C-2), 117.6 (C-1), 116.4 (CH<sub>2</sub> Bu), 112.3 (C-4), 108.7 and 109.0 (C<sub>a</sub>), 72.9, 79.0, 79.9 and 84.6 (C-5,6,7,8), 66.9 (C-9), 59.6 (CH<sub>2</sub> Bu), 34.1 (C-3), 34.0 (CH<sub>2</sub> Bu), 24.4, 25.1, 25.8 and 26.7 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>19</sub>H<sub>30</sub>O<sub>6</sub> 354.2, found *m/z* 377.2  $(M + Na)^{+}$ .

(2*R*,3*S*,4*S*,5*S*)-3,4-Dihydroxy-3,4-*O*-isopropylidene-2-({(*R*)-1,2-dihydroxy-1,2-*O*-isopropylidene}ethyl)-1,6-dioxa-spiro[4.6]undec-9-ene (29). Obtained from 28 by RCM according to the general procedure (method B) as a greenish syrup (0.27 g, 0.83 mmol, 99%);  $[a]_D$ +9.2 (*c* 1 in CHCl<sub>3</sub>);  $\delta_H$  (600 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 5.65 (m, 1H, H-9), 5.53 (m, 1H, H-10), 4.74 (dd, 1H, *J* 3.8 and 5.9 Hz, H-3), 4.36 (d, 1H, *J* 5.9 Hz, H-4), 4.29–4.32 (m, 1H, CH Et), 4.03 (dd, 1H, *J* 6.3 and 8.7 Hz, CH<sub>2</sub> Et), 3.94 (dd, 1H, *J* 4.3 and 8.7 Hz, CH<sub>2</sub> Et), 3.89 (dd, 1H, *J* 2.2 and 12.0 Hz, H-7), 3.75 (dd, 1H, *J* 3.8 and 8.2 Hz, H-2), 3.55 (dd, 1H, *J* 3.7 and 12.3 Hz, H-7), 2.66 (dd, 1H, *J* 6.5 and 16.3 Hz, H-11), 2.46 (dd, 1H, *J* 5.3 and 16.3 Hz, H-11), 2.38–2.43 (m, 1H, H-8), 2.17 (br d, 1H, *J* 17.2 Hz, H-8), 1.26, 1.31, 1.38 and 1.42 (4s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 130.0 (C-10), 124.3 (C-9), 112.3 (C<sub>q</sub>), 108.9 (C-5), 72.9, 78.6, 79.9 and 85.4 (C-2,3,4 and CH Et), 67.1 (CH<sub>2</sub> Et), 60.4 (C-7), 31.8 (C-11), 31.1 (C-8), 24.3, 25.0, 25.7 and 26.7 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>17</sub>H<sub>26</sub>O<sub>6</sub> 326.2, found *m*/z 349.1 (M + Na)<sup>+</sup>.

(1R,3R,4R,5S,6R)-4,5-Bis-benzyloxy-3-hydroxy-3-vinyl-2,7dioxa-bicyclo[4.4.0]dec-9-ene (32). Bicyclic methyl glycoside 30<sup>22</sup> (630 mg, 1.7 mmol) was dissolved in acetic acid (20 ml) and 3 M sulfuric acid (2.4 ml). The mixture was stirred at 80 °C for 6 h, then allowed to cool to room temperature and diluted with ethyl acetate (30 ml). The organic phase was washed with sat. NaHCO<sub>3</sub> until pH 7, dried, filtered and concentrated. The residue was purified by silica gel chromatography (15% EtOAclight petroleum ether). The purified lactol was dissolved in dimethylsulfoxide-acetic anhydride 2:1 v/v (6 ml) and stirred for 17 h, when TLC analysis (25% EtOAc-light petroleum ether) revealed complete consumption of the starting material. The mixture was concentrated, dissolved in ethyl acetate and extracted with sat. NaHCO<sub>3</sub>. Organics were dried, filtered and concentrated to leave a brown syrup (300 mg, 0.8 mmol, 50%), which was subjected to a Grignard reaction with vinylmagnesium bromide as described in the general procedure to give 32 as a colourless syrup (117 mg, 0.30 mmol, 21%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.26–7.31 (m, 10H, CH<sub>arom</sub>), 5.89 (dd, 1H, J 10.2 and 17 Hz, CH vinyl), 5.68-5.77 (m, 2H, H-9,10), 5.59 (dd, 1H, J 1.5 and 16.8 Hz, CH<sub>2</sub> vinyl), 5.28 (dd, 1H, J 1.5 and 10.2 Hz, CH<sub>2</sub> vinyl), 4.61, 4.77, 4.82 and 4.98 (4d, each 1H, CH<sub>2</sub> Bn), 4.48 (m, 1H, H-1), 4.25 (br s, 2H, 2 × H-8), 3.89 (dd, 1H, J 8.8 and 9.5 Hz, H-5), 3.43 (d, 1H, J 8.8 Hz, H-4), 3.33 (dd, 1H, J 8.8 and 9.5 Hz, H-6);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.8 (CH vinyl), 127.7–128.7 (C-9 and CH<sub>arom</sub>), 126.1 (C-10), 117.0 (CH<sub>2</sub> vinyl), 96.2 (C-3), 65.1, 78.8, 80.5 and 82.7 (C-1,4,5,6), 74.9 and 75.8 (CH<sub>2</sub> Bn), 66.0 (C-8); MS (ESI): calc. for C<sub>24</sub>H<sub>26</sub>O<sub>5</sub> 394.2, found m/z 417.3 (M + Na)<sup>+</sup>.

(1*R*,3*R*,4*R*,5*S*,6*R*)-3-Allyloxy-4,5-bis-benzyloxy-3-vinyl-2,7dioxa-bicyclo[4.4.0]dec-9-ene (33). Prepared from 32 by K-10mediated glycosidation according to the general procedure. Colourless syrup (65 mg, 0.15 mmol, 50%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.22–7.40 (m, 10H, CH<sub>arom</sub>), 5.73–6.00 (m, 2H, CH vinyl and allyl), 5.52 (dd, 1H, *J* 2.2 and 17.5 Hz, CH<sub>2</sub> allyl), 5.27 (dd, 1H, *J* 1.5 and 16.8 Hz, CH<sub>2</sub> vinyl), 5.26 (dd, 1H, *J* 2.2 and 11.0 Hz, CH<sub>2</sub> allyl), 5.12 (dd, 1H, *J* 1.5 and 10.2 Hz, CH<sub>2</sub> vinyl), 4.62, 4.78, 4.91 and 4.96 (4d, each 1H, CH<sub>2</sub> Bn), 4.26 (br s, 2H, 2 × H-8), 4.18 (m, 1H, H-1), 4.06 (dd, 1H, *J* 8.8 and 9.5 Hz, H-5), 3.94 (m, 2H, CH<sub>2</sub> allyl), 3.35 (d, 1H, *J* 8.8 Hz, H-4), 3.31 (dd, 1H, *J* 8.8 and 9.5 Hz, H-6); MS (ESI): calc. for C<sub>27</sub>H<sub>30</sub>O<sub>5</sub> 434.2, found *m*/z 457.2 (M + Na)<sup>+</sup>.

(1*R*,3*S*,4*R*,5*S*,6*R*)-4,5-Bis-benzyloxy-2,7-dioxa-bicyclo[4.4.0]dec-9-enylspiro-3,2'-[2',5'-dihydrofuran] (34). Obtained from 33 by RCM according to the general procedure (method B) as a greenish syrup (58 mg, 0.14 mmol, 95%);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.20–7.29 (m, 10H, CH<sub>arom</sub>), 6.26 (m, 1H, H-3'), 5.77 (m, 2H, H-9,10), 5.55 (m, 1H, H-4'), 4.77 (br s, 2H, 2 × H-5'), 4.59, 4.80 and 4.98 (3d, 4H, CH<sub>2</sub> Bn), 4.40 (m, 1H, H-1), 4.26 (m, 2H, 2 × H-8), 3.97 (br t, 1H, *J* 9.5 Hz, H-6), 3.61 (d, 1H, *J* 8.8 Hz, H-4), 3.33 (dd, 1H, *J* 8.8 and 9.5 Hz, H-5);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.1 and 138.9 (C<sub>q</sub> Bn), 132.6 (C-3'), 126.1–128.1 (C-4',9,10 and CH<sub>arom</sub>), 112.8 (C-3), 78.9, 80.8 and 81.2 (C-4,5,6), 75.1 and 75.9 (CH<sub>2</sub> Bn), 66.1 (C-8), 66.0 (C-1); MS (ESI): calc. for C<sub>25</sub>H<sub>26</sub>O<sub>5</sub> 406.2, found *m*/*z* 429.1 (M + Na)<sup>+</sup>.

3-O-(4',5',6',8'-Tetra-O-benzyl-1',2'-dideoxy-α-D-gluco-oct-1'-eno-3'-ulopyranosyl)-5,6-dideoxy-1,2-O-isopropylidene-α-Dxylo-hex-5-enofuranoside (36). K-10 mediated glycosidation, as described in the general procedure, of 2 (0.39 g, 0.68 mmol) 5,6-dideoxy-1,2-O-isopropylidene-a-D-xylo-hex-5-enowith furanose<sup>22</sup> (35, 0.76 g, 4.09 mmol) afforded 36 as a colourless syrup (0.23 g, 0.31 mmol, 45%);  $\delta_{\rm H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.18-7.35 (m, 20H, CH<sub>arom</sub>), 6.21 (m, 1H, H-5), 5.95 (dd, 1H, J 11.0 and 17.5 Hz, H-2'), 5.93 (dd, 1H, J 2.2 and 12.4 Hz, H-6), 5.64 (dd, 1H, J 2.2 and 17.5 Hz, H-6), 5.38 (d, 1H, J 2.9 Hz, H-1), 5.31–5.39 (m, 1H, H-1), 5.17 (dd, 1H, J 1.5 and 10.2 Hz, H-1), 4.52-4.91 (m, 10H, H-3,4 and CH<sub>2</sub> Bn), 4.26 (d, 1H, J 2.9 Hz, H-2), 4.07 (t, 1H, J 9.5 Hz, H-5'), 3.94 (m, 1H, H-8'), 3.63-3.82 (m, 2H, H-7',8'), 3.53 (t, 1H, J 9.5 Hz, H-6'), 3.30 (d, 1H, J 9.5 Hz, H-4'), 1.18 and 1.47 (2s, each 3H, CH<sub>3</sub>);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.9 and 138.5 (C<sub>q</sub> Bn), 133.2 and 135.0 (C-2',5), 127.1-128.4 (CH<sub>arom</sub>), 119.5 and 119.7 (C-1',6), 111.2 (Cq), 104.7 (C-1), 100.5 (C-3'), 73.5, 75.2, 75.4 and 75.6 (CH<sub>2</sub> Bn), 72.5, 78.4, 78.6, 82.3, 82.5, 83.6 and 85.1 (C-2,3,4,4',5',6',7'), 69.1 (C-8'), 26.0 and 26.7 (CH<sub>3</sub>); MS (ESI): calc. for C<sub>45</sub>H<sub>50</sub>O<sub>9</sub> 734.3, found *m*/*z* 757.4  $(M + Na)^{+}$ .

2,3,4,6-Tetra-O-benzyl-α-D-glucopyranosylspiro-1'-[(3'S,4'R, 5'R,7'R)-4',5'-dihydroxy-4',5'-O-isopropylidene-2',6'-dioxabicyclo[3.4.0]non-8'-ene] (37). Obtained from 36 by RCM according to the general procedure (method B) as a colourless syrup (85 mg, 0.12 mmol, 64%); [a]<sub>D</sub>+64.4 (c 1 in CHCl<sub>3</sub>); δ<sub>H</sub> (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.12–7.31 (m, 20H, CH<sub>arom</sub>), 6.18 (dd, 1H, J 5.1 and 10.2 Hz, H-9'), 6.08 (d, 1H, J 3.6 Hz, H-1), 5.78 (d, 1H, J 10.2 Hz, H-8'), 4.69 (br s, 2H, CH<sub>2</sub> Bn), 4.47, 4.53, 4.60, 4.81, 4.83 and 4.91 (6d, each 1H, CH<sub>2</sub> Bn), 4.40 (m, 1H, H-5'), 4.29 (d, 1H, J 3.0 Hz, H-4'), 4.05 (t, 1H, J 8.8 Hz, H-3), 3.61-3.88 (m, 4H, H-1,7 and  $2 \times$  H-8), 3.71 (t, 1H, J 8.8 Hz, H-4), 3.51 (d, 1H, J 8.8 Hz, H-2), 1.32 and 1.52 (2s, each 3H, CH<sub>3</sub>); δ<sub>C</sub> (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.9, 138.1 and 138.7 (C<sub>q</sub> Bn), 127.7–128.3 (CH<sub>arom</sub>), 125.4 and 132.2 (C-8',9'), 111.3 (C<sub>a</sub>), 105.7 (C-3'), 95.5 (C-1'), 73.4, 74.7, 74.9 and 75.5 (CH<sub>2</sub> Bn), 69.5, 71.8, 73.7, 77.8, 82.6 and 84.2 (C-1',2,3,4,4',5,5'), 68.3 (C-6), 26.1 and 26.8 (CH<sub>3</sub>); MS (ESI): calc. for  $C_{43}H_{46}O_{9}$  706.3, found *m*/*z* 729.5 (M + Na)<sup>+</sup>.

(2R,3R,4S,5R,6S)-3,4,5-Tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.5]undec-8-ene (38). To a stirred solution of 8 (330 mg, 0.56 mmol) in anhydrous ethanol (30 mL) were added Wilkinson's catalyst (129 mg, 0.13 mmol) and DBU (2 drops) under a nitrogen atmosphere. The mixture was heated to 60 °C for 1 h after which TLC analysis (25% EtOAc-light petroleum ether) revealed completed conversion. The mixture was allowed to cool to room temperature and concentrated. Silica gel chromatography (15% EtOAc-light petroleum ether) of the residue yielded **38** as a colourless syrup (296 mg, 0.50 mmol, 90%);  $[a]_{D}$ +8.4 (c 1 in CHCl<sub>3</sub>);  $\delta_{H}$  (200 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.16-7.26 (m, 20H, CH<sub>arom</sub>), 6.25 (d, 1H, J 6.2 Hz, H-8), 4.50, 4.55, 4.57, 4.60, 4.68 4.86, 4.91 and 4.92 (8d, each 1H, CH<sub>2</sub> Bn), 4.79 (m, 1H, H-9), 4.16 (t, 1H, J 9.1 Hz, H-4), 4.11-4.21 (m, 1H, CH<sub>2</sub>OBn), 3.62–3.79 (m, 3H, H-2,3 and CH<sub>2</sub>OBn), 3.40 (d, 1H, J 9.9 Hz, H-5), 1.49–2.05 (m, 4H,  $2 \times$  H-10 and  $2 \times$  H-11);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si)  $\delta$  139.6 (C-8), 137.7, 138.2 and 138.6 (C<sub>q</sub> Bn), 127.5–128.4 (CH<sub>arom</sub>), 102.3 (C-9), 97.6 (C-6), 71.8, 78.3, 82.6 and 83.0 (C-2,3,4,5), 73.1, 74.8, 75.5 and 75.6 (CH<sub>2</sub> Bn), 68.4 (CH<sub>2</sub>OBn), 26.8 (C-11), 16.0 (C-10); MS (ESI): calc. for  $C_{38}H_{40}O_6$  592.3, found *m*/*z* 615.3 (M + Na)<sup>+</sup>.

# (2R,3R,4S,5R,6S,8R,9S)-8-Allyl-3,4,5-tris-benzyloxy-2-

**benzyloxymethyl-1,7-dioxa-spiro[5.5]-undecan-9-ol (39).** To a chilled (0 °C) solution of spiroketal **38** (370 mg, 0.63 mmol) in anhydrous dichloromethane (3 mL) was added dropwise a freshly prepared solution of 3,3-dimethyldioxirane (9 mL, 0.09 M, 0.8 mmol) in acetone.<sup>25</sup> After stirring the mixture for

10 min, TLC analysis (20% EtOAc-light petroleum ether) showed complete consumption of the starting material. The mixture was concentrated, redissolved in anhydrous toluene (5 mL) and concentrated again. The crude epoxide was dissolved in THF (5 mL), cooled to 0 °C under argon atmosphere, after which a solution of allylmagnesium chloride (0.58 mL, 2 M, 1.16 mmol) was added. After stirring for 15 min., TLC analysis (33% EtOAc/light petroleum ether) showed a nearly completed reaction. The reaction was quenched by addition of sat. NH<sub>4</sub>Cl and extracted with diethyl ether. The organic layer was washed with sat. NH<sub>4</sub>Cl, dried and concentrated. Silica gel chromatography (16 to 40% EtOAc-light petroleum ether) yielded compound 39 as a colourless syrup (246 mg, 0.37 mmol, 61%);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.7, 138.1 and 138.5 (C<sub>a</sub> Bn), 134.8 (CH allyl), 127.4–128.2 (CH<sub>arom</sub>), 117.1 (CH<sub>2</sub> allyl), 98.3 (C-6), 73.2, 74.8, 75.3 and 75.7 (CH<sub>2</sub> Bn), 68.7 (CH<sub>2</sub>OBn), 65.9, 71.1, 77.9, 78.5, 82.9 and 83.7 (C-2,3,4,5,8,9), 37.4 (CH<sub>2</sub> allyl), 25.0 (C-11), 21.8 (C-10); MS (ESI): calc. for C41H46O7 650.3, found m/z 673.3 (M + Na)<sup>+</sup>.

(2R,3R,4S,5R,6S,8R,9S)-8-Allyl-9-allyloxy-3,4,5-tris-benzyloxy-2-benzyloxymethyl-1,7-dioxa-spiro[5.5]undecane (40). To a solution of compound 39 (246 mg, 0.38 mmol) in anhydrous N,N-dimethylformamide (3 mL) were added allyl bromide (50 µl, 0.57 mmol) and sodium hydride (20 mg, 60% dispersion in oil, 0.49 mmol). After stirring for 16 h, TLC analysis (33% EtOAc-light petroleum ether) showed complete disappearance of 39. The excess sodium hydride was destroyed by addition of methanol (5 mL). The mixture was partitioned between diethyl ether and brine. The organic phase was extracted with brine, dried and concentrated. Purification was effected by silica gel chromatography (20% EtOAc-light petroleum ether) to afford 40 as a colourless syrup (242 mg, 0.35 mmol, 93%);  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 138.2 and 138.7 (C<sub>q</sub> Bn), 135.1 (CH allyl), 127.3–128.2 (CH<sub>arom</sub>), 116.1 and 116.9 (CH<sub>2</sub> allyl), 98.6 (C-6), 73.1, 74.7, 75.3 and 75.5 (CH<sub>2</sub> Bn), 70.9, 72.9, 75.4, 78.6, 82.9 and 84.0 (C-2,3,4,5,8,9), 69.1 (CH2 allyl), 68.8 (CH2OBn), 37.6 (CH<sub>2</sub> allyl), 25.5 (C-11), 19.4 (C-10); MS (ESI): calc. for  $C_{44}H_{50}O_7$  690.3, found *m*/*z* 713.3 (M + Na)<sup>+</sup>.

2,3,4,6-Tetra-O-benzyl-α-D-glucopyranosylspiro-3'-[(1'R,

6'S)-2',7'-dioxa-bicyclo[4.5.0]undec-9'-ene] (42). To a solution of diene 40 (240 mg, 0.35 mmol) in anhydrous dichloromethane (5 mL) was added Grubbs' catalyst 41<sup>27</sup> (0.1 mg, 0.18 µmol) under argon atmosphere and the mixture was stirred at 40 °C for 2 h. TLC analysis (33% EtOAc-light petroleum ether) revealed a completed conversion, after which the mixture was concentrated and purified by silica gel chromatography (20%) EtOAc-light petroleum ether) to give a colourless syrup of 42 (185 mg, 0.28 mmol, 80%);  $\delta_{\rm H}$  (300 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 7.16– 7.31 (m, 20H, CH<sub>arom</sub>), 5.76–5.86 (m, 2H, H-9', 10'), 4.88 (br s, 2H, CH<sub>2</sub> Bn), 4.51, 4.54, 4.60, 4.67, 4.84 and 4.89 (6d, each 1H, CH<sub>2</sub> Bn), 4.14 (d, 1H, J 4.9 and 14.7 Hz, H-8'), 4.04 (t, 1H, J 9.2 Hz, H-3), 3.84-4.01 (m, 1H, H-8'), 3.57-3.76 (m, 5H, H-1,4,5 and 2 × H-6), 3.33 (d, 1H, J 9.5 Hz, H-2), 2.61–2.69 (m, 1H, H-11'), 2.11-2.33 (m, 3H, H-4',5',11'), 1.42-1.83 (m, 2H, H-4',5');  $\delta_{\rm C}$  (50 MHz; CDCl<sub>3</sub>; Me<sub>4</sub>Si) 137.9, 138.1, 138.2 and 138.6 (C<sub>q</sub> Bn), 131.7 and 131.9 (C-9',10'), 127.4-128.8 (CH<sub>arom</sub>), 98.6 (C-1), 73.2, 74.9 and 75.4 (CH<sub>2</sub> Bn), 70.7, 73.4, 78.6, 81.0, 83.3 and 83.7 (C-1',2,3,4,5,6'), 68.9 (C-6), 67.1 (C-8'), 34.3 (C-11'), 27.5 (C-4'), 24.8 (C-5'); MS (ESI): calc. for  $C_{42}H_{46}O_7$  662.3, found *m*/*z* 685.3 (M + Na)<sup>+</sup>.

### Acknowledgements

The authors thank Fons Lefeber for recording the nOe-difference spectra, and Hans van den Elst and Nico Meeuwenoord for recording the mass spectra. H.S.O. thanks the Netherlands Organization for Scientific Research (NWO) for financial support.

- For reviews on the synthesis of spiroketals, see: (a) F. Perron and K. F. Albizati, Chem. Rev., 1989, 89, 1617; (b) T. L. B. Boivin, Tetrahedron, 1987, 43, 3309; (c) A. F. Kluge, Heterocycles, 1986, 24, 1699; (d) I. E. Markó, A. Mekhalfia, F. Murphy, D. J. Bayston, M. Bailey, Z. Janousek and S. Dolan, Pure Appl. Chem., 1997, 69, 565.
- 2 S. Hanessian and A. Ugolini, Carbohydr. Res., 1984, 130, 261.
- 3 (a) I. I. Cubero, M. T. P. López-Espinosa and N. Kari, *Carbohydr. Res.*, 1995, **268**, 187; (b) I. I. Cubero, M. T. P. López-Espinosa and N. Kari, *Carbohydr. Res.*, 1994, **261**, 231; (c) I. I. Cubero, M. T. P. López-Espinosa, A. C. Richardson and K. H. Aamlid, *Carbohydr. Res.*, 1993, **242**, 281.
- 4 (a) R. L. Dorta, A. Martín, J. A. Salazar and E. Suárez, J. Org. Chem., 1998, 63, 2251; (b) R. L. Dorta, A. Martín, J. A. Salazar and E. Suárez, *Tetrahedron Lett.*, 1996, 37, 6021; (c) A. Martín, J. A. Salazar and E. Suárez, J. Org. Chem., 1996, 61, 3999; (d) A. Martín, J. A. Salazar and E. Suárez, *Tetrahedron Lett.*, 1995, 36, 4489.
- 5 A. Haudrechy and P. Sinaÿ, Carbohydr. Res., 1991, 216, 375.
- 6 A synthesis of carbohydrate-derived spiroketals, analogous to reference 5, was recently published: G. V. M. Sharma, Rakesh, A. S. Chander, V. G. Reddy, M. V. H. R. Rao and A. C. Kunwar, *Tetrahedron: Asymmetry*, 2003, 14, 2991.
- 7 (a) P. A. V. van Hooft, M. A. Leeuwenburgh, H. S. Overkleeft, G. A. van der Marel, C. A. A. van Boeckel and J. H. van Boom, *Tetrahedron Lett.*, 1998, **39**, 6061; (b) M. A. Leeuwenburgh, C. C. M. Appeldoorn, P. A. V. van Hooft, H. S. Overkleeft, G. A. van der Marel and J. H. van Boom, *Eur. J. Org. Chem.*, 2000, 873.
- 8 B. M. Trost and E. D. Edstrom, Angew. Chem., Int. Ed. Engl., 1990, 29, 520.
- 9 P. Schwab, R. H. Grubbs and J. W. Ziller, J. Am. Chem. Soc., 1996, 118, 100.
- 10 Assignment according to the CIP system: H. Hirschmann and K. R. Hanson, J. Org. Chem., 1971, 36, 3293.
- 11 Previous assignment of absolute configuration at the spirocentre in this compound was incorrect, see references 5 and 10.

- 13 K. Tomooka, Y. Nakamura and T. Nakai, Synlett, 1995, 321.
- 14 M. D. Lewis, J. K. Cha and Y. Kishi, J. Am. Chem. Soc., 1982, 104, 4976.
- 15 The absolute configurations at the spirocentres of all compounds described in this paper were determined by NOESY-experiments.
- 16 M. L. Alcaraz, F. K. Griffin, D. E. Paterson and R. J. K. Taylor, *Tetrahedron Lett.*, 1998, **39**, 8183.
  17 The synthesis of compound **11** via a different approach was recently
- 17 The synthesis of compound 11 via a different approach was recently reported: R. Caputo, U. Ciriello, P. Festa, A. Guaragna, G. Palumbo and S. Pedatella, *Eur. J. Org. Chem.*, 2003, 2617.
- 18 S. Cai, M. R. Stroud, S. Hakomori and T. Toyokuni, J. Org. Chem., 1992, 57, 6693.
- 19 A. J. Norris and T. Toyokuni, J. Carbohydr. Chem., 1999, 18, 1097.
- 20 R. S. Tipson, H. S. Isbell and J. E. Stewart, J. Res. Natl. Bur. Stand., 1959, 62, 257.
- 21 Similar bis-adducts have been isolated after Grignard reactions with γ-valerolactone and δ-butyrolactone; see: (a) H. Ogura, H. Takahashi and Itoh, J. Org. Chem., 1972, **37**, 72; (b) M. Sprinzl, J. Farkas and F. Šorm, *Tetrahedron Lett.*, 1966, 597; (c) M. S. Kharash in *Grignard Reactions of Non-Metallic Substances*, Prentice Hall, Englewood Cliffs, New York, 1954, and references cited herein.
- 22 (a) M. A. Leeuwenburgh, H. S. Overkleeft, G. A. van der Marel and J. H. van Boom, *Synlett*, 1997, 1263; (b) M. A. Leeuwenburgh, C. Kulker, H. I. Duynstee, H. S. Overkleeft, G. A. van der Marel and J. H. van Boom, *Tetrahedron*, 1999, **55**, 8253–8262.
- 23 G. V. M. Sharma and K. Krishnudu, *Tetrahedron Lett.*, 1995, 36, 2661.
- 24 (a) E. J. Corey and J. W. Suggs, J. Org. Chem., 1973, 38, 3224; (b) J. S. Clark and J. G. Kettle, *Tetrahedron Lett.*, 1997, 38, 127.
- 25 W. Adam, J. Bialas and L. Hadjiarapoglou, Chem. Ber., 1991, 2377.
- 26 (a) W. M. Best, V. Ferro, J. Harle, R. V. Stick and D. M. G. Tilbrook, *Aust. J. Chem.*, 1997, **50**, 463; (b) D. A. Evans, B. W. Trotter and B. Côté, *Tetrahedron Lett.*, 1998, **39**, 1709.
- 27 M. Scholl, S. Ding, C. W. Ling and R. H. Grubbs, *Org. Lett.*, 2000, **2**, 953.
- 28 S. P. Allwein, J. M. Cox, B. E. Howard, H. W. B. Johnson and J. D. Rainier, *Tetrahedron*, 2002, 58, 1997.