

Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine: $[N(PPh_2)_2]_2Ni$, $(Ph_3P)_2Ni[(Ph_2P)_2NH]$. Synthesis, characterization, and some properties

Vyacheslav V. Sushev^a, Alexander N. Kornev^{a,*}, Yana V. Fedotova^a,
Yurii A. Kursky^a, Tatiana G. Mushtina^a, Gleb A. Abakumov^a, Lev N. Zakharov^b,
Arnold L. Rheingold^b

^a Razuvaev Institute of Organometallic Chemistry, Russian Academy of Sciences, 49 Tropinin Street, 603950 Nizhny Novgorod, Russia

^b Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

Received 14 March 2003; received in revised form 2 April 2003; accepted 17 April 2003

Abstract

Disproportionation of nickel(I) bis(triphenylphosphino)bis(trimethylsilyl)amide, $(Ph_3P)_2Ni-N(SiMe_3)_2$, in the presence of bis(diphenylphosphino)amine, $(Ph_2P)_2NH$, yields Ni(II) and Ni(0) phosphinoamide complexes: $[N(Ph_2P)_2]_2Ni$ (**1**), $(Ph_3P)_2Ni[(Ph_2P)_2NH]$ (**2**). Ether solution, containing **2** and Ph_3P (1:2) reacts with dioxygen (one equivalent) to form triphenylphosphinoxide adduct $(Ph_3P)_2Ni[(Ph_2P)_2NH \cdots OPPh_3]$ (**3**) in high yield. The crystal structures of compounds **1** and **3** have been determined by X-ray diffraction method.


© 2003 Elsevier Science B.V. All rights reserved.

Keywords: Transition metals; Nickel; Phosphazanes; Phosphinoamides; X-ray diffraction


1. Introduction

bis(Diphenylphosphino)amine (dppa) is one of the most used and intriguing phosphorus–nitrogen ligands in transition metal chemistry [1]. It demonstrates coordinative versatility both in its neutral form and as an anion $[Ph_2PNPPh_2]^-$ [1,2]. However, one of the problems arising is the easy implication of dppa into redox processes. Particularly, it is specific to the bis(diphenylphosphino)amide anion $[Ph_2PNPPh_2]^-$. So, lithium salt of dppa reacts with some inorganic halides to form unpredictable products of phosphazene type. This is the situation, for example, in the reaction of $LiN(PPh_2)_2$ with PCl_3 , leading to the mixture of cyclic $\{N(Ph_2P=P-PPh_2)_2N\}$, and linear $\{(Ph_2P-N=PPh_2)_2\}$ phosphazenes [3]. Reactions of nickel and cobalt chlorides with $LiN(PPh_2)_2$ represent a combina-

tion of redox and substitution processes. Prolonged heating of $LiN(Ph_2P)_2$ with cobalt(II) or nickel(II) chlorides in toluene yields phosphazene complexes $[(Ph_2P-N)_2PPh_2]_2M$ ($M = Co$ [4], Ni [5]):



It remains unclear, however, whether homoleptic complexes $[N(Ph_2P)_2]_2M$ were formed in the beginning of the reaction or not. The same reaction ($M = Ni$) in the presence of trimethylphosphine ligands yields dimer **4**.



Further substitution of the remaining chlorine atom for dppa is difficult apparently due to considerable steric

* Corresponding author.

Table 1
Summary of crystal and refinement data for complexes



	[N(PPh ₂) ₂] ₂ Ni (1)	(Ph ₃ P) ₂ Ni[(Ph ₂ P) ₂ NH⋯OPPh ₃] (3)
Empirical formula	C ₄₈ H ₄₀ N ₂ NiP ₄	C ₇₈ H ₆₆ NNiOP ₅
Formula weight	827.41	1246.88
Temperature (K)	150(2)	150(2)
Wavelength (Å)	0.71073	0.71073
Crystal system	monoclinic	triclinic
Space group	<i>I</i> 2/ <i>a</i>	<i>P</i> $\bar{1}$
Unit cell dimensions		
<i>a</i> (Å)	17.5853(8)	13.0058(7)
<i>b</i> (Å)	12.3427(6)	13.3297(8)
<i>c</i> (Å)	18.5459(9)	19.5573(11)
α (°)	90	76.8900(10)
β (°)	96.5050(10)	80.1530(10)
γ (°)	90	78.8340(10)
<i>V</i> (Å ³)	3999.5(3)	3210.9(3)
<i>Z</i>	4	2
<i>D</i> _{calc} (Mg m ^{−3})	1.374	1.290
Absorption coefficient (mm ^{−1})	0.683	0.474
Crystal size (mm ³)	0.25 × 0.20 × 0.10	0.35 × 0.30 × 0.25
Reflections collected	12152	20623
Independent reflections	4590	14378
	[<i>R</i> _{int} = 0.0169]	[<i>R</i> _{int} = 0.0190]
Absorption correction	SADABS	SADABS
Max. and min. transmission	1.000; 0.898	1.000; 0.837
Refinement method	Full-matrix least-squares on <i>F</i> ²	Full-matrix least-squares on <i>F</i> ²
Data/restraints/parameters	4590/0/395	14378/0/779
Final <i>R</i> indices	<i>R</i> = 0.1260 ^a , <i>wR</i> = 0.3394 ^b	<i>R</i> = 0.0438 ^a , <i>wR</i> = 0.1053 ^b
[<i>I</i> > 2σ(<i>I</i>)]		
<i>R</i> indices (all data)	<i>R</i> = 0.1269 ^a , <i>wR</i> = 0.3397 ^b	<i>R</i> ₁ = 0.0559 ^a , <i>wR</i> = 0.1122 ^b
<i>S</i> ^c	1.154	1.014
Largest difference peak and hole (e Å ^{−3})	0.890; −1.630	0.618; −0.608

^a *R* = Σ||*F*_o| − |*F*_c||/Σ|*F*_o|.

^b *wR* = *R*(*wF*²) = {Σ[*w*(*F*_o² − *F*_c²)]/Σ[*w*(*F*_o²)]}^{1/2}; *w* = 1/[σ²(*F*_o²) + (*aP*)² + *bP*], *P* = [2*F*_c² + max(*F*_o, 0)]/3.

^c *S* = Goodness-of-fit on *F*² = {Σ[*w*(*F*_o² − *F*_c²)]/(*n* − *p*)^{1/2}, where *n* is the number of reflections, and *p* is the number of refined parameters.

ing **2** and triphenylphosphine (reaction mixture after separation of the complex **1**). To our surprise we succeeded to separate not a free oxidized ligands but in the complex with **2**. The compound **3** contained Ph₃P=O molecule associated with hydrogen atom of dppa ligand by hydrogen bond. Since the Ph₃P=O molecule was not coordinate to a nickel atom, the oxidation process may conditionally consider as catalytic one.



Addition of an excess of dioxygen to the reaction

mixture at room temperature instantly affords a lot of colorless crystals of pure triphenylphosphine oxide. No other free phosphine oxides (particularly Ph₂PNHP(O)Ph₂, possible products of dppa oxidation) were detected.

2.2. Structures

The structures of **1** and **3** have been determined by X-ray diffraction methods. The crystal data and some details of the data collection and refinement for **1** and **3** are given in Table 1. Selected bond distances and angles for **1** and **3** are in Tables 2 and 3, respectively.

The Ni atom in **1** is on a center of symmetry; i.e. the NP₂NiP₂N fragment is planar (Fig. 1). In **3** the Ni atom has a distorted tetrahedral environment (Fig. 2). The N–P bond lengths in a free (Ph₂P)₂NH molecule are 1.692(2) Å [7]. The close distances [1.6916(18) and 1.6935(18) Å] were observed in Ni(0) complex **3**. So the chelation of dppa to the nickel atom in **3** does not change the length and the order of N–P bond. At the same time the valence PNP angle is contracted significantly from 118.9(2)° in the free ligand [7] to 101.97(10)° in the Ni(0) complex **3**. The average Ni–P distance of 2.22 Å in **1** is within the typical range observed for Ni(II) complexes and slightly longer than those (2.1926(6) and 2.1987(5) Å) founded in Ni(0) complex **3**.

In going from **1** to **3** we observed the shortening of the P–N bonds from 1.6935(18) and 1.6916(18) Å to 1.653(7) and 1.656(9) Å, respectively. This can be rationalized by a higher bond order of the N–P bonds when the metal center is more electropositive, as electron density is transferred from the metal to appropriate N–P π-bonding orbitals. It causes in turn further decreasing of PNP and chelating PNiP angles to 96.5(4) and 67.43(8)°, respectively. To the best of our knowledge, the chelating PNiP angle 67.43(8)° in **1** is the smallest known. For example, in the similar compound [CH₂(PPh₂)₂]₂Ni[BF₄]₂ [6] chelating PNiP angle is significantly larger, 73.2(3)°. Interestingly, the complex **1** is paramagnetic at room temperature with magnetic moment 3.1 μ_B, while the other planar phosphine complexes are diamagnetic [6]. Perhaps, this phenomenon is associated with temperature dependent change in geo-

Table 2
Selected bond lengths (Å) and angles (°) for [N(PPh₂)₂]₂Ni (**1**)


Ni(1)–P(1)	2.226(2)	P(2) ^a –Ni(1)–P(1)	112.57(8)
Ni(1)–P(2)	2.222(2)	P(2)–Ni(1)–P(1)	67.43(8)
P(1)–N(1)	1.653(7)	N(1)–P(1)–Ni(1)	98.0(3)
P(1)–C(1)	1.822(9)	N(1)–P(2)–Ni(1)	98.0(3)
P(1)–C(7)	1.832(9)	P(1)–N(1)–P(2)	96.5(4)
P(1)⋯P(2)	2.469(3)		
P(2)–N(1)	1.656(9)		

^a Symmetry equivalent atom.

Table 3

Selected bond lengths (Å) and angles (°) for $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH} \cdots \text{OPPh}_3]$ (**3**)

Ni(1)–P(1)	2.1736(6)	P(1)–Ni(1)–P(2)	113.77(2)
Ni(1)–P(2)	2.1741(6)	P(1)–Ni(1)–P(3)	112.92(2)
Ni(1)–P(3)	2.1926(6)	P(2)–Ni(1)–P(3)	118.35(2)
Ni(1)–P(4)	2.1987(5)	P(1)–Ni(1)–P(4)	116.25(2)
		P(2)–Ni(1)–P(4)	116.26(2)
P(3)–N(1)	1.6935(18)	P(3)–Ni(1)–P(4)	73.59(2)
P(3)···P(4)	2.6301(7)	N(1)–P(4)–Ni(1)	92.13(6)
P(4)–N(1)	1.6916(18)	N(1)–P(3)–Ni(1)	92.29(6)
P(5)–O(1)	1.444(2)	P(4)–N(1)–P(3)	101.97(10)
P(5)–C(73)	1.808(2)	P(4)–N(1)–H(1N)	128.8(16)
N(1)–H(1N)	0.83(2)	P(3)–N(1)–H(1N)	128.9(16)
O(1)···H(1N)	1.99(2)	N(1)–H(1N)···O(1)	170(2)
N(1)···O(1)	2.820(3)		


Fig. 1. An ORTEP view of **1** with 30% probability ellipsoids. Only one position of the disordered Ph-rings is shown.

metry of **1** from tetrahedral (at room temperature) to plane-square at 150 K. This study now is in progress.

3. Experimental

3.1. General considerations

Solvents were purified following standard methods [8]. Toluene was thoroughly dried and distilled over P_2O_5 prior to use. Ether was dried and distilled over Na/benzophenone; the compounds $(\text{Ph}_3\text{P})_2\text{Ni}-\text{N}(\text{SiMe}_3)_2$ [9], $(\text{Ph}_2\text{P})_2\text{NH}$ [10], were prepared according to known methods. All manipulations were performed with rigorous exclusion of oxygen and moisture, in vacuum or

Fig. 2. An ORTEP view of **3** with 30% probability ellipsoids.

under an argon atmosphere using standard Schlenk techniques.

Spectrophotometric determination of nickel with dimethylglyoxime was provided by the method [11];

Infrared spectra were recorded on a Perkin–Elmer 577 spectrometer from 4000 to 400 cm^{-1} in nujol. Room-temperature magnetic moments were measured by the Faraday method.

NMR spectra were recorded in CDCl_3 or C_6D_6 solutions using ‘Bruker DPX-200’ instrument, with Me_4Si as internal standard.

3.2. Reaction of $(\text{Ph}_3\text{P})\text{Ni}-\text{N}(\text{SiMe}_3)_2$ with bis(diphenylphosphino)amide

A mixture of toluene solutions of $(\text{Ph}_2\text{P})_2\text{NH}$ (0.29 g, 0.75 mmol, 5 ml) and $(\text{Ph}_3\text{P})_2\text{Ni}-\text{N}(\text{SiMe}_3)_2$ (0.37 g, 0.50 mmol, 5 ml) was maintained at 20°C . The yellow mixture turned dark-red. Red crystalline precipitate of **1** was formed after 3 h. It was filtered, washed with toluene and dried in vacuum. Yield 0.15 g (73%). Anal. Calc. for $\text{C}_{48}\text{H}_{40}\text{P}_4\text{N}_2\text{Ni}$ (**1**): C, 69.67, H, 4.87, Ni, 7.09. Found: C, 70.00; H, 5.04, Ni, 6.93%. IR (cm^{-1}): 1430 s, 1300 w, 1220 w, 1180 vw, 1100 s, 1020 w, 1000 w, 920 vs, 870 s, 730 s, 700 s, 550 s, 510 s, 490 s. $\mu_{\text{eff}} = 3.1\ \mu_{\text{B}}$.

Toluene was pumped in vacuum from filtrate and changed for ether (8 ml). Dark-brown crystals of **2** formed overnight were filtered, washed with cold ether and dried in vacuum. Yield 0.18 g (75%) of $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$. Anal. Calc. For $\text{C}_{60}\text{H}_{51}\text{P}_4\text{NNi}$: C, 74.40; H, 5.31; Ni, 6.06. Found: C, 74.56; H, 5.43; Ni, 5.93%. IR (cm^{-1}): 3050 w, 1580 w, 1430 m, 1300 m, 1200 m, 1180 m, 1120 m, 1090 s, 1070 w, 1020 m, 790 m, 730 s, 700 vs, 550 m, 510 vs. ^{31}P -NMR (C_6D_6), δ ppm: 61.9 (t, $^2J_{\text{PP}}$ 20.3, Ph_2P), 33.5 (t, $^2J_{\text{PP}}$ 20.3, Ph_3P), ^1H -NMR: 2.2 (s, 1H, NH), 6.3–8.1 (m, 50H).

3.3. Oxidation of **2** in the presence of Ph_3P

The reaction of $(\text{Ph}_2\text{P})_2\text{NH}$ (0.75 mmol) with $(\text{Ph}_3\text{P})_2\text{Ni}-\text{N}(\text{SiMe}_3)_2$ (0.50 mmol) was carried out as described above. The toluene filtrate after separation of **1** contained 0.25 mmol of **2** and 0.5 mmol of Ph_3P . Toluene was changed for ether. Dioxygen (2.8 ml, 0.125 mmol) was allowed to react with solution at 20 °C. Large brown–red crystals were grown overnight. The crystals were separated, washed with cold ether and dried in vacuum. Yield 0.24 g (77%) of **3**. Anal. Calc. For $\text{C}_{78}\text{H}_{66}\text{P}_5\text{NONi}$: C, 75.13; H, 5.34; Ni, 4.71. Found: C, 74.96; H, 5.40; Ni, 4.75%. IR (cm^{-1}): 1430 m, 1170 s ($\text{P}=\text{O}$), 1120 m, 1080 m, 1020 w, 970 w, 820 m, 740 w, 720 m, 700 s, 540 s, 510 s. ^{31}P -NMR (C_6D_6), δ ppm: 61.9 (t, $^2J_{\text{PP}}$ 19.7 Hz, Ph_2P), 33.6 (t, $^2J_{\text{PP}}$ 19.7 Hz, Ph_3P), 27.3 (s, Ph_3PO). ^1H -NMR: 2.3 (s, 1H, NH), 6.3–8.1 (m, 65H).

3.4. X-ray diffraction studies

X-ray data were collected on a Bruker Smart Apex CCD diffractometer at 150(2) K. The crystal data and some details of the data collection and refinement for **1** and **3** are given in Table 1. Both structures were determined using a combination of direct methods and calculations of Fourier maps and refined by full-matrix least-squared procedures based on the structural factors F^2 . The positions of the H atoms were calculated using general geometrical conditions and the H atoms were refined in a rigid group model. Selected bond distances and angles in **1** and **3** are given in Tables 2 and 3, respectively. In the crystal structure of **1** there are two possible arrangements of the main molecule with different positions of Ph-rings. The disordered positions of Ph-rings corresponding to two such arrangements were found from the difference F-map and refined without additional restrictions for Ph-rings. Occupation multiplicities of these two arrangements were refined and gave ratio 49/51. The high value of R -factor for this structure seems to be related with existence of such disorder.

SADABS absorption corrections were applied to both structures [12]. All software and sources scattering factors are contained in the SHELXTL (5.10) program package [13].

4. Conclusions

Nickel (I) bis(triphenylphosphino)bis(trimethylsilyl)amide turned out to be a convenient reagent in prepara-

tion of novel dppa complexes of nickel, $[\text{N}(\text{Ph}_2\text{P})_2]_2\text{Ni}$ (**1**), $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ (**2**). The molecule of bis[bis(diphenylphosphino)amido]nickel (II) (**1**) shows the smallest of known chelating PNiP angle $67.43(8)^\circ$ and possess of magnetic moment $3.1 \mu_{\text{B}}$ at 20 °C. The partial oxidation of **2** with dioxygen in the presence of Ph_3P affords hydrogen bonded triphenylphosphine oxide adduct $\{\text{Ph}_3\text{PO} \cdots \text{HN}(\text{PPh}_2)_2\text{Ni}(\text{PPh}_3)_2\}$ (**3**). Oxidation of **2** with an excess of O_2 affords pure triphenylphosphine oxide.

5. Supplementary material

Crystallographic data have been deposited with the Cambridge Crystallographic Data Centre, CCDC 204878 for **1** and CCDC 204879 for **3**. Copies of this information may be obtained free of charge from The Director, CCDC, 12, Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-336033); e-mail deposit@ccdc.cam.ac.uk or <http://www.ccdc.cam.ac.uk>.

Acknowledgements

We are grateful to the Russian Foundation for Basic Research (Grant 03-03-32051 and 1649.2003.3) for financial support of this work.

References

- [1] P. Bhattacharyya, J.D. Woollins, *Polyhedron* 14 (1995) 3367.
- [2] M. Witt, H.W. Roesky, *Chem. Rev.* 94 (1994) 1163.
- [3] P. Braunstein, R. Hasselbring, A. Tiripicchio, F. Ugozzoli, *J. Chem. Soc. Chem. Commun.* (1995) 37.
- [4] J. Ellerman, J. Sutter, F.A. Knoch, M. Moll, *Angew. Chem. Int. Ed. Engl.* 32 (1993) 700.
- [5] J. Ellerman, J. Sutter, C. Schelle, F.A. Knoch, M. Moll, *Z. Anorg. Allg. Chem.* 619 (1993) 2006.
- [6] A. Miedaner, R.C. Haltiwanger, D.L. DuBois, *Inorg. Chem.* 30 (1991) 417.
- [7] H. Noth, E. Fluck, *Z. Naturforsch.* 39b (1984) 744.
- [8] D.D. Perrin, W.L.F. Armarego, D.R. Perrin, *Purification of Laboratory Chemicals*, Pergamon, Oxford, 1980.
- [9] D.C. Bradley, M.B. Hursthouse, R.J. Smallwood, A.J. Welch, *J. Chem. Soc. Chem. Commun.* (1972) 872.
- [10] H. Noth, L. Meinel, *Z. Anorg. Allg. Chem.* 349 (1967) 225.
- [11] E. Upor, M. Mohai, Gy. Novak, *Photometric Methods in Inorganic Trace Analysis*, Akademiai Kiado, Budapest, 1985.
- [12] G.M. Sheldrick (1998). SADABS (2.01), BRUKER/Siemens Area Detector Absorption Correction Program, Bruker AXS, Madison, WI, USA.
- [13] G. Sheldrick, Bruker XRD, Madison, WI, USA.