# Ligand vs. Metal Basicity: Reactions of 2-(Diphenylphosphanyl)anilido and 2-(Diphenylphosphanyl)phenolato Complexes of Rhodium(I) and Iridium(I) with HBF<sub>4</sub>

Lutz Dahlenburg<sup>a</sup> and Konrad Herbst<sup>b</sup>

<sup>a</sup> Department Chemie und Pharmazie, Universität Erlangen-Nürnberg, Egerlandstraße 1,

91058 Erlangen, Germany <sup>b</sup> Haldor Topsøe A/S, Nymøllevej 55, 2800 Lyngby, Denmark

Reprint requests to Prof. Dr. Lutz Dahlenburg. Fax: +49 9131 85 27387. E-mail: Lutz.Dahlenburg@chemie.uni-erlangen.de

Z. Naturforsch. 2010, 65b, 376-382; received November 14, 2009

Dedicated to Professor Rolf W. Saalfrank on the occasion of his 70<sup>th</sup> birthday

Treatment of [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NR- $\kappa N, \kappa P$ )], where M/NR = Rh/NH (1), Rh/NCH<sub>3</sub> (2), Ir/NH (3), and Ir/NCH<sub>3</sub> (4), with Et<sub>2</sub>O·HBF<sub>4</sub> in CH<sub>2</sub>Cl<sub>2</sub> resulted in protonation at nitrogen with formation of [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NHR- $\kappa N, \kappa P$ )]BF<sub>4</sub> [M/NHR = Rh/NH<sub>2</sub> (7), Rh/NHCH<sub>3</sub> (8), Ir/NH<sub>2</sub> (9), Ir/NHCH<sub>3</sub> (10)]. Similar protonation of [Rh(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )] (5) in CH<sub>2</sub>Cl<sub>2</sub> afforded [Rh(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH- $\kappa O, \kappa P$ )]BF<sub>4</sub> (11), but furnished [Rh(CO)(PPh<sub>3</sub>)-(NCCH<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH- $\kappa P$ )]BF<sub>4</sub> (12) if carried out in CH<sub>3</sub>CN. [Ir(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )] (6) reacted with HBF<sub>4</sub> by protonation at the central metal atom and oxidative addition to give [IrH(FBF<sub>3</sub>)(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )] (13), the substitutionally labile BF<sub>4</sub><sup>-</sup> ligand of which underwent smooth exchange with neutral donors L producing [IrH(CO)(L)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )]BF<sub>4</sub> with L = H<sub>2</sub>O (14), CH<sub>3</sub>CN (15) and PPh<sub>3</sub> (16). The structures of 6 and 15 were determined by single-crystal X-ray crystallography.

Key words: P,N Ligands, P,O Ligands, Rhodium Complexes, Iridium Complexes, Protonation

# Introduction

In previous work we have been investigating several aspects of the coordination chemistry of bidentate 2-(diphenylphosphanyl)phenol, -thiophenol and -aniline ligands, both in their neutral 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>XH (X = O, S, NH, NCH<sub>3</sub>) and deprotonated 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>X<sup>-</sup> forms [1-5], in particular with regard to the reactivity of their Ir(I) and Rh(I) complexes towards selected Brønsted and Lewis acids [1,2]. In this context, the 2-(diphenylphosphanyl)anilido-substituted iridium(I) complex [Ir(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NH- $\kappa N, \kappa P$ )] was seen to react with HCl in CHCl<sub>3</sub> or toluene solution at -60 °C by oxidative addition to the central metal atom as well as protonation at nitrogen to form the ionic chelate complex [IrH(Cl)- $(CO)(PPh_3)(2-Ph_2PC_6H_4NH_2-\kappa N,\kappa P)]Cl,$  containing one of its NH groups hydrogen-bonded to Ir-Cl [2]. Treatment of the N-methylanilido compound  $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4NCH_3-\kappa N,\kappa P)]$  with hydrogen chloride under the conditions chosen for

the reaction of its NH analog with HCl also resulted in oxidative addition to iridium and protonation at nitrogen. However, different from the conversion of  $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4NH-\kappa N,\kappa P)]$ into the stable ionic product [IrH(Cl)(CO)(PPh3)- $(2-Ph_2PC_6H_4NH_2-\kappa N,\kappa P)$ ]Cl, the protonation of the N-methylanilido ligand was followed by dissociation of the NHCH<sub>3</sub> group from the metal, allowing the chloride ion to coordinate with formation of the covalent ring-opened product [IrHCl<sub>2</sub>(CO)(PPh<sub>3</sub>)- $(2-Ph_2PC_6H_4NHCH_3-\kappa P)]$ , stabilized by intramolecular -N(CH<sub>3</sub>)H····Cl-Ir hydrogen bonding [2]. In contrast, the ring-opened compound [IrHCl<sub>2</sub>(CO)(PPh<sub>3</sub>)- $(2-Ph_2PC_6H_4POH-\kappa P)$ ], resulting from combination of the chelated phenolato complex [Ir(CO)(PPh<sub>3</sub>)- $(2-Ph_2PC_6H_4O-\kappa O,\kappa P)$  with HCl in chloroform between -60 and +20 °C proved to be stable only in the presence of excess hydrogen chloride but otherwise was transformed by elimination of HCl and ring closure into [IrH(Cl)(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ ] [1]. With the aim of weighing the O- and N-

0932–0776 / 10 / 0300–0376 \$ 06.00 © 2010 Verlag der Zeitschrift für Naturforschung, Tübingen · http://znaturforsch.com

basicities of 2-(diphenylphosphanyl)anilido and -phenolato ligands against those displayed by coordinatively unsaturated metal centers, we have extended our previous studies to the reactivity of the rhodium and iridium chelate complexes [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>X- $\kappa P, \kappa X$ )] (X = NH, NHCH<sub>3</sub>, O) towards Brønsted acids which, different from HCl, possess non-coordinating or, at best, weakly coordination anions, *e. g.*, HBF<sub>4</sub>.

# **Results and Discussion**

Treatment of CH<sub>2</sub>Cl<sub>2</sub> solutions of the 16e chelate complexes [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NR- $\kappa N, \kappa P$ )] [2], where M/NR = Rh/NH (1), Rh/NCH<sub>3</sub> (2), Ir/NH (3), and Ir/NCH<sub>3</sub> (4), with one molar equivalent of HBF<sub>4</sub> (54% in diethyl ether) resulted in smooth protonation of the amide functions to form the corresponding tetrafluoroborate salts [M(CO)-(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NHR- $\kappa N, \kappa P$ )]BF<sub>4</sub> (Scheme 1, **7**–**10**). When allowed to interact with one additional equivalent of HBF<sub>4</sub>, none of the four  $d^8$  complexes **7**–**10** underwent any clean further transformation such as, *e. g.*, protonation at rhodium or iridium with or without decoordination of the NH<sub>2</sub> and NHCH<sub>3</sub> donors. Instead, intractable mixtures of products were obtained.

In the infrared spectra, the moderately air-stable yellow complex salts **7**–**10** exhibit a single carbonyl stretch band in the 1980–2010 cm<sup>-1</sup> region, each positioned *ca*. 50 cm<sup>-1</sup> at higher wavenumbers than those of the neutral starting compounds **1**–**4** [2], along with a very strong absorption between 1050 and 1070 cm<sup>-1</sup> arising from the triply degenerate v(BF) vibration of the anion. The NHCH<sub>3</sub> and NH<sub>2</sub>







groups manifest themselves by single and, respectively, split v(NH) absorptions around 3200 cm<sup>-1</sup> as well as by proton resonances observed as broad signals at  $\delta$  ca. 5.0 and 5.5 for the two aniline complexes 7 and 9, and at  $\delta$  ca. 4.3 and 5.1 for their N-methylated homologs 8 and 10. P,N-chelation in the [M(CO)- $(PPh_3)(2-Ph_2PC_6H_4NHR-\kappa N,\kappa P)]^+$  cations is evident from the pronounced downfield shifts [6] of their Ph<sub>2</sub>P <sup>31</sup>P{<sup>1</sup>H} resonances ( $\delta$  ca. 47 for M = Rh;  $\delta$  ca. 44 for M = Ir), which are approximately 66-69 ppm at lower field than those of the free 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NHR ligands ( $\delta$  ca. -22) and thus closely resemble the <sup>31</sup>P shift values observed for the two deprotonated P,N bidentates in the respective  $[M(CO)(PPh_3)(2-$ Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NR- $\kappa N$ , $\kappa P$ )] precursors ( $\delta$  ca. 49 for M = Rh;  $\delta$  ca. 38 for M = Ir) [2]. Coupling constants  ${}^{2}J_{P,P}$ of 280 to 285 Hz indicate the PPh3 and Ph2P donors to be coordinated in mutual *trans* position [7].

Rhodium complex **5** similarly underwent protonation of its anionic 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O<sup>-</sup> ligand when combined in CH<sub>2</sub>Cl<sub>2</sub> with ethereal HBF<sub>4</sub> in 1:1 stoichiometry. Both the *P*,*O*-chelated identity and the *trans*-P-Rh-P geometry of the resulting ionic product, [Rh(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH- $\kappa O, \kappa P$ )]BF<sub>4</sub> (**11**), have been established by IR and <sup>31</sup>P-NMR spectroscopy as outlined above for **7**–**10**: *v*(BF) = 1098, *v*(CO) = 2000 cm<sup>-1</sup>;  $\delta$ (Ph<sub>2</sub>P) = 43.6, <sup>2</sup>J<sub>P,P</sub> = 284.4 Hz. The presence of the phenolic OH group was evident from a broad <sup>1</sup>H resonance at  $\delta$  = 10.1.

Dissociation of the OH function from the metal atom with formation of [Rh(CO)(PPh<sub>3</sub>)(NCCH<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH- $\kappa P$ )]BF<sub>4</sub> (**12**) was observed when the reaction between **5** and an equimolar quantity of Et<sub>2</sub>O·HBF<sub>4</sub> was conducted in acetonitrile as a strongly donating solvent rather than in dichloromethane, which has only poor coordinating abilities [8–11]. The overall geometry of the cation [Rh(CO)(PPh<sub>3</sub>)(NCCH<sub>3</sub>)-(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH- $\kappa P$ )]<sup>+</sup> shown in Scheme 1 was confirmed by spectral data. In particular, the <sup>31</sup>P{<sup>1</sup>H}-NMR spectrum revealed very similar chemical shifts for the two *trans*-positioned phosphorus nuclei ( $\delta$  = 25.3 and 31.5; <sup>2</sup>J<sub>P,P</sub> = 292.2 Hz), which proves the 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>OH ligand to be bonded in a monodentate fashion.

Whereas the phenolato rhodium complex **5** reacted with HBF<sub>4</sub> in different solvents by protonation at oxygen to give **11** or **12**, the very same reaction of the phenolato iridium(I) compound [Ir(CO)-(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )] **(6)** afforded the hydrido complex [IrH(FBF<sub>3</sub>)(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O-

 $\kappa O, \kappa P$ ] (13) by oxidative addition of the acid. The outcome of this reaction, which parallels the oxidative addition of HBF<sub>4</sub> to trans-[IrCl(CO)(PPh<sub>3</sub>)<sub>2</sub>] [12], clearly shows the metal basicity of 6 to surpass that of the  $2-Ph_2PC_6H_4O^-$  chelate ligand. In agreement with the assignment of 13 as an iridium(III) complex containing a coordinated  $BF_4^-$  ion of  $C_{3v}$  symmetry, the IR-active triply degenerate v(BF) vibration of the free anion is now seen to be split into two components absorbing at 924 and 1125 cm<sup>-1</sup> ( $F_2 \rightarrow A_1 + E$ ). Coordination of the hydride cis to two mutually transpositioned P-donors and trans to Ir-FBF3 follows from both the  ${}^{31}P{}^{1}H$ -NMR spectrum consisting of two doublets split by 313.5 Hz and the <sup>1</sup>H-NMR spectrum containing a *cis-P*.*P*-coupled IrH pseudotriplet at  $\delta$  = -26.60, *i. e.*, close to the chemical shift previously reported as  $\delta = -26.5$  for trans-(H,F)-, trans-(P,P)-[IrH(Cl)(FBF<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>] [12].

As observed for other transition metal tetrafluoroborato complexes, the  $BF_4^-$  ligand of 13 is substitutionally labile and is easily displaced by diverse hard and soft donors to produce aqua, nitrile and phosphane derivatives such as, e.g., [IrH(CO)(OH<sub>2</sub>)- $(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$  (14), [IrH(CO)- $(NCCH_3)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$ (15), $[IrH(CO)(PPh_3)_2(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$ and (16). While the preparation of 14 required complex 13 to be isolated and purified prior to the  $BF_4$ <sup>-</sup>/H<sub>2</sub>O exchange reaction, compounds 15 and 16 were also obtained from the in situ protonation, with HBF4, of  $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]$  (6) either in neat acetonitrile or in CH<sub>2</sub>Cl<sub>2</sub> in the presence of added PPh<sub>3</sub>.

Notwithstanding that the aqua complex 14 contains a non-coordinated  $BF_4^-$  ion, the infrared spectrum displays three BF stretch bands at 995, 1064, and 1097  $cm^{-1}$ , which points to symmetry lowering from  $T_d$  to  $C_{2v}$   $(F_2 \rightarrow A_1 + B_1 + B_2)$  as a result of O-H···F hydrogen bonding between the complex cation and the  $BF_4^-$  counterion. Such hydrogen bonding interactions have previously been established, by vibrational spectroscopy and X-ray crystallography, for several related aqua complexes, representative examples of which are given by  $[IrH(Cl)(OH_2)(CO)(PPh_3)_2]BF_4$ [12],  $[Cr(CCH_3)(CO)_3 \{P(CH_3)_3\}(OH_2)]BF_4$  [13],  $[(\eta^7 C_7H_7$ )Mo(acac)(OH<sub>2</sub>)]BF<sub>4</sub> [14], and [Re(CO)<sub>3</sub>- $\{(CH_3)_2NCH_2CH_2N(CH_3)_2\}(OH_2)\}BF_4$  [15].

The geometry of the cation  $16^+$  has been concluded from its NMR features displaying (i)  ${}^{31}P{}^{1}H{}$ 



Fig. 1. Molecular structure of  $[IrH(CO)(NCCH_3)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$  (15) in the crystal (aryl H atoms omitted for clarity). Selected bond lengths (Å) and angles (deg): Ir–P(1) 2.322(2), Ir–P(2) 2.381(2), Ir–O(2) 2.053(4), Ir–N 2.134(5), Ir–C(1) 1.831(8), Ir–H, 1.6 (calcd.); P(1)–Ir–P(2) 169.26(6), P(1)–Ir–O(2) 84.0(1), P(1)–Ir–N 90.3(1), P(1)–Ir–C(1) 92.7(2), P(2)–Ir–O(2) 85.9(1), P(2)–Ir–N 92.5(1), P(2)–Ir–C(1) 97.2(2), O(2)–Ir–N 86.3(2), O(2)–Ir–C(1) 175.3(2), N–Ir–C(1) 97.1(3).

*ABX*-type splitting with *trans*- and *cis*-P,P couplings of 291.0, 15.3, and 13.3 Hz, respectively, (ii) pseudoquartet multiplicity of the <sup>13</sup>C{<sup>1</sup>H} carbonyl resonance, and (iii) an IrH doublet of virtual triplets characterized by *trans*-<sup>2</sup> $J_{P,H}$  = 148.0 Hz and |cis-<sup>2</sup> $J_{P,H}$  + *cis*-<sup>2</sup> $J_{P',H}|$  = 31.8 Hz.

Complex 15 was isolated as the addition compound  $15 \cdot 2C_3H_6O$  by crystallizing the residue of an evaporated reaction mixture of 6 and an equimolar quantity of Et<sub>2</sub>O·HBF<sub>4</sub> in CH<sub>3</sub>CN from an acetone/pentane solvent mixture. X-Ray crystal structure analysis confirmed the presence of a distorted octahedral cation in which the acetonitrile ligand is bonded trans to Ir-H (Fig. 1). The Ir-N distance, 2.134(5) Å, is within the range of 2.10 to 2.15 Å previously measured for various other cationic iridium(III) complexes possessing trans-H-Ir-NCCH<sub>3</sub> building blocks [16–19]. Ir–P and Ir–O bond lengths within the fivemembered chelate ring, 2.322(2) and 2.053(4) Å, are slightly longer than the Ir-P and Ir-O distances of 2.297(2) and 2.039(4) Å observed for [Ir(CO)- $(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)$ ] (6), the crystal structure of which has been determined for comparison (Fig. 2).

Molecule **6** displays the expected four-coordinate planar coordination geometry about the central metal atom as evidenced from the sum of the four interligand *cis* angles,  $360.2^{\circ}$ . Metal-to-ligand bond lengths and valence angles at the central metal atom reveal a close relation to [Rh(CO)-



Fig. 2. Molecular structure of  $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P]$  (6) in the crystal (H atoms omitted for clarity). Selected bond lengths (Å) and angles (deg): Ir–P(1) 2.336(2), Ir–P(2) 2.297(2), Ir–O(1) 2.039(4), Ir–C(37) 1.785(8); P(1)–Ir–P(2) 167.27(7), P(1)–Ir–O(1) 85.2(1), P(1)–Ir–C(37) 96.0(2), P(2)–Ir–O(1) 82.6(1), P(2)–Ir–C(37) 96.4(2), O(1)–Ir–C(37) 177.1(3).

 $(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)$ ] and  $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4NHCH_3-\kappa N,\kappa P)]$ , the structures of which have been reported previously [1, 2].

# **Concluding Remarks**

A comparative investigation of the reaction of HBF<sub>4</sub> with Vaska-type 2-(diphenylphosphanyl)anilido complexes of rhodium(I) and iridium(I), [M(CO)- $(PPh_3)(2-Ph_2PC_6H_4NR-\kappa N,\kappa P)]$  (R = H, CH<sub>3</sub>), has shown that, irrespective of the nature of the central metal atom, protonation occurs at nitrogen to form ionic 2-(diphenylphosphanyl)aniline derivatives  $[M(CO)(PPh_3)(2-Ph_2PC_6H_4NHR-\kappa N,\kappa P)]BF_4.$ 2-(diphenylphosphanyl)phenolato precursors The  $[Rh(CO)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]$  and [Ir(CO)- $(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]$  displayed divergent reactivity towards HBF<sub>4</sub>: the rhodium(I) compound, dissolved in CH<sub>2</sub>Cl<sub>2</sub> or CH<sub>3</sub>CN, was protonated by the acid at the phenolato function producing  $[Rh(CO)(PPh_3)(2-Ph_2PC_6H_4OH-\kappa O,\kappa P)]BF_4$ and  $[Rh(CO)(PPh_3)(NCCH_3)(2-Ph_2PC_6H_4OH-\kappa P)]BF_4,$ respectively. In marked contrast, the iridium(I) complex underwent protonation at the central metal atom with formation of hydridoiridium(III) products,  $[IrH(FBF_3)(CO)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]$  and  $[IrH(CO)(L)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4 (L =$ H<sub>2</sub>O, CH<sub>3</sub>CN, PPh<sub>3</sub>), under all conditions studied.

In conclusion, it has been demonstrated that the Brønsted basicity towards HBF<sub>4</sub> of the anionic anilido chelate ligands of the 16e complexes [M(CO)(PPh<sub>3</sub>)-(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NR- $\kappa N$ , $\kappa P$ )] (R = H, CH<sub>3</sub>) exceeds

the metal basicity of both Rh(I) and Ir(I), whereas the ligand basicity of the 2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O<sup>-</sup> chelate is superior to that of the central metal atom of [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ )] for M = Rh, but inferior for M = Ir.

#### **Experimental Section**

All manipulations were performed under nitrogen using standard Schlenk techniques. Solvents were distilled from the appropriate drying agents prior to use. – IR: Mattson Polaris – NMR: Bruker DPX 300 (300.1 MHz for <sup>1</sup>H, 75.5 MHz for <sup>13</sup>C, and 121.5 MHz for <sup>31</sup>P) with SiMe<sub>4</sub> as internal or H<sub>3</sub>PO<sub>4</sub> as external standards (downfield positive) at ambient temperature ("m" = deceptively simple multiplet). Complexes [M(CO)(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>NR- $\kappa N, \kappa P$ )] [M = Rh: R = H (1), CH<sub>3</sub> (2); M = Ir: R = H (3), CH<sub>3</sub> (4)] and [M(CO)-(PPh<sub>3</sub>)(2-Ph<sub>2</sub>PC<sub>6</sub>H<sub>4</sub>O- $\kappa O, \kappa P$ ] [M = Rh (5), Ir (6)] were prepared as previously described [1, 2].

# $[Rh(CO)(PPh_3)(2-Ph_2PC_6H_4NH_2-\kappa N,\kappa P)]BF_4(7)$

A solution of 100 mg (0.15 mmol) of **1** in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated with 21  $\mu$ L of HBF<sub>4</sub> (54% in diethyl ether; 0.15 mmol) and stirred for 1 h at r.t. Evaporation of all volatiles left a pale-yellow oil which was triturated with pentane to give 103 mg (90%) of **7** as yellow microcrystals. – IR (KBr): v = 3218/3120 (NH<sub>2</sub>), 2004 (CO), 1067 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 4.97$  (br, 2 H, NH<sub>2</sub>), 7.5 (m, 29 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 147.8$  (dd, <sup>2</sup>*J*<sub>P,C</sub> = 23.1 Hz, <sup>3</sup>*J*<sub>P,C</sub> = 5.3 Hz, phenylene C-1), 190.0 ("dt", <sup>1</sup>*J*<sub>Rh,C</sub> = 70.8 Hz,  $|cis^{-2}J_{P,C} + cis^{-2}J_{P',C}|$ = 31.0 Hz, CO). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 31.0$  (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 126.5 Hz, *trans*-<sup>2</sup>*J*<sub>P,P</sub> = 285.3 Hz, PPh<sub>3</sub>), 47.1 (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 123.9 Hz, Ph<sub>2</sub>P). – C<sub>37</sub>H<sub>31</sub>BF<sub>4</sub>NOP<sub>2</sub>Rh (757.32): calcd. C 58.68, H 4.13, N 1.85; found C 58.86, H 4.09, N 1.54.

Similar procedures were used for the preparation of complexes 8-10.

# $[Rh(CO)(PPh_3)(2-Ph_2PC_6H_4NHCH_3-\kappa N,\kappa P)]BF_4$ (8)

From 140 mg (0.21 mmol) of **2** and 28  $\mu$ L of 54% ethereal HBF<sub>4</sub> (0.21 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub>. – Yield: 140 mg (86%). – IR (KBr): v = 3244 (NH), 1996 (CO), 1057 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 2.15$  (d, <sup>3</sup>*J*<sub>H,H</sub> = 6.0 Hz, 3 H, CH<sub>3</sub>), 4.33 (br, 1 H, NH), 7.6 (m, 29 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 45.4$  (s, CH<sub>3</sub>), 154.7 (dd, <sup>2</sup>*J*<sub>P,C</sub> = 22.0 Hz, <sup>3</sup>*J*<sub>P,C</sub> = 3.5 Hz, phenylene C-1), 189.9 ("dt", <sup>1</sup>*J*<sub>Rh,C</sub> = 69.7 Hz, |*cis*-<sup>2</sup>*J*<sub>P,C</sub> + *cis*-<sup>2</sup>*J*<sub>P',C</sub>|= 32.4 Hz, CO). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 31.1$  (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 128.8 Hz, *trans*-<sup>2</sup>*J*<sub>P,P</sub> = 280.1 Hz, PPh<sub>3</sub>), 44.2 (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 126.4 Hz, Ph<sub>2</sub>P). – C<sub>38</sub>H<sub>33</sub>BF<sub>4</sub>NOP<sub>2</sub>Rh (771.34): calcd. C 59.17, H 4.31, N 1.82; found C 59.16, H 4.33, N 1.79.

#### $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4NH_2-\kappa N,\kappa P)]BF_4$ (9)

From 150 mg (0.20 mmol) of **3** and 27  $\mu$ L of 54 % ethereal HBF<sub>4</sub> (0.21 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub>. – Yield: 145 mg (86 %). – IR (KBr):  $\nu$  = 3244/3176 (NH<sub>2</sub>), 1980 (CO), 1069 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 5.45 (br, 2 H, NH<sub>2</sub>), 7.5 (m, 29 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = 147.5 (dd, <sup>2</sup>*J*<sub>P,C</sub> = 21.0 Hz, <sup>3</sup>*J*<sub>P,C</sub> = 6.3 Hz, phenylene C-1), 175.1 ("t", |*cis*-<sup>2</sup>*J*<sub>P,C</sub>+*cis*-<sup>2</sup>*J*<sub>P',C</sub>| = 20.8 Hz, CO). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = 28.1 (PPh<sub>3</sub>), 42.6 (Ph<sub>2</sub>P); both d, *trans*-<sup>2</sup>*J*<sub>P,P</sub> = 283.1 Hz. – C<sub>37</sub>H<sub>31</sub>BF<sub>4</sub>IrNOP<sub>2</sub> (846.63): calcd. C 52.49, H 3.69, N 1.65; found C 51.69, H 3.81, N 1.51.

#### $[Ir(CO)(PPh_3)(2-Ph_2PC_6H_4NHCH_3-\kappa N,\kappa P)]BF_4$ (10)

From 130 mg (0.18 mmol) of **4** and 24  $\mu$ L of 54% ethereal HBF<sub>4</sub> (0.18 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub>. – Yield: 140 mg (96%). – IR (KBr):  $\nu$  = 3227 (NH), 1987 (CO), 1057 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.20 (d, <sup>3</sup>J<sub>H,H</sub> = 5.8 Hz, 3 H, CH<sub>3</sub>), 5.07 (br, 1 H, NH), 7.7 (m, 29 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = 47.1 (s, CH<sub>3</sub>), 155.7 (dd, <sup>2</sup>J<sub>P,C</sub> = 20.4 Hz, <sup>3</sup>J<sub>P,C</sub> = 2.0 Hz, phenylene C-1), 174.3 ("t", |*cis*-<sup>2</sup>J<sub>P,C</sub>+*cis*-<sup>2</sup>J<sub>P',C</sub>| = 21.2 Hz, CO). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = 28.6 (PPh<sub>3</sub>), 40.0 (Ph<sub>2</sub>P); both d, *trans*-<sup>2</sup>J<sub>P,P</sub> = 281.2 Hz. – C<sub>38</sub>H<sub>33</sub>BF<sub>4</sub>IrNOP<sub>2</sub> (860.66): calcd. C 53.03, H 3.86, N 1.63; found C 52.73, H 4.03, N 1.13.

#### $[Rh(CO)(PPh_3)(2-Ph_2PC_6H_4OH-\kappa O,\kappa P)]BF_4(11)$

160 mg (0.24 mmol) of **5** and 33 μL of 54% ethereal HBF<sub>4</sub> (0.24 mmol) were combined in 10 mL of CH<sub>2</sub>Cl<sub>2</sub>. Stirring at ambient conditions and subsequent work-up as described for compound **7** afforded 169 mg (91%) of **11** as yellow crystals. – Yield: 165 mg (91%). – IR (KBr): v = 2000 (CO), 1098 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 7.1-7.6$  (m, 29 H, aryl H), 10.1 (br, 1 H, OH). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 161.2$  (dd, <sup>2</sup>*J*<sub>P,C</sub> = 18.0 Hz, <sup>3</sup>*J*<sub>P,C</sub> = 3.6 Hz, phenylene C-1), 188.6 ("dt", <sup>1</sup>*J*<sub>Rh,C</sub> = 82.1 Hz, |*cis*<sup>-2</sup>*J*<sub>P,C</sub> + *cis*<sup>-2</sup>*J*<sub>P',C</sub>| = 32.2 Hz, CO). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 28.0$  (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 125.9 Hz, *trans*-<sup>2</sup>*J*<sub>P,P</sub> = 284.4 Hz, PPh<sub>3</sub>), 43.6 (dd, <sup>1</sup>*J*<sub>Rh,P</sub> = 118.1 Hz, Ph<sub>2</sub>P). – C<sub>37</sub>H<sub>30</sub>BF<sub>4</sub>O<sub>2</sub>P<sub>2</sub>Rh (758.30): calcd. C 58.61, H 3.99; found C 57.58, H 3.87.

# $[Rh(CO)(PPh_3)(NCCH_3)(2-Ph_2PC_6H_4OH-\kappa P)]BF_4$ (12)

Treatment of a suspension of 160 mg (0.24 mmol) of **5** in 10 mL of acetonitrile with 33  $\mu$ L of 54 % ethereal HBF<sub>4</sub> (0.24 mmol) gave a clear solution which was stirred for 1 h at ambient conditions. Removal of the volatiles left the product as a bright-yellow solid which was washed with pentane and dried under vacuum. – Yield: 188 mg (98 %). – IR (KBr): v = 3394 (OH), 2296 (CN), 1999 (CO), 1094 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.98$  (s, 3 H, CH<sub>3</sub>), 2.6 (br, 1 H, OH), 7.0–8.1 (m, 29 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 1.6$  (s, CH<sub>3</sub>), 120.6 (s, CN), 188.0 (unresolved, CO).  $-{}^{31}P{^{1}H}$  NMR (CDCl<sub>3</sub>):  $\delta = 25.3$  (dd,  ${}^{1}J_{Rh,P} = 121.2$  Hz, *trans*- ${}^{2}J_{P,P} = 292.2$  Hz, PPh<sub>3</sub>), 31.5 (dd,  ${}^{1}J_{Rh,P} = 121.2$  Hz, Ph<sub>2</sub>P).  $- C_{39}H_{33}BF_{4}NO_{2}P_{2}Rh$  (799.34): calcd. C 58.60, H 4.16, N 1.75; found C 59.06, H 4.62, N 1.56.

## $[IrH(FBF_3)(CO)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]$ (13)

A mixture of 180 mg (0.24 mmol) of **6** and 33  $\mu$ L of 54 % ethereal HBF<sub>4</sub> (0.24 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was stirred for 1 h at r. t. Evaporation of the solvent followed by treatment of the residue with pentane left 195 mg (96 %) of **13** as a pale-yellow solid. – IR (KBr):  $\nu = 2054$  (CO), 1125/924 (FBF<sub>3</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = -26.60$  ("t" (br), 1 H, IrH), 6.8, 7.2 (both m, 2 H each, C<sub>6</sub>H<sub>4</sub>), 7.6 (m, 25 H, C<sub>6</sub>H<sub>5</sub>). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 14.6$  (PPh<sub>3</sub>), 36.6 (Ph<sub>2</sub>P); both d, *trans*-<sup>2</sup>*J*<sub>P,P</sub> = 313.5 Hz. – C<sub>37</sub>H<sub>30</sub>BF<sub>4</sub>IrO<sub>2</sub>P<sub>2</sub> (847.62): calcd. C 52.43, H 3.57; found C 52.53, H 3.80.

### $[IrH(CO)(OH_2)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$ (14)

A solution of 140 mg (0.17 mmol) of **13** in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated with 3  $\mu$ L of water. Stirring the mixture for 1 h at r. t. followed by evaporation to dryness afforded 143 mg (99%) of compound **14** as a pale-yellow solid which was washed with pentane and dried. – IR (KBr): v = 3408 (OH), 2276 (IrH), 2045 (CO), 1097/1064/995 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = -21.57$  ("t",  $|cis^{-2}J_{P,H} + cis^{-2}J_{P',H}| = 22.2$  Hz, 1 H, IrH), 3.5 (br, 2 H, H<sub>2</sub>O), 6.5, 7.0 (both m, 2 H each, C<sub>6</sub>H<sub>4</sub>), 7.3 (m, 25 H, C<sub>6</sub>H<sub>5</sub>). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta = 13.4$  (PPh<sub>3</sub>), 35.3 (Ph<sub>2</sub>P); both d, *trans*-<sup>2</sup>J<sub>P,P</sub> = 314.0 Hz. – C<sub>37</sub>H<sub>32</sub>BF<sub>4</sub>IrO<sub>3</sub>P<sub>2</sub> (865.63): calcd. C 51.34, H 3.73; found C 51.12, H 3.69.

# $[IrH(CO)(NCCH_3)(PPh_3)(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$ (15)

This compound was obtained from equimolar quantities of **6** and 54 % ethereal HBF<sub>4</sub> in acetonitrile using a procedure similar to the one outlined above for the rhodium complex **12**. The oily residue remaining after evaporation of the solvent was re-dissolved in acetone. Addition of pentane resulted in the gradual deposition of some off-white cystals which were identified as the acetone solvate  $15 \cdot 2C_3H_6O$  by X-ray structure analysis.

## $[IrH(CO)(PPh_3)_2(2-Ph_2PC_6H_4O-\kappa O,\kappa P)]BF_4$ (16)

Addition of 100  $\mu$ L of 54 % ethereal HBF<sub>4</sub> (0.73 mmol) and 83 mg (0.32 mmol) of PPh<sub>3</sub> to 240 mg (0.17 mmol) of **6**, dissolved in 10 mL of CH<sub>2</sub>Cl<sub>2</sub>, produced a colorless solution which was stirred at ambient conditions for 1 h and then evaporated. Complex **16** was left as an off-white residue which was washed with pentane and dried. – Yield: 355 mg (quantitative). – IR (KBr):  $\nu = 2127$  (IrH), 2040 (CO), 1092 (BF<sub>4</sub>) cm<sup>-1</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = –9.01 ("dt", trans-<sup>2</sup>J<sub>P,H</sub> = 148.0 Hz, |*cis*-<sup>2</sup>J<sub>P,H</sub> + *cis*-<sup>2</sup>J<sub>P',H</sub>| = 31.8 Hz, 1 H, IrH), 6.8 – 7.8 (m, 44 H, aryl H). – <sup>13</sup>C{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = 164.0 ("q",  $\Sigma$ |*cis*-<sup>2</sup>J<sub>P,C</sub>| = 24.0 Hz, CO), 176.9 ("dt", <sup>2</sup>J<sub>P,C</sub> = 18.9 Hz, |<sup>3</sup>J<sub>P,C</sub> + <sup>3</sup>J<sub>P',C</sub>| = 11.8 Hz, phenylene C-1). – <sup>31</sup>P{<sup>1</sup>H} NMR (CDCl<sub>3</sub>):  $\delta$  = –13.2, –0.6, 22.8 (*ABX* system, trans-<sup>2</sup>J<sub>P,P</sub> = 291.0 Hz, *cis*-<sup>2</sup>J<sub>P,P</sub> = 15.3 and 13.3 Hz). – C<sub>55</sub>H<sub>45</sub>BF<sub>4</sub>IrO<sub>2</sub>P<sub>3</sub> (1109.91): calcd. C 59.52, H 4.09; found C 58.90, H 3.85.

#### X-Ray structure determinations

Single crystals of **6**  $(0.30 \times 0.15 \times 0.13 \text{ mm}^3)$ and  $15 \cdot 2C_3H_6O$  (0.45 × 0.35 × 0.25 mm<sup>3</sup>) were grown from toluene/pentane and, respectively, acetone/pentane solvent mixtures. Diffraction measurements were made at ambient temperature on an Enraf-Nonius CAD-4 MACH 3 diffractometer using graphite-monochromatized  $MoK_{\alpha}$ radiation ( $\lambda = 0.71073$  Å); orientation matrices and unit cell parameters from the setting angles of 25 centered medium-angle reflections; collection of the diffraction intensities by  $\omega$  scans; data empirically corrected for absorption using  $\psi$  scans [20] (6: T<sub>min</sub> = 0.349, T<sub>max</sub> = 0.596;  $15 \cdot 2C_3H_6O$ :  $T_{min} = 0.332$ ,  $T_{max} = 0.507$ ). The structures were solved by Direct Methods and subsequently refined by full-matrix least-squares procedures on  $F^2$ with allowance for anisotropic thermal motion of all non-hydrogen atoms employing the WINGX package [21] with the programs SIR-97 [22], SHELXL-97 [23], and ORTEP-3 [24] implemented therein. - 6: C<sub>37</sub>H<sub>29</sub>IrO<sub>2</sub>P<sub>2</sub> (759.74); monoclinic,  $P2_1/n$ , a = 10.022(9), b = 14.861(4),

- L. Dahlenburg, K. Herbst, M. Kühnlein, Z. Anorg. Allg. Chem. 1997, 623, 250–258.
- [2] L. Dahlenburg, K. Herbst, Chem. Ber./Recueil 1997, 130, 1693-1698.
- [3] L. Dahlenburg, K. Herbst, H. Berke, J. Organomet. Chem. 1999, 585, 225–233.
- [4] L. Dahlenburg, K. Herbst, J. Chem. Soc., Dalton Trans. 1999, 3935 – 3939.
- [5] L. Dahlenburg, K. Herbst, A. Zahl, J. Organomet. Chem. 2000, 616, 19–28.
- [6] P.E. Garrou, Chem. Rev. 1981, 81, 229-266.
- [7] P. S. Pregosin, R. W. Kunz in *NMR Basic Principles and Progress*, Vol 16: <sup>31</sup>P and <sup>13</sup>C NMR of Transition Metal Phosphine Complexes (Eds.: P. Diehl, E. Fluck, R. Kosfeld), Springer, Berlin, **1979**, chapter B.
- [8] K. Sünkel, G. Urban, W. Beck, J. Organomet. Chem. 1983, 252, 187–197.
- [9] J. M. Fernandez, J. A. Gladysz, Organometallics 1989, 8, 207–219.

c = 20.903(5) Å,  $\beta = 94.71(4)^{\circ}$ , V = 3103(3) Å<sup>3</sup>, Z = 4,  $D_{\text{calcd}} = 1.63 \text{ g cm}^{-3}, \ \mu(\text{Mo}K_{\alpha}) = 4.4 \text{ mm}^{-1}, \ F(000) =$ 1496;  $2.58^\circ < \Theta < 25.02^\circ$ , 5618 reflections collected  $(-11 \le h \le +11, 0 \le k \le +17, 0 \le l \le +24)$ , 5463 unique  $(R_{int} = 0.0381); wR(F^2) = 0.0492$  for all data and 379 parameters, R(F) = 0.0349 for 2504 structure factors  $F_0 \ge 4\sigma(F_0)$ ; weighting scheme applied:  $w = 1/[\sigma^2(F_0) + (0.0094P)^2],$ where  $P = (F_0^2 + 2F_c^2)/3$ ; largest peak and hole in final difference map: 0.871 and  $-0.863 \text{ e} \text{ Å}^{-3}$ .  $-15 \cdot 2C_3 H_6 O$ :  $C_{45}H_{45}BF_4IrNO_4P_2$  (1004.77); triclinic,  $P\bar{1}$ , a = 9.164(6), b = 14.700(2), c = 16.515(2) Å,  $\alpha = 91.26(1), \beta =$ 93.28(3),  $\gamma = 94.74(2)^{\circ}$ , V = 2212.7(15) Å<sup>3</sup>, Z = 2,  $D_{\text{calcd}} = 1.51 \text{ g cm}^{-3}, \ \mu(\text{Mo}K_{\alpha}) = 3.1 \text{ mm}^{-1}, \ F(000) =$ 1004;  $2.53^{\circ} \leq \Theta \leq 26.96^{\circ}$ , 9966 reflections collected  $(-11 \le h \le +11, -18 \le k \le +18, 0 \le l \le +21),$ 9626 unique ( $R_{int} = 0.0284$ );  $wR(F^2) = 0.1098$  for all data, 529 parameters, and 16 restraints, R(F) = 0.0493for 7160 structure factors  $F_{\rm o} \ge 4\sigma(F_{\rm o})$ ; weighting scheme applied:  $w = 1/[\sigma^2(F_0) + (0.0512P)^2]$ , where  $P = (F_0^2 + 2F_c^2)/3$ ; largest peak and hole in final difference map: 1.746 and  $-1.035 \text{ e} \text{ Å}^{-3}$ .

CCDC 753417 (6) and CCDC 753418 ( $15 \cdot 2C_3H_6O$ ) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre *via* www.ccdc.cam.ac.uk/data\_request/cif.

#### Acknowledgements

Support of this work by the Deutsche Forschungsgemeinschaft (Bonn, SFB 583) is gratefully acknowledged. We are also indebted to Mrs. S. Hoffmann for her skilful assistance.

- [10] J. Huhmann-Vincent, B.L. Scott, G. Kubas, *Inorg. Chem.* 1999, 38, 115–124.
- B. Olgemüller, H. Bauer, H. Löbermann, U. Nagel, W. Beck, *Chem. Ber.* 1982, 115, 2271 – 2286.
- [12] H. Bauer, U. Nagel, W. Beck, J. Organomet. Chem. 1985, 290, 219–229.
- [13] K. Richter, E. O. Fischer, C. G. Kreiter, J. Organomet. Chem. 1976, 122, 187–196.
- [14] M. Green, H. P. Kirsch, F. G. A. Stone, A. J. Welch, J. Chem. Soc., Dalton Trans. 1977, 1755 – 1761.
- [15] E. Horn, M. R. Snow, Aust. J. Chem. 1984, 37, 1475– 1393.
- [16] X. D. He, J. Fernandez-Baeza, B. Chaudret, K. Folting, K. G. Caulton, *Inorg. Chem.* **1990**, *29*, 5000 – 5002.
- [17] D. Hermann, M. Gandelman, H. Rozenberg, L. J. W. Shimon, D. Milstein, *Organometallics* 2002, 21, 812– 818.
- [18] F. Acha, M. A. Garralda, L. Ibarlucea, E. Pinilla, M. R. Torres, *Inorg. Chem.* **2005**, *44*, 9084–9091.

- [19] E. Ben-Ari, R. Cohen, M. Gandelman, L. J. W. Shimon, J. M. L. Martin, D. Milstein, *Organometallics* **2006**, *25*, 3190–3210.
- [20] A.C.T. North, D.C. Phillips, F.S. Mathews, Acta Crystallogr. 1968, A24, 351–359.
- [21] L. J. Farrugia, WINGX, A MS-Windows System of Programs for Solving, Refining and Analysing Single Crystal X-ray Diffraction Data for Small Molecules, University of Glasgow, Glasgow, Scotland (U.K.)
  2005. See also: L. J. Farrugia, J. Appl. Crystallogr. 1999, 32, 837-838.
- [22] A. Altomare, M. C. Burla, M. Camalli, G. L. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. C. Moliterni, G. Polidori, R. Spagna, SIR97, A Program for

the Automatic Solution of Crystal Structures by Direct Methods; see also: *J. Appl. Crystallogr.* **1999**, *32*, 115–119.

- [23] G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures, University of Göttingen, Göttingen (Germany) 1997. See also: G. M. Sheldrick, *Acta Crystallogr.* 2008, A64, 112–122.
- [24] C. K. Johnson, M. N. Burnett, ORTEP-3 (version 1.0.2), Rep. ORNL-6895, Oak Ridge National Laboratory, Oak Ridge, TN (USA) **1996**. Windows version: L. J. Farrugia, University of Glasgow, Glasgow, Scotland (U.K.) **1999**. See also: L. J. Farrugia, *J. Appl. Crystallogr.* **1997**, *30*, 565.