

Photochemistry. X.¹⁾ Photolysis of Tetrazolo[1,5-*b*]pyridazines

TAKASHI TSUCHIYA, HEIHACHIRO ARAI, and HIROSHI IGETA

School of Pharmaceutical Sciences, Showa University²⁾

(Received March 24, 1973)

Photolysis of 3-azidopyridazine, *i.e.*, tetrazolo[1,5-*b*]pyridazine (I) was examined. Unsubstituted and methyl-substituted compounds (Ia—d) afforded 3-cyanocyclopropenes (III). On the other hand, 6-methoxy compound (Ie) gave the triene (IV) and the ethylene (V and VI). These compounds are considered to be formed *via* the diazo intermediates (VIII) and the carbenes (X).


Concerning the photolyses of azides³⁾ and polyaza compounds such as tetrazoles⁴⁾ and triazoles,⁵⁾ many reports have been published. Most of them referred to the interesting result that the key intermediates were nitrenes and carbenes formed by elimination of molecular nitrogen. Similarly, photolysis of the diazo compounds⁶⁾ also gave various compounds *via* the carbenes.

Meanwhile, as for pyridazines, photolysis^{2,7,8)} of their N-oxide resulted in the formation of key intermediate of diazo compound and carbene. However, there have been no reports on the photolysis of tetrazolopyridazines.

On the azide-tetrazole tautomerism of 3-azidopyridazine, Wentrup⁹⁾ and Sasaki, *et al.*¹⁰⁾ reported that in solution it existed completely in the tetrazole form. And some reports have been already published on gas phase pyrolysis¹¹⁾ and thermal reaction¹⁰⁾ of 3-azidopyridazine. Thus, we have been interested in the photolysis of this compound and have obtained some interesting results, on which we now report.¹²⁾

Tetrazolo[1,5-*b*]pyridazine (I) was irradiated in dichloromethane for 6—7 hr and from Ia—d, the corresponding 3-cyanocyclopropenes (IIa—d) were obtained in 20—25% yields, respectively. The compound Ib gave also 3-cyano-5-methylpyrazole (IIIb) in a trace amount (0.1%) along with IIb. Except for Id, other compounds (Ia—c) afforded trace amounts of the compounds presumably to be the corresponding III, which were recognized by gas-liquid chromatography (GLC) and thin-layer chromatography (TLC) but could not be isolated. In all cases, an irradiation for 6—7 hr resulted in the recovery of the starting material in 20%,

- 1) Part IX: T. Tsuchiya, H. Arai, and H. Igeta, *Tetrahedron*, **29**, 2747 (1973).
- 2) Location: 1-5-8, Hatanodai, Shinagawa-ku, Tokyo, 142, Japan.
- 3) R.L. Whistler and A.K.M. Anisuzzaman, *J. Org. Chem.*, **34**, 3823 (1969); R.A. Abramovitch, G.N. Knaus, and V. Uma, *J. Am. Chem. Soc.*, **91**, 7532 (1969).
- 4) P. Scheider, *J. Org. Chem.*, **34**, 199 (1969); R.R. Fraser, *ibid.*, **34**, 4118 (1969); F.L. Bach, J. Karliner, and G.E. van Lear, *Chem. Commun.*, **1969**, 1110; J. Hyatt and J.S. Swenton, *J. Heterocyclic Chem.*, **9**, 409 (1972).
- 5) E.M. Burgess, R. Carithers, and L. McCullagh, *J. Am. Chem. Soc.*, **90**, 1923 (1968); A.J. Hubert, *Chem. Commun.*, **1969**, 328; R.H. Spector and M.M. Joullie, *J. Heterocyclic Chem.*, **6**, 605 (1969).
- 6) G.L. Closs and W.A. Böll, *J. Am. Chem. Soc.*, **85**, 3904 (1963); R.W. Hoffman and H.J. Luthardt, *Chem. Ber.*, **101**, 3861 (1968).
- 7) P.L. Kumler and O. Buchardt, *J. Am. Chem. Soc.*, **90**, 5640 (1968).
- 8) T. Tsuchiya, H. Arai, and H. Igeta, *Tetrahedron Letters*, **1971**, 2579; T. Tsuchiya, H. Arai, T. Tonami, and H. Igeta, *Chem. Pharm. Bull.* (Tokyo), **20**, 300 (1972).
- 9) C. Wentrup, *Tetrahedron*, **26**, 4969 (1970).
- 10) T. Sasaki, K. Kanematsu, and M. Murata, *J. Org. Chem.*, **36**, 446 (1971).
- 11) C. Wentrup and W. D. Crow, *Tetrahedron*, **26**, 4915 (1970).
- 12) A part of this work has been reported in a preliminary communication; T. Tsuchiya, H. Arai, and H. Igeta, *Chem. Commun.*, **1972**, 1059.



larger contribution of the structure (XII), it may not afford cyclopropene but the triene (IV) by dimerization and ethylene (V and VI) by addition¹⁴⁾ of the solvent alcohol.

Experimental

Photolyses were carried out in an immersion apparatus equipped with 200W high pressure mercury lamp (Nikko Sekiei Co., Japan) and cooled internally with running water. IR spectra were determined with a JASCO IRA-1 spectrometer and mass spectra were recorded on a Hitachi RMS-4 instrument. NMR spectra were recorded on Hitachi R-20 and R-22 spectrometers in CCl_4 or CDCl_3 solution using TMS as internal standard. Melting points were measured on a Yamato MP-1 apparatus and are uncorrected. Column and thin layer chromatography were carried out with alumina and silica gel obtained from Merck Co., Ltd.

Preparation of Tetrazolo[1,5-*b*]pyridazines (I)—According to the method¹⁵⁾ described by Itai and Kamiya, one drop of conc. HCl was added to a solution of 3-chloropyridazine derivatives and sodium azide dissolved in $\text{EtOH-H}_2\text{O}$ (2: 1) and heated in a sealed tube at $80-90^\circ$ for 7–8 hr. After cooling the deposited crystals were collected and recrystallized from benzene to give I in 40–60% yield.

i) Tetrazolo[1,5-*b*]pyridazine (Ia): This compound was prepared from 3-chloropyridazine, mp $108-109^\circ$ (lit.¹³⁾ $110-111^\circ$.

ii) 6-Methyltetrazolo[1,5-*b*]pyridazine (Ib): This compound was prepared from 3-chloro-6-methylpyridazine,¹⁶⁾ mp $139-141^\circ$. Mass Spectrum m/e : 135 (M^+). Anal. Calcd. for $\text{C}_5\text{H}_5\text{N}_5$: C, 44.44; H, 3.73; N, 51.83. Found: C, 44.71; H, 3.83; N, 51.54.

iii) 5-Methyltetrazolo[1,5-*b*]pyridazine (Ic): This compound was prepared from 3-chloro-5-methylpyridazine,¹⁷⁾ mp $130-131^\circ$. Mass Spectrum m/e : 135 (M^+). Anal. Calcd. for $\text{C}_5\text{H}_5\text{N}_5$: C, 44.44; H, 3.73; N, 51.83. Found: C, 44.57; H, 3.61; N, 51.73.

iv) 4-Methyltetrazolo[1,5-*b*]pyridazine (Id): This compound was prepared from 3-chloro-4-methylpyridazine,¹⁷⁾ mp $119-121^\circ$. Mass Spectrum m/e : 135 (M^+). Anal. Calcd. for $\text{C}_5\text{H}_5\text{N}_5$: C, 44.44; H, 3.73; N, 51.83. Found: C, 44.38; H, 3.52; N, 51.82.

v) 6-Methyltetrazolo[1,5-*b*]pyridazine (Ie): This compound was prepared from 3-chloro-6-methylpyridazine, mp $155-156^\circ$ (lit.¹⁰⁾ 154.5° .

14) D. Bethell and R.D. Howard, *J. Chem. Soc. (B)*, **1969**, 745; P.K. Freeman and R.C. Johnson, *J. Org. Chem.*, **34**, 1746 (1969); H. Chaimovich, R.J. Vaughan, and F.H. Westheimer, *J. Am. Chem. Soc.*, **90**, 4088 (1968).

15) T. Itai and S. Kamiya, *Chem. Pharm. Bull. (Tokyo)*, **11**, 342 (1963).

16) M. Kumagai, *Nippon Kagaku Zasshi*, **81**, 489 (1960); N.K. Basu and F.L. Rose, *J. Chem. Soc.*, **1963**, 5560.

17) T. Takabayashi, *Chem. Pharm. Bull. (Tokyo)*, **5**, 229 (1957).

Photolysis of Tetrazolo[1,5-*b*]pyridazine (Ia—d)—A solution of I (2 g) dissolved in dichloromethane (250 ml) was irradiated for 6—7 hr and the reaction mixture was evaporated *in vacuo*. The residue was chromatographed on alumina. From the eluate with benzene, 3-cyanocyclopropene (IIa—d) was obtained. Their spectral data are collected in Table I. Then, from the eluate with dichloromethane, the starting material (I) was recovered in 18—20% yield. In the case of Ib, from the first eluate with CH₂Cl₂, 3-cyanopyrazole (IIIb) was obtained.

3-Cyanocyclopropene (IIa): Yield 24%, bp₁₀ 55° (bath temp.). *Anal.* Calcd. for C₄H₃N: C, 73.83; H, 4.65; N, 21.53. Found: C, 73.78; H, 4.88; N, 21.31.

1-Methyl-3-cyanocyclopropene (IIb): Yields 20% from Ib, 23% from Ic, bp₁₀ 60° (bath temp.). *Anal.* Calcd. for C₅H₅N: C, 75.92; H, 6.37; N, 17.71. Found: C, 76.25; H, 6.13; N, 17.70.

3-Methyl-3-cyanocyclopropene (IIc): Yield 26%, bp₁₀ 60° (bath temp.). *Anal.* Calcd. for C₅H₅N: C, 75.92; H, 6.37; N, 17.71. Found: C, 75.66; H, 6.47; N, 17.43.

3-Cyano-5-methylpyrazole (IIIb): Yield 0.1%, mp 112—114° (from iso-Pr₂O), Mass Spectrum *m/e*: 107 (M⁺), IR $\nu_{\text{max}}^{\text{KBr}}$: 2270 cm⁻¹ (CN), NMR δ (CDCl₃): 2.48 (3H, s, 5-CH₃), 6.49 (1H, s, H₄). *Anal.* Calcd. for C₅H₅N₃: C, 56.06; H, 4.71; N, 39.23. Found: C, 55.81; H, 4.75; N, 39.11.

Photolysis of 6-Methoxytetrazolo[1,5-*b*]pyridazine (Ie)—i) Dichloromethane Solution: Similar to the cases of Ia—d, the residue was chromatographed on alumina. From the eluate with benzene, 1,6-dicyano-3,4-dimethoxy-1,3,5-hexatriene (IV) was obtained in ca. 5% yield. Colorless crystals, mp 177—179° (from benzene), Mass Spectrum *m/e*: 190 (M⁺), IR $\nu_{\text{max}}^{\text{KBr}}$: 2220 cm⁻¹ (CN), NMR δ (CDCl₃): 3.68 (6H, s, -OCH₃), 5.72 (2H, d, *J*=16.0, H₁ and H₆), 7.25 (2H, d, *J*=16.0, H₂ and H₅). *Anal.* Calcd. C₁₀H₁₀O₂N₂: C, 63.15; H, 5.30; N, 14.73. Found: C, 63.06; H, 5.10; N, 14.67. Then, from the eluate with CH₂Cl₂, the starting material was recovered in 17% yield.

ii) Methanol Solution: Under similar condition to i), irradiation was continued until the starting material was consumed (about 3 hr). The reaction mixture was evaporated and the residue was chromatographed on alumina. The eluate with benzene was again chromatographed on silica gel. From the eluates with benzene containing 5% AcOEt, 1-cyano-2-dimethoxymethyl-ethylene (V) and then IV were obtained in 21% and 2—3% yields, respectively. V: bp₁₀ 80° (bath temp.), Mass Spectrum *m/e*: 127 (M⁺), IR $\nu_{\text{max}}^{\text{KBr}}$: 2230 cm⁻¹ (CN), NMR δ (CDCl₃): 3.73 (6H, s, -OCH₃), 5.02 (1H, d, *J*=6.0, -CH(OCH₃)₂), 5.47 (1H, d, *J*=12.0, H₁), 6.28 (1H, d.d, *J*=6.0 and 12.0, H₂). *Anal.* Calcd. for C₆H₉O₂N: C, 56.68; H, 7.14; N, 11.02. Found: C, 56.39; H, 7.03; N, 11.35.

iii) Ethanol Solution: Similar to ii), 1-cyano-2-(ethoxymethoxymethyl)-ethylene (VI) and IV were obtained in 19% and 2—3% yields, respectively. VI: bp₃ 80° (bath temp.), Mass Spectrum *m/e*: 141 (M⁺), IR $\nu_{\text{max}}^{\text{KBr}}$: 2230 cm⁻¹ (CN), NMR δ (CDCl₃): 1.23 (3H, t, -CH₂CH₃), 3.36 (3H, s, -OCH₃), 3.60 (2H, q, -CH₂CH₃), 5.07 (1H, d, *J*=6.0, -CH-), 5.45 (1H, d, *J*=12.0, H₁), 6.28 (1H, d.d, *J*=6.0 and 12.0, H₂). *Anal.* Calcd. for C₇H₁₁O₂N: C, 59.55; H, 7.85; N, 9.92. Found: C, 59.64; H, 7.60; N, 10.16.