

# Azido[1,1'-bis(diphenylphosphino)-ferrocene](pentamethylcyclopentadienyl)rhodium(III) hexafluorophosphate

Won Seok Han and Soon W. Lee\*

Department of Chemistry (BK21), Institute of Basic Science, Sungkyunkwan University, Natural Science Campus, Suwon 440-746, South Korea  
Correspondence e-mail: soonwlee@skku.edu

Received 19 February 2004


Accepted 2 March 2004

Online 31 March 2004

In the title compound, azido-2κN-bis[μ-(1η<sup>5</sup>:2κP)-diphenylphosphinocyclopentadienyl][2(η<sup>5</sup>-pentamethylcyclopentadienyl)iron(III)rhodium(III) hexafluorophosphate,  $[\{\text{Rh}(\text{C}_{10}\text{H}_{15}(\text{N}_3)\}\{\text{Fe}(\mu\text{-C}_{17}\text{H}_{14}\text{P})_2\})\text{PF}_6$  or  $[\text{FeRh}(\text{C}_{10}\text{H}_{15})(\mu\text{-C}_{17}\text{H}_{14}\text{P})_2(\text{N}_3)]\text{PF}_6$ , the coordination sphere of  $\text{Rh}^{\text{III}}$  can be described as pseudo-tetrahedral, composed of two P atoms from a 1,1'-bis(diphenylphosphino)ferrocene (dppf) ligand, an azido N atom and the centroid of the ring of a  $\text{C}_5\text{Me}_5$  ( $\text{Cp}^*$ ) ligand. The two cyclopentadienyl rings in the dppf moiety adopt an eclipsed conformation. The  $\text{Rh}\cdots\text{Fe}$  distance is 4.340 (2) Å.

## Comment

Various redox-active ligands are employed to control the reactivities of transition-metal complexes (Gan & Hor, 1995). The dppf ligand [1,1'-bis(diphenylphosphino)ferrocene,  $\text{Fe}(\text{C}_5\text{H}_4\text{PPh}_2)_2$ ] and its derivatives are well known redox-active ligands, and their complexes usually exhibit a ferrocene-based oxidation process, together with additional redox



processes at other metal centres. Recently, an interesting Rh-dppf chloro complex,  $[\text{Cp}^*\text{RhCl}(\text{dppf})]\text{PF}_6$ , was reported ( $\text{Cp}^*$  is  $\text{C}_5\text{Me}_5$ ), which was prepared by treating a chloro-bridged  $\text{Rh}^{\text{III}}$  dimer,  $[\text{Cp}^*\text{RhCl}_2]_2$ , with dppf and  $\text{NaPF}_6$  (Ma & Yamamoto, 1999). In order to compare structural features and to examine the redox properties of the Rh-dppf azide complex with respect to its chloro analogue, we have prepared the title compound, (I), and present its structure here.

The coordination sphere of the Rh metal in (I) can be described as pseudo-tetrahedral. The two Cp (cyclopentadienyl) rings of the dppf ligand are not perfectly parallel, but are twisted from each other with a dihedral angle of 3.4 (1)°. The  $\text{P1}-\text{C11}\cdots\text{C16}-\text{P2}$  pseudo-torsion angle is 3.0 (4)°, indicating that the two Cp rings adopt an eclipsed conformation. For comparison, the ideal torsion angle for a *gauche* (or staggered) conformation is 36°. Both  $\text{Fe}-\text{Cg}$  ( $\text{Cg}$  is the centroid of a Cp ring) distances are 1.63 Å, and the  $\text{Cg1}-\text{Fe}-\text{Cg2}$  angle ( $\text{Cg1}$  is the centroid of the  $\text{C11}-\text{C15}$  ring and  $\text{Cg2}$  is the centroid of the  $\text{C16}-\text{C20}$  ring) is 177.5°. The  $\text{Cg}^*\cdots\text{Rh1}-\text{P1}$ ,  $\text{Cg}^*\cdots\text{Rh1}-\text{P2}$  and  $\text{Cg}^*\cdots\text{Rh1}-\text{N1}$  angles ( $\text{Cg}^*$  is the centroid of the  $\text{C1}-\text{C5}$  ring) are 125.5, 127.01 and 126.8°, respectively. The  $\text{P1}\cdots\text{Fe}\cdots\text{P2}$  bite angle is 60.68 (5)°, and the  $\text{P1}\cdots\text{P2}$  distance is 3.526 (3) Å. These bonding parameters within a ferrocene moiety are consistent with those found in the chloro analogue  $[\text{Cp}^*\text{RhCl}(\text{dppf})]\text{PF}_6$  (Ma & Yamamoto, 1999).




Figure 1

The molecular structure of (I), showing 50% probability displacement ellipsoids. H atoms have been omitted for clarity.

The  $\text{Rh}-\text{N1}-\text{N2}$  bond angle of 127.6 (7)° in (I) agrees well with the values found in six-coordinate  $\text{Rh}^{\text{III}}$ -azide complexes (Seok *et al.*, 2002). Atom N3 in the azide ligand has a large displacement ellipsoid, probably due to its terminal position. The  $\text{Rh}-\text{N}$  bond length of 2.111 (7) Å indicates an  $\text{Rh}-\text{N}$  single bond (Davis *et al.*, 1969; Lee & Lee, 1999). The  $\text{Rh}\cdots\text{Fe}$  distance is 4.340 (2) Å, which clearly rules out a direct bonding interaction between the two metal atoms.

## Experimental

A solution of  $[\text{Cp}^*\text{RhCl}(\text{dppf})]\text{PF}_6$  (100 mg, 0.12 mmol) and  $\text{AgNO}_3$  (62 mg, 0.36 mmol) in a mixture of dichloromethane and acetone (1:1, 30 ml) was stirred for 3 h at room temperature, and the solvent was removed under vacuum. The resulting solid was extracted with

dichloromethane and neat  $\text{N}_3\text{SiMe}_3$  (16  $\mu\text{l}$ , 0.12 mmol) was added. The resulting solution was then stirred for 18 h and the solvent was removed. The remaining solid was washed with diethyl ether (20 ml  $\times$  2) to give the title compound, (I), which was recrystallized from dichloromethane–hexane (1:1) to give red crystals (0.074 g, 62.6%; m.p. 471–473 K). Spectroscopic analysis, IR (KBr,  $\nu$ ,  $\text{cm}^{-1}$ ): 2029 ( $\text{N}_3$ ), 843 ( $\text{PF}_6$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , p.p.m.): 7.74–7.33 (20H, *m*, Ph), 4.91, 4.39, 4.29, 4.12 (8H,  $\text{C}_5\text{H}_4$ ), 1.15 (15H, *t*,  $J_{\text{PH}} = 4.0$  Hz,  $\text{C}_5\text{Me}_5$ );  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , p.p.m.): 135.7–128.8 (Ph), 104.7 (*d*,  $J_{\text{RhC}} = 6.4$  Hz,  $\text{Cp}^* \text{C}_5$ ), 77.5–76.6 ( $\text{C}_5\text{H}_4$ ), 8.59 ( $\text{Cp}^* \text{Me}_5$ );  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{CDCl}_3$ , p.p.m.): 38.79 (*d*,  $J_{\text{RhP}} = 146.5$  Hz, dppf), –143.8 (*sep*,  $J_{\text{PF}} = 708.2$  Hz); analysis calculated for  $\text{C}_{44}\text{H}_{43}\text{F}_6\text{Fe}-\text{N}_3\text{P}_3\text{Rh}$ : C 53.95, H 4.43, N 4.29%; found: C 54.23, H 4.39, N 4.24%.

## Crystal data

|                                                                                                          |                                           |
|----------------------------------------------------------------------------------------------------------|-------------------------------------------|
| $[\text{FeRh}(\text{C}_{10}\text{H}_{15})(\text{C}_{17}\text{H}_{14}\text{P})_2(\text{N}_3)]\text{PF}_6$ | Mo $K\alpha$ radiation                    |
| $M_r = 979.48$                                                                                           | Cell parameters from 26 reflections       |
| Monoclinic, $P2_1/n$                                                                                     | $\theta = 5.1\text{--}13.1^\circ$         |
| $a = 14.759$ (3) Å                                                                                       | $\mu = 0.93 \text{ mm}^{-1}$              |
| $b = 17.829$ (4) Å                                                                                       | $T = 295$ (2) K                           |
| $c = 15.736$ (4) Å                                                                                       | Block, red                                |
| $\beta = 92.669$ (12)°                                                                                   | $0.26 \times 0.24 \times 0.18 \text{ mm}$ |
| $V = 4136.3$ (16) Å <sup>3</sup>                                                                         |                                           |
| $Z = 4$                                                                                                  |                                           |
| $D_x = 1.573 \text{ Mg m}^{-3}$                                                                          |                                           |

## Data collection

|                                                                 |                                    |
|-----------------------------------------------------------------|------------------------------------|
| Siemens P4 diffractometer                                       | $R_{\text{int}} = 0.052$           |
| $\omega$ scans                                                  | $\theta_{\text{max}} = 25.0^\circ$ |
| Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968) | $h = -17 \rightarrow 0$            |
| $T_{\text{min}} = 0.698$ , $T_{\text{max}} = 0.914$             | $k = 0 \rightarrow 21$             |
| 7547 measured reflections                                       | $l = -18 \rightarrow 18$           |
| 7252 independent reflections                                    | 3 standard reflections             |
| 3968 reflections with $I > 2\sigma(I)$                          | every 97 reflections               |
|                                                                 | intensity decay: none              |

## Refinement

|                               |                                                    |
|-------------------------------|----------------------------------------------------|
| Refinement on $F^2$           | $w = 1/[\sigma^2(F_o^2) + (0.0584P)^2 + 2.6079P]$  |
| $R(F) = 0.065$                | where $P = (F_o^2 + 2F_c^2)/3$                     |
| $wR(F^2) = 0.159$             | $(\Delta/\sigma)_{\text{max}} < 0.001$             |
| $S = 1.01$                    | $\Delta\rho_{\text{max}} = 0.58 \text{ e Å}^{-3}$  |
| 7252 reflections              | $\Delta\rho_{\text{min}} = -0.46 \text{ e Å}^{-3}$ |
| 523 parameters                |                                                    |
| H-atom parameters constrained |                                                    |

**Table 1**

Selected geometric parameters (Å, °).

|           |           |           |            |
|-----------|-----------|-----------|------------|
| Rh1–N1    | 2.111 (7) | N1–N2     | 1.114 (10) |
| Rh1–P2    | 2.366 (2) | N2–N3     | 1.155 (12) |
| Rh1–P1    | 2.373 (2) |           |            |
| N1–Rh1–P2 | 85.5 (2)  | N2–N1–Rh1 | 127.6 (7)  |
| N1–Rh1–P1 | 82.3 (2)  | N1–N2–N3  | 174.3 (13) |
| P2–Rh1–P1 | 96.19 (7) |           |            |

All H atoms were generated in ideal positions and refined in a riding model, with C–H distances in the range 0.93–0.96 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . Difference maps show that the methyl H atoms were not well resolved and for the final refinement they were positioned using the HFIX 33 instruction of *SHELXTL* (Sheldrick, 1997).

Data collection: *XSCANS* (Siemens, 1995); cell refinement: *XSCANS*; data reduction: *SHELXTL* (Sheldrick, 1997); program(s) used to solve structure: *SHELXTL*; program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Korean Ministry of Education (KRF-2003-015-C00309).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: OB1171). Services for accessing these data are described at the back of the journal.

## References

- Davis, B. R., Payne, N. C. & Ibers, J. A. (1969). *Inorg. Chem.* **8**, 2719–2728.
- Gan, K.-S. & Hor, T. S. A. (1995). *Ferrocenes*, edited by A. Tongni & T. Hayashi, pp. 3–104. New York: VCH.
- Lee, S. W. & Lee, J.-K. (1999). *Acta Cryst.* **C55**, 722–724.
- Ma, J.-F. & Yamamoto, Y. (1999). *J. Organomet. Chem.* **574**, 148–154.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Seok, W. K., Lee, H. N., Kim, M. Y., Klapotke, T. M., Dong, Y. & Yun, H. S. (2002). *J. Organomet. Chem.* **654**, 170–175.
- Sheldrick, G. M. (1997). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Siemens (1995). *XSCANS User's Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.