

Synthesis of 2-Amino-2,5-dideoxy-5-hydroxyphosphinyl-D-mannopyranose Derivatives: New Phospho-sugar Analogues of D-Mannosamine†

J. Chem. Research (S),
1998, 790–791†


Tadashi Hanaya,^{a*} Yasushi Fujii^b and Hiroshi Yamamoto^{a,b}

^aCenter of Instrumental Analysis, Okayama University, Tsushima, Okayama 700, Japan

^bDepartment of Chemistry, Faculty of Science, Okayama University, Tsushima, Okayama 700, Japan


3,6-Di-*O*-benzyl-5-deoxy-5-dimethylphosphinyl-1,2-*O*-isopropylidene- α -D-glucofuranose **5** was converted, in four steps, into methyl 2-acetamido-3,6-di-*O*-benzyl-2,5-dideoxy-5-dimethoxyphosphinyl- α -D-mannofuranoside **7**, which led to 3,5-di-*O*-benzyl derivatives **9** of a P-in-the-ring D-mannosamine analogue.

Various sugar analogues containing a heteroatom instead of oxygen in the ring have been prepared because of the wide interest in their chemical and biochemical properties.^{1,2} Heteroatom-in-the-ring sugar analogues of 2-amino- and 2-acetamido-2-deoxyhexopyranoses, which widely occur as a component of many natural products, have also attracted considerable interest. Thiasugar analogues of 2-acetamido-2-deoxy-D-glucose **1**³ and D-mannose **2**,⁴ for example, are potentially useful owing to their inhibitory activity in the biosynthesis of important constituents of higher animal cell walls.⁵ In view of such a chemical modification by heteroatoms, we have prepared various sugar analogues having a phosphorus atom in the ring (phospho-sugar),^{6,7} such as D-glucopyranose **3**⁸ and D-mannopyranose analogues **4**.⁹ We describe herein the first synthetic route to a new phospho-sugar of a mannosamine analogue.



Treatment of the 5-phosphinyl-D-glucofuranose derivative **5**¹⁰ with methanol in the presence of an acidic ion-exchange resin gave methyl α -D-glucofuranoside **6a** (39%) and its β -anomer **6b** (42%). The conversion of **6a** into the 2-*O*-triflate, followed by the treatment with sodium azide, afforded the corresponding 2-azido-2-deoxy- α -D-mannofuranoside; a byproduct, 5-(2-*O*-cyclo-methoxyphosphinyl)- α -D-mannofuranoside **8**, was present but it was separable only after the subsequent step (see below). The treatment of the above mixture with thioacetic acid provided, after chromatographic separation, the 2-acetamido-2-deoxy- α -D-mannofuranoside **7** (36% overall yield from **6a**) and **8** (20%). Attempts to convert the β -anomer **6b** into the corresponding 2-*C*-azide (the β -anomer of **7**) using similar procedures have remained unsuccessful.


Compound **7** was then reduced with sodium dihydrosulfide (SDMA) to give its 5-phosphino derivative, which was immediately subjected to acid hydrolysis and then to oxidation with hydrogen peroxide, affording 5-hydroxyphosphinyl-D-mannopyranose **9**. This was characterized, after peracetylation and methylation, as 2-acetamido-5-[(*R*)-methoxyphosphinyl]- α -D-mannopyranose **10a** (13% overall yield from **7**) and its 5-[(*S*)-methoxyphosphinyl] isomer **10b** (12%). A small amount of an inseparable mixture (1.7%) was obtained as a colourless



Scheme 1

syrup, which was presumed by NMR spectra to be the β -anomers of **10a** and **10b**.

The structural assignments of **10a,b** were made by the analysis of the ¹H NMR spectra. Both compounds have medium $J_{2,P}$ and $J_{4,P}$ values (14–21 Hz), indicating that they existed as a conformational mixture of ⁴C₁ and ¹C₄ forms.^{7,9} By employing the additivity rule¹¹ for vicinal coupling constants, the equilibrium populations of ⁴C₁ and ¹C₄ conformers were estimated to be 33:67 for **10a** and 42:58 for **10b**.[‡] Such equilibria were observed in the case of only the α -anomer of D-mannopyranose phospho-sugars **4**.⁹ A slight downfield shift of H-2 signal of **10b** in comparison with that of **10a** indicates an axial P=O orientation for the ¹C₄ form of **10b**. A similar downfield shift of the H-4 signal of **10a** compared with that of **10b** shows an axial P=O orientation for the ⁴C₁ form of **10a**.



Experimental

The general methods followed those described earlier,⁷ the TLC solvent system being (A) AcOEt, (B) 1:19 (v/v) EtOH–AcOEt and (C) 1:19 (v/v) AcOEt–CHCl₃. All the products were purified by

*To receive any correspondence.

†This is a **Short Paper** as defined in the Instructions for Authors, Section 5.0 [see *J. Chem. Research (S)*, 1998, Issue 1]; there is therefore no corresponding material in *J. Chem. Research (M)*.

‡The ratios of the conformers were estimated by use of the following values, $J_{1,2} = J_{2,3} = 3.8$, $J_{3,4} = 9.8$, $J_{4,5} = 11.55$ for the pure ⁴C₁ form, $J_{1,2} = 12.1$, $J_{2,3} = 2.9$, $J_{3,4} = 3.6$, $J_{4,5} = 3.5$ for the pure ¹C₄ form.

column chromatography on silica gel. NMR spectra were measured in CDCl₃ with Varian VXR-500 (500 MHz for ¹H) and VXR-200 (81 MHz for ³¹P) instruments (SC-NMR Lab., Okayama Univ.) at 23 °C. *J* values are given in Hz. The mass spectra were taken on a VG 70-SE instrument and are given as *m/z* (relative intensity).

Methyl 3,6-Di-O-benzyl-5-deoxy-5-dimethoxyphosphinyl-α-D-glucofuranoside 6a and its β-Anomer 6b.—A mixture of **5**¹⁰ (4.79 g, 9.73 mmol) and Amberlite IR-120(H⁺) in dry methanol (96 cm³) was stirred at 60 °C for 8 h. The resin was filtered off and the filtrate was evaporated *in vacuo* to give **6a** and **6b**.

6a: Colourless prisms (1.78 g, 39%), mp 84–85 °C [Found: (M⁺ + H), 467.1850. C₂₃H₃₂O₈P requires (M + 1), 467.1836]; *R*_F = 0.50 (C); δ_H 2.81 (1 H, dddd, *J*_{5,P} 20.1, *J*_{4,5} 9.8, *J*_{5,6'} 5.5, *J*_{5,6} 3.1, H-5), 2.98 (1 H, d, *J*_{2,OH} 4.9, HO-2), 3.46 (3 H, s, MeO-1), 3.67, 3.675 [3 H each, 2 d, *J*_{POMe} 10.8, P(OMe)₂], 3.90 (1 H, ddd, *J*_{6,P} 27.5, *J*_{6,6'} 9.5, H-6'), 3.95 (1 H, ddd, *J*_{6,P} 17.1, H-6), 3.98 (1 H, dd, *J*_{3,4} 4.2, *J*_{2,3} 1.8, H-3), 4.24 (1 H, td, *J*_{1,2} 4.8, H-2), 4.53, 4.59 (1 H each, 2 d, *J* 11.9, CH₂O-3 or -6), 4.55 (1 H, ddd, *J*_{4,P} 6.2, H-4), 4.54, 4.68 (1 H each, 2 d, *J* 11.0, CH₂O-6 or -3), 5.01 (1 H, d, H-1) and 7.25–7.35 (10 H, m, Ph); δ_P 32.5; FAB *m/z* 467 (M⁺ + 1, 23), 387 (6), 297 (12), 181 (9) and 91 (100).

6b: Colourless needles (1.93 g, 42%), mp 123–124 °C [Found: (M⁺ + H), 467.1828. C₂₃H₃₂O₈P requires (M + 1), 467.1836]; *R*_F = 0.40 (C); δ_H 2.83 (1 H, d, *J*_{2,OH} 4.0, HO-2), 2.90 (1 H, dddd, *J*_{5,P} 19.2, *J*_{4,5} 9.5, *J*_{5,6'} 5.5, *J*_{5,6} 3.4, H-5), 3.36 (3 H, s, MeO-1), 3.62, 3.67 [3 H each, 2 d, *J*_{POMe} 10.8, P(OMe)₂], 3.93 (1 H, ddd, *J*_{6,P} 25.6, *J*_{6,6'} 9.5, H-6'), 3.955 (1 H, ddd, *J*_{6,P} 19.5, H-6), 3.96 (1 H, dd, *J*_{3,4} 4.3, *J*_{2,3} 1.0, H-3), 4.17 (1 H, td, *J*_{1,2} 0.8, H-2), 4.53, 4.58 (1 H each, 2 d, *J* 12.2, CH₂O-3 or -6), 4.55, 4.65 (1 H each, 2 d, *J* 11.6, PhCH₂O-6 or -3), 4.65 (1 H, ddd, *J*_{4,P} 7.0, H-4), 4.80 (1 H, d, H-1) and 7.23–7.36 (10 H, m, Ph); δ_P 32.7; FAB *m/z* (M⁺ + 1, 24), 435 (15), 297 (39), 207 (9) and 91 (100).

Methyl 2-Acetamido-3,6-di-O-benzyl-2,5-dideoxy-5-dimethoxyphosphinyl-α-D-glucofuranoside 7 and Methyl 3-O-Benzyl-5-deoxy-5-[(5*R*,*S*)-2-O-cyclomethoxyphosphinyl]-α-D-mannofuranoside 8.—Trifluoromethanesulfonic anhydride (0.54 cm³, 3.21 mmol) was added to a solution of **6a** (750 mg, 1.61 mmol) in pyridine (0.52 cm³, 6.43 mmol) in dry CH₂Cl₂ (25 cm³) at –40 °C. The mixture was stirred at –40 °C for 30 min under argon, poured into water and extracted with CH₂Cl₂. The organic layer was evaporated *in vacuo* to give the crude 2-O-triflate as a colourless syrup [*R*_F = 0.63 (A)]. A mixture of this syrup and sodium azide (520 mg, 8.00 mmol) in dry DMF (25 cm³) was stirred at 40 °C for 3 d under argon, poured into water and extracted with CHCl₃. The organic layer was evaporated *in vacuo*, giving an inseparable mixture (573 mg) of the 2-azido compound and **8**. This was dissolved in thioacetic acid (5.0 cm³, 7.0 mmol) and stirred at 23 °C for 1 d under argon. The excess thioacetic acid was evaporated *in vacuo*, giving **7** and **8**.

7: Colourless prisms (295 mg, 36% from **6a**), mp 42–44 °C [Found: (M⁺ + H), 508.2118. C₂₅H₃₅O₈NP requires (M + 1), 508.2102]; *R*_F = 0.18 (A); δ_H 1.72 (3 H, s, Ac), 2.92 (1 H, dtd, *J*_{5,P} 21.7, *J*_{5,6} 6.7, *J*_{4,5} 6.1, *J*_{5,6'} 4.3, H-5), 3.30 (3 H, s, MeO-1), 3.62, 3.70 [3 H each, 2 d, *J*_{POMe} 10.7, P(OMe)₂], 3.81 (1 H, ddd, *J*_{6,P} 16.5, *J*_{6,6'} 10.1, H-6'), 3.98 (1 H, td, *J*_{6,P} 10.1, H-6), 4.41 (1 H, dd, *J*_{3,4} 6.5, *J*_{2,3} 6.0, H-3), 4.43, 4.52 (1 H each, 2 d, *J* 11.3, CH₂O-3 or -6), 4.44, 4.52 (1 H each, 2 d, *J* 11.6, CH₂O-6 or -3), 4.54 (1 H, ddd, *J*_{2,NH} 8.9, *J*_{1,2} 1.2, H-2), 4.60 (1 H, td, *J*_{4,P} 18.0, H-4), 4.76 (1 H, d, H-1), 7.04 (1 H, br d, HN-2) and 7.25–7.35 (10 H, m, Ph); δ_P 32.1; FAB *m/z* 508 (M⁺ + 1, 15), 476 (14), 368 (9), 338 (38), 248 (10) and 91 (100).

8: Yellow syrup (143 mg, 20%) [Found: (M⁺ + H), 435.1560. C₂₂H₂₈O₇P requires (M + 1), 435.1573]; *R*_F = 0.55 (A); (for the main isomer) δ_H 2.79 (1 H, dddd, *J*_{5,P} 19.5, *J*_{5,6} 9.2, *J*_{5,6'} 6.1, *J*_{4,5} 3.0, H-5), 3.50 (3 H, s, MeO-1), 3.78 [3 H, d, *J*_{POMe} 12.0, P(OMe)], 3.88 (1 H, ddd, *J*_{6,6'} 10.4, *J*_{6,P} 7.9, H-6'), 3.92 (1 H, td, *J*_{6,P} 11.0, H-6), 4.29 (1 H, ddd, *J*_{3,4} 6.1, *J*_{2,3} 3.1, *J*_{3,P} 1.2, H-3), 4.41, 4.54 (1 H each, 2 d, *J* 12.2, CH₂O-3 or -6), 4.49, 4.77 (1 H each, 2 d, *J* 11.6, CH₂O-6 or -3), 4.51 (1 H, dd, *J*_{2,P} 18.9, *J*_{1,2} ≈ 0, H-2), 4.71 (1 H, ddd, *J*_{4,P} 31.1, H-4), 5.11 (1 H, br s, H-1) and 7.23–7.38 (10 H, m, Ph); δ_P 22.6; FAB *m/z* 435 (M⁺ + 1, 16), 345 (5), 237 (5), 181 (10) and 91 (100).

2-Acetamido-1,4-di-O-acetyl-3,6-di-O-benzyl-2,5-dideoxy-5-[(*R*)-methoxyphosphinyl]-α-D-mannopyranose 10a and its 5-[(*S*)-Methoxyphosphinyl] Isomer 10b.—To a solution of **7** (157 mg, 0.309 mmol) in dry toluene (1.5 cm³) was added SDMA (0.34 M in toluene, 3.20 cm³, 1.09 mmol) at 0 °C under argon. The mixture was stirred at 0 °C for 1 h and then water (0.12 cm³) was added. The mixture

was centrifuged and the precipitate was extracted with benzene. The organic layer was evaporated *in vacuo*, giving the 5-phosphino derivative as a colourless syrup; *R*_F = 0.70 (A). This syrup was immediately treated with 1:1 (v/v) propan-2-ol–0.5 M HCl (3.0 cm³) at 90 °C for 2 h under argon. After cooling, 30% hydrogen peroxide (1.0 cm³) was added. The solution was stirred at 23 °C for 18 h and concentrated *in vacuo*. The residue was dissolved in MeOH (1.5 cm³), treated with propylene oxide (0.8 ml) at 23 °C for 6 h and evaporated *in vacuo* to give crude **9** as a colourless syrup. This was dissolved in dry pyridine (1.5 cm³) and acetic anhydride (0.8 cm³) at 23 °C for 20 h, and concentrated *in vacuo*. The residue was dissolved in ethanol and passed through a column of Amberlite IR-120(H⁺). The eluent was evaporated *in vacuo* and the residue was methylated with ethereal diazomethane in dry CH₂Cl₂ (2.0 cm³) at 0 °C. After evaporation of the solvent, the residue was separated by chromatography into three fractions A–C.

Fraction A [*R*_F = 0.27 (B)] gave **10a** as colourless needles, mp 168–169 °C (21.3 mg, 13% from **7**) [Found: (M⁺ + 1), 548.2062. C₂₇H₃₄O₉NP requires (M + 1), 548.2051]; δ_H 1.73, 2.03, 1.78 (3 H each, 3 s, Ac-1,2,4), 2.66 (1 H, dtd, *J*_{5,P} 19.2, *J*_{5,6'} 9.5, *J*_{4,5} 6.1, *J*_{5,6} 5.8, H-5), 3.73 (1 H, dd, *J*_{3,4} 5.8, *J*_{2,3} 3.1, H-3), 3.81 (3 H, d, *J*_{POMe} 11.0, P(OMe)), 3.87 (1 H, q, *J*_{6,6'} 9.8, *J*_{6,P} 9.4, H-6'), 3.92 (1 H, dt, *J*_{6,P} 6.4, H-6), 4.45, 4.54 (1 H each, 2 d, *J* 11.9, CH₂O-3 or -6), 4.51, 4.57 (1 H each, 2 d, *J* 11.9, CH₂O-6 or -3), 4.79 (1 H, dtd, *J*_{2,P} 14.7, *J*_{2,NH} 9.8, *J*_{1,2} 9.4, H-2), 5.39 (1 H, dd, *J*_{1,P} 5.3, H-1), 5.63 (1 H, dt, *J*_{4,P} 21.4, H-4), 5.92 (1 H, br d, NH-2), 7.25–7.36 (10 H, m, Ph); δ_P 41.4; FAB *m/z* 548 (M⁺ + 1, 22), 506 (8), 488 (10), 181 (14) and 91 (100).

Fraction B [*R*_F = 0.22 (B)] gave an inseparable mixture (1:1) of (5-[(*R*)- and (*S*)-methoxyphosphinyl]-β-isomers (3.2 mg, 1.8%) containing a small amount of **10a,b**.

Fraction C [*R*_F = 0.18 (B)] gave **10b** as a colourless syrup (20.2 mg, 12% from **7**); δ_H 1.81, 2.03, 2.13 (3 H each, 3 s, Ac-1,2,4), 2.70 (1 H, dtd, *J*_{5,P} 18.9, *J*_{4,5} 7.1, *J*_{5,6} 6.9, *J*_{5,6'} 5.2, H-5), 3.72 (1 H, ddd, *J*_{6,P} 9.8, H-6'), 3.73 (1 H, dd, *J*_{3,4} 6.0, *J*_{2,3} 3.4 Hz, H-3), 3.74 (3 H, d, *J*_{POMe} 10.7, P(OMe)), 3.84 (1 H, td, *J*_{6,P} 9.6, H-6), 4.49 (2 H, s, CH₂O-6), 4.50, 4.61 (1 H each, 2 d, *J* 11.9, CH₂O-3), 4.84 (1 H, dtd, *J*_{2,P} 14.3, *J*_{2,NH} 9.2, *J*_{1,2} 8.7, H-2), 5.32 (1 H, dd, *J*_{1,P} 7.9, H-1), 5.54 (1 H, ddd, *J*_{4,P} 16.9, H-4), 5.63 (1 H, br d, NH-2), 7.26–7.36 (10 H, m, Ph); δ_P 41.3.

Received, 29th June 1998; Accepted, 10th August 1998

Paper E/8/04961K

References

- 1 H. Paulsen, *Angew. Chem., Int. Ed. Engl.*, 1966, **5**, 495; G. Legler and E. Jülich, *Carbohydr. Res.*, 1984, **108**, 61.
- 2 R. L. Whistler and W. C. Lake, *Methods Carbohydr. Chem.*, 1972, **6**, 286; M. J. Pitts, M. Chmielewski, M. S. Chen, M. M. A. Abd El-rahman and R. L. Whistler, *Arch. Biochem. Biophys.*, 1975, **169**, 384.
- 3 R. Bognar, P. Herczegh, R. L. Whistler and E. B. Madumelu, *Carbohydr. Res.*, 1981, **70**, 138; R. D. Guthrie and K. O'Shea, *Aust. J. Chem.*, 1981, **34**, 2225; E. Tanahashi, M. Kiso and A. Hasegawa, *Carbohydr. Res.*, 1983, **97**, 304.
- 4 A. Hasegawa, E. Tanahashi, Y. Hioki and M. Kiso, *Carbohydr. Res.*, 1983, **102**, 168; R. Csuk and B. I. Glänzer, *J. Chem. Soc., Chem. Commun.*, 1986, 343.
- 5 P. K. Ray, *Adv. Appl. Microbiol.*, 1971, **21**, 227.
- 6 T. Hanaya and H. Yamamoto, *Yuki Gosei Kagaku Kyokai Shi*, 1993, **51**, 377 and references therein.
- 7 T. Hanaya, K. Yasuda, H. Yamamoto and H. Yamamoto, *Bull. Chem. Soc. Jpn.*, 1993, **66**, 2315.
- 8 H. Yamamoto, T. Hanaya, H. Kawamoto, S. Inokawa, M. Yamashita, M.-A. Armour and T. T. Nakashima, *J. Org. Chem.*, 1985, **50**, 3516; T. Hanaya, A. Akamatsu, H. Kawamoto, M.-A. Armour, A. M. Hogg and H. Yamamoto, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 2398.
- 9 T. Hanaya, K. Hirose and H. Yamamoto, *Heterocycles*, 1993, **36**, 2557; T. Hanaya, K. Ohmori, H. Yamamoto, M.-A. Armour and A. M. Hogg, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 1174.
- 10 T. Hanaya, Y. Fujii, S. Ikejiri and H. Yamamoto, *Heterocycles*, 1999, **50**, in press.
- 11 C. Altona and C. A. G. Haasnoot, *Org. Magn. Reson.*, 1980, **13**, 417; T. Hanaya and H. Yamamoto, *Carbohydr. Res.*, in press.